


**COMMONWEALTH OF VIRGINIA  
DEPARTMENT OF ENVIRONMENTAL QUALITY  
WATER PLANNING DIVISION  
JUTTA SCHNEIDER, DIRECTOR**

**P.O. BOX 1105**

**Richmond, VA 23218**

**Subject:** Guidance Memo No. GM23-2002  
**2024 Water Quality Assessment Guidance Manual**

**To:** Regional Directors

**From:** Jutta Schneider, Water Planning Division Director 

**Date:** 5/30/2023

**Copies:** Water Quality Planning Managers and Staff, Water Permitting Division Director, Water Division Director, Director of Central Operations, Director of Regional Operations

**Summary:**

The Environmental Protection Agency's (EPA) 2006 Integrated Report Guidance recommends that states submit an "Integrated Report" (IR) that will satisfy Clean Water Act (CWA) requirements for both Sections 305(b) water quality reports and 303(d) impaired waters lists. According to EPA this Integrated Report should include the following information:

- delineation of water quality assessment units (AUs) based on the National Hydrography Dataset (NHD);
- status of and progress toward achieving comprehensive assessments of all waters;
- Water Quality Standard attainment determination for every AU;
- additional monitoring that may be needed to determine Water Quality Standard attainment status and, if necessary, to support development of Total Maximum Daily Loads (TMDLs) for each pollutant/AU combination;
- schedules for additional monitoring planned for AUs;
- pollutant/AU combinations still requiring TMDLs;
- TMDL development schedules reflecting the priority ranking of each pollutant/AU combination.

DEQ has incorporated the EPA Integrated Reporting guidance into the Virginia 2024 Water Quality Assessment Guidance Manual. The 2024 IR guidance is designed to integrate or combine the 305(b) overall assessment of Virginia's waters and separate out those waters impaired and needing a TMDL as per Section 303(d) of the Clean Water Act. The EPA Integrated Report Guidance and the Assessment, Total Maximum Daily Load (TMDL) Tracking and Implementation System (ATTAINS V1.0.6) have five different assessment categories in which every segment or "assessment unit" (AU) will be placed. The EPA Guidance allows the states to subdivide the federal categories in order to address state programmatic needs. Virginia's 2024 IR guidance contains the categories and subcategories Virginia has chosen for enhanced tracking and data management purposes.

The 2024 water quality assessment guidance manual contains a number of changes, all enumerated in Part II. A few notable updates include an assessment methodology for assessing benthic chlorophyll-a in parts of the North and South Forks and mainstem of the Shenandoah River, the addition of Category 3E to categorize stressors identified in TMDL development that do not have water quality standards and a streamlined data submittal process for volunteer and non-agency data using the new Virginia Data Explorer database.

The data window to be used in the development of the 2024 IR is generally January 1, 2017 through December 31, 2022. A Notice of Public Solicitation of Water Quality Data for the 2024 IR was published in the Virginia Register on January 16, 2023, asking for all data to be reviewed and considered be submitted by March 6, 2023. Water quality data received after this date will be reviewed for the 2026 IR. The manual uses excerpts from the “EPA 2006 Integrated Report Guidance”, “2008, 2010, 2012, 2014, 2016, 2018 and 2024 EPA Integrated Report Clarification Guidance Memoranda”, and “EPA 1997 Guidelines for the Preparation of the 1998 State Water Quality Assessment 305(b) Reports”, along with other state and federal guidelines. The assessment methodologies in this draft guidance were developed based on final triennial review amendments to the water quality standards regulation approved by EPA on April 18, 2023.

### **Electronic Copy:**

An electronic copy of this guidance in PDF format is available for staff internally on the Water Quality Assessments SharePoint site, and for the general public on DEQ's website at:  
<https://www.deq.virginia.gov/water/water-quality/assessments/wqa-guidance-manual>.

### **Contact information:**

If you have any questions regarding the guidance manual, you can contact Amanda Shaver, Department of Environmental Quality, P.O. Box 1105, Richmond, Virginia 23218. Telephone (804) 774-8416 or via e-mail [amanda.shaver@deq.virginia.gov](mailto:amanda.shaver@deq.virginia.gov).

### **Certification:**

As required by Subsection B of § 2.2-4002.1 of the APA, the agency certifies that this guidance document conforms to the definition of a guidance document in § 2.2-4101 of the Code of Virginia.

### **Disclaimer:**

**This document has been developed based on Virginia’s Water Quality Standards Regulation (9 VAC 25-260), with amendments approved by the State Water Control Board resulting from iterative Triennial Reviews or periodic rulemakings. It is provided as guidance and, as such, sets forth standard operating procedures for the agency. However, it does not mandate nor prohibit any particular method for the analysis of data, establishment of a wasteload allocation, or establishment of a permit limit. If alternative proposals are made, such proposals should be reviewed and accepted or denied based on their technical adequacy and compliance with appropriate laws and regulations.**



WATER QUALITY  
ASSESSMENT GUIDANCE MANUAL  
for  
2024

305(b)/303(d) Integrated Water Quality Report

May 2023

## TABLE OF CONTENTS

<u>PART</u>		<u>PAGE</u>
<b>I.</b>	<b>REGULATORY REQUIREMENTS AND OBJECTIVES.....</b>	<b><u>1</u></b>
<b>II.</b>	<b>MODIFICATIONS TO PREVIOUS GUIDANCE.....</b>	<b><u>3</u></b>
<b>III.</b>	<b>FEDERAL AND VIRGINIA ASSESSMENT CATEGORIES.....</b>	<b><u>4</u></b>
<b>IV.</b>	<b>GENERAL RULES OF WATER QUALITY ASSESSMENT.....</b>	<b><u>6</u></b>
<b>V.</b>	<b>ASSESSMENT METHODOLOGY</b>	
	5.1 Monitoring Station Siting and Delineation.....	<u>12</u>
	5.2 Evaluation of Designated Uses.....	<u>13</u>
	5.3 Chesapeake Bay Assessment.....	<u>22</u>
	5.4 Biological Assessments.....	<u>27</u>
	5.5 Toxics Assessment.....	<u>30</u>
	5.6 Swamp Waters Assessment.....	<u>33</u>
	5.7 Lakes/Reservoirs Assessment.....	<u>33</u>
	5.8 Nonpoint Source Assessment.....	<u>42</u>
	5.9 Coastal Assessment.....	<u>42</u>
	5.10 Wetlands Assessment.....	<u>42</u>
	5.11 Freshwater Probabilistic Monitoring Assessment.....	<u>52</u>
	5.12 Continuous Monitoring Assessment.....	<u>52</u>
<b>VI.</b>	<b>PROCEDURES FOR NON-AGENCY DATA.....</b>	<b><u>55</u></b>
<b>VII.</b>	<b>303(D) LISTING/DELISTING AND TMDL PRIORITY RANKING.....</b>	<b><u>60</u></b>
<b>VIII.</b>	<b>APPENDICES</b>	
	A – Clean Water Act References.....	<u>65</u>
	B – Regional Biologist Assessment Fact Sheet.....	<u>67</u>
	C – Classification of Virginia’s Shellfish Growing Areas.....	<u>69</u>
	D-1– Incorporating the Proactive Approach for Impaired Waters Delisting.....	<u>73</u>
	D-2- Category 5R Requirements.....	<u>75</u>
	E-1 – Fish Tissue Values (TVs).....	<u>76</u>
	E-2 – Fish Tissue Screening Values (TSVs).....	<u>79</u>
	F – Consensus-Based and ER-M Sediment Screen Values.....	<u>81</u>
	G – Estuarine Aquatic Life Use Assessment using the Sediment Quality Triad.....	<u>84</u>
	H – Significant Lakes by DEQ Regional Offices.....	<u>103</u>
	I – Memo to Standardize Rounding Method for Assessment.....	<u>106</u>

## **List of Tables**

Table 1	Designated Use Matrix	P. 19
Table 2	Designated Use Assessment Methodology	P. 20
Table 3	Trophic Status in ATTAINS	P. 39
Table 4	Lake/Reservoir Designated Use Assessment Summary	P. 41
Table 5	Wetland Monitoring and Assessment Program Elements	P. 47

## **List of Figures**

Figure 1	Annual temperature cycles in stratified lakes	P. 35
Figure 2	Multi-tiered sampling design of wetlands	P. 47
Figure 3	Wetland Water Quality Stress Condition	P. 49
Figure 4	Comparison of Wetland Water Quality Condition Scores	P. 50
Figure 5	Comparison of In-Stream Water Quality Parameters	P. 50
Figure 6	Non-tidal Wetlands Data Viewer	P. 51

## **PART I. REGULATORY REQUIREMENTS AND OBJECTIVES**

Under the [Clean Water Act](#), the United States Environmental Protection Agency (EPA) requires that each state develop a program to monitor the quality of its surface and ground waters and prepare a report every two years describing the status of its water quality. Each state identifies waters of concern and schedules additional monitoring, if appropriate, to determine if designated uses are being met. EPA issues guidelines for States to use during the reporting cycle for national consistency purposes. States are encouraged to use these guidelines to prepare these water quality reports for EPA. EPA compiles the data from the state reports, summarizes them, and transmits the summaries to Congress, including an analysis of water quality nationwide. The 305(b)/303(d) integrated process is the principal means by which the EPA, Congress, and the public evaluate current water quality, the progress made maintaining and restoring water quality, and the extent of remaining work to be done. Many states, including Virginia, rely on the 305(b)/303(d) process for information needed to conduct water quality planning. The 305(b)/303(d) process is an integral part of Virginia's water quality management program, requirements for which are set forth in [40 CFR 130](#). The Department of Environmental Quality (DEQ) and the Department of Conservation and Recreation (DCR) are the principal state agencies charged with conducting water quality assessment and associated activities.

In 1997, the General Assembly enacted the Water Quality Monitoring, Information and Restoration Act (WQMIRA; VA Code [§62.1- 44.19:4 through §62.1- 44.19:8](#)). This legislation supplements the CWA 305(b)/303(d) federal requirements. The requirements of this State legislation for assessment procedures or processes are briefly outlined as follows:

1. The Act requires the 303(d) portion of the Integrated Report to identify geographically defined water segments as impaired if monitoring or other evidence shows:
  - a. exceedances of ambient [water quality standards](#) for aquatic life or human health;
  - b. fishing restrictions or advisories;
  - c. shellfish consumption restrictions due to contamination;
  - d. nutrient over-enrichment;
  - e. significant declines in aquatic life biodiversity or populations; and/or
  - f. contamination of sediments at levels which exceed water quality standards or threaten aquatic life or human health.
2. Waters identified as “naturally impaired”, “fully supporting but threatened” or “evaluated” (without monitoring) as impaired shall be set out in the 303(d) portion of the Integrated Report in the same format as those listed as “impaired.”
3. The 303(d) portion of the Integrated Report shall include an assessment, conducted in conjunction with other appropriate state agencies, for the attribution of impairment to point and nonpoint sources. The absence of point source permit violations at or near the impaired water shall not conclusively support a determination that impairment is due to nonpoint sources. In determining the cause for impairment, the Board shall consider the cumulative impact of 1) multiple point source discharges, 2) individual discharges over time, and 3) nonpoint sources.
4. The Board shall develop and publish a procedure governing its process for defining and determining impaired water segments and shall provide for public comment on the procedure.

5. The Integrated Report, inclusive of CWA sections 305(b) and 303(d) shall be produced in accordance with the schedule required by federal law and shall incorporate at least the preceding five years of data, where appropriate. Data older than five years shall be incorporated when scientifically appropriate for trend analysis or other longer-term considerations.
6. The Integrated Report, inclusive of CWA sections 305(b) and 303(d), shall be developed in consultation with scientists from state universities prior to submission by the Board to EPA.
7. The Integrated Report, inclusive of CWA sections 305(b) and 303(d), shall indicate water quality trends for specific, easily identifiable, geographically defined water segments and provide summaries of the trends using available data and evaluations. This will allow the citizens of the Commonwealth to easily interpret and understand the conditions of the geographically defined water segments.
8. Based on the information in the Integrated Report, inclusive of CWA sections 303(d) and 305(b), the Board shall request the Department of Wildlife Resources (DWR) or the Virginia Marine Resources Commission (VMRC) to post notices at public access points for all “toxic” impaired waters. The notice, prepared by the Board, shall contain the basis for the impaired designation and a statement of potential health risks. The Board shall coordinate with the DWR and VMRC to assure that adequate notice of posted waters is provided to those purchasing hunting and fishing licenses.

The Water Quality Monitoring, Information, and Restoration Act directs DEQ to develop and publish a water quality assessment guidance document governing the process for defining and determining impaired waters, and to provide an opportunity for public comment on the assessment guidance.

The purpose of this guidance manual is to guide DEQ staff in the development and reporting of the 2024 Integrated Report (305(b) Water Quality Assessment/303(d) Impaired Waters). It is also intended to assist the public in understanding the monitoring and assessment process.

Section 305(b) of the Clean Water Act requires each state to submit a biennial report to EPA describing the quality of its navigable waters. The 305(b) report provides DEQ’s best overall assessment of water quality conditions and trends in the Commonwealth. The report is intended to be used as a tool in planning and management of water quality in Virginia. The report also directs continuous planning and implementation activities in coordination with the State Water Quality Management Plan and the Continuous Planning Process (CPP).

Primary objectives of the Integrated Report are:

1. To educate and inform citizens and public officials about Virginia’s overall water quality.
2. To analyze water quality data in order to determine the extent to which Virginia’s waters are supporting the applicable designated uses and to compare the results to WQ Standards and other appropriate criteria and guidelines.
3. To determine the causes for the “failure to support” the designated uses of the State’s waters.
4. To determine the nature and recognizable extent of point and nonpoint source impacts in accordance with state and federal guidelines.

Section 303(d) of the Clean Water Act and the Environmental Protection Agency’s regulation [40 CFR Section 130.7 \(d\)](#), promulgated in July 1992, requires each state to submit a Total Maximum Daily Load (TMDL) Priority List to EPA on April 1 of even numbered years.

Category 5 signifies waters that are impaired and need a TMDL.

Impaired waters needing a TMDL are those waters that do not meet water quality standards due to a pollutant(s). A pollutant, as defined in [40 CFR 122.2](#), means: *any dredged spoil, solid waste, incinerator residue, filter backwash, sewage, garbage, sewage sludge, munitions, chemical wastes, biological materials, radioactive materials (except those regulated under the Atomic Energy Act of 1954, as amended (42 U.S.C. 2011 et seq.)), heat, wrecked or discarded equipment, rock, sand, cellar dirt and industrial, municipal, and agricultural waste discharged into water.*

EPA’s Integrated Report Guidance recommends that states submit an “Integrated Report” that will satisfy Clean Water Act (CWA) requirements for Sections 305(b) overall water quality report, 303(d) Impaired Waters List and Section 314 assessment of publicly owned lakes. This Integrated Report shows the following information:

- delineation of water quality assessment units (AUs) based on [National Hydrography Dataset \(NHD\)](#);
- status of and progress toward achieving comprehensive assessments of all waters;
- attainment status of water quality standards (WQS) for every AU assessed;
- additional monitoring that may be needed to determine WQS attainment status and, if necessary, to support development of TMDLs for each pollutant/AU combination;
- schedules for additional monitoring planned for AUs;
- pollutant/AU combinations still requiring TMDLs; and
- TMDL development schedules reflecting the priority ranking of each pollutant/AU combination.

Virginia’s biennial water quality assessment is conducted by the Department of Environmental Quality (DEQ), with the assistance of the Department of Conservation and Recreation (DCR) and the Virginia Department of Health (VDH), to determine the water quality conditions in the Commonwealth. The results of this water quality analysis are usually reported to the EPA by April 1 of even numbered years. The Integrated Report describes the aggregated water quality conditions of the State and contains the individual listing of those waters identified as “impaired” for one or more designated uses and needing a Total Maximum Daily Load (TMDL). As per EPA guidance, the former 305(b) Water Quality Assessment Report and the 303(d) Impaired Waters List are now combined into a single Integrated Report. EPA compiles the data from all State reports into a national water quality status report that is presented to Congress.

## **PART II. MODIFICATIONS TO PREVIOUS GUIDANCE**

DEQ has incorporated EPA Integrated Reporting guidance initially developed in 2004, with all subsequent versions current to January 2023. DEQ’s guidance for the 2024 Integrated Report contains the following notable modifications:

1. Assessment methodology for assessing benthic chlorophyll-a in parts of the North and South Forks and mainstem of the Shenandoah River.
2. The addition of Virginia Category 3E to categorize parameters identified as most probable stressors, had a TMDL equation assigned in an EPA approved TMDL study, but do not have water quality standards.



3. Implementing a streamlined data submittal process for volunteer and non-agency data using the new Virginia Data Explorer database.
4. The recreational bacteria assessment will be based on data collected in the two most recent calendar years of the assessment data window (e.g. 2021-2022).

### **PART III. FEDERAL AND VIRGINIA ASSESSMENT CATEGORIES**

The 2024 EPA Integrated Report Guidance and the Assessment, Total Maximum Daily Load (TMDL) Tracking and Implementation System (ATTAINS V1.0.6) have five major categories and three subcategories which every “assessment unit” (AU) are placed based on designated use attainment. Additionally, Virginia has created several subcategories to supplement the federal categories, enabling a more precise water quality tracking and reporting mechanism.

Below are the US EPA-defined categories and associated Virginia-defined subcategories:

#### ***FULLY SUPPORTING - Waters are supporting one or more designated uses***

- **EPA Category 1** - Attaining all associated designated uses and no designated use is threatened
  - Va. Category 1A** - waters are attaining all uses and a TMDL has been developed for one or more uses.
- **EPA Category 2** – Available data and/or other information indicate that some, but not all of the designated uses are supported.
  - Va. Category 2A** - waters are supporting all of the uses for which they are monitored.
  - Va. Category 2B** - the Shellfish Use is fully supporting, but harvesting is limited because of seasonal or conditional closures per VDH Department of Shellfish Sanitation.
  - Va. Category 2C** - waters are now attaining the use(s) for which they were originally 303(d) listed and the TMDL is EPA approved but other applicable use(s) were not monitored and assessed.

#### ***INDETERMINATE - Waters needing additional information***

- **EPA Category 3** - Insufficient data and/or information to determine whether any designated uses are met.
  - Va. Category 3A** - no data are available within the data window of the current assessment to determine if any designated use is attained and the water is not currently listed as impaired.
  - Va. Category 3B** - some data exist but are insufficient to determine support of designated uses. Such waters will be prioritized for follow up monitoring, as needed.
  - Va. Category 3C**- data collected by a citizen monitoring or another organization indicating water quality problems may exist but the methodology and/or data quality has not been approved for a determination of support of designated use(s). These waters are considered as having insufficient data and are prioritized by DEQ for follow up monitoring.

**Va. Category 3D** - data collected by a citizen monitoring or other organization indicating designated use(s) are being attained but the methodology and/or data quality has not been approved for such a determination.

**Va. Category 3E** – waters are of concern to the state and data exist, but no Water Quality Standard has been established for the pollutant. The waterbody either exceeds a state screening value, failed a toxicity test or the pollutant has been identified as a most probable stressor and had a TMDL equation assigned during development of an EPA approved TMDL.

***IMPAIRED - Waters are impaired or threatened but a TMDL is not required.***

- **EPA Category 4A** – water is impaired or threatened for one or more designated uses but does not require a TMDL (an EPA approved TMDL already exists or the waterbody has been nested within an approved TMDL). In the case of a nested water, a new TMDL is not necessary to address the newly impaired water if the nesting procedure is followed (see Part VII, Rule 3).
- **EPA Category 4B** - water is impaired or threatened for one or more designated uses but does not require the development of a TMDL because other pollution control requirements are reasonably expected to result in attainment of the Water Quality Standard by the next reporting period-
- **EPA Category 4C** - water is impaired or threatened for one or more designated uses but does not require a TMDL because the impairment is not caused by a pollutant. This category includes Virginia waters that are swampwaters awaiting aquatic life criteria reclassification because the impairment has been determined to be caused by natural conditions.

**Va. Category 4D** – part(s) of a water quality standard is attained for a pollutant with a TMDL, but the remaining criteria for the standard were not assessed due to insufficient information. (Only to be applied to dissolved oxygen in tidal waters of the Chesapeake Bay).

***IMPAIRED - Waters are impaired or threatened and require a TMDL***

- **EPA Category 5** - Waters are impaired or threatened and a TMDL is needed.

**Va. Category 5A** - a Water Quality Standard is not attained. The water is impaired or threatened for one or more designated uses by a pollutant(s) and requires a TMDL (303d list).

**Va. Category 5B** - the Water Quality Standard for shellfish use is not attained. One or more pollutants causing impairment require TMDL development (303d list).

**Va. Category 5C** - the Water Quality Standard is not attained due to “suspected” natural conditions. The water is impaired for one or more designated uses by a pollutant(s) and may require a TMDL (303d list). Water Quality Standards for these waters may be re-evaluated due to the presence of natural conditions.

**Va. Category 5D** - the Water Quality Standard is not attained where TMDLs for a pollutant(s) have been developed and one or more pollutants are still causing impairment requiring additional TMDL development (303d list).

**Va. Category 5F** - the Water Quality Standard is attained for a pollutant(s) with a TMDL and 303(d) delisting has been approved but the water remains impaired for additional pollutant(s) requiring TMDL development (303d list).

**Va. Category 5R** - the Water Quality Standard is not attained and the water is impaired, and implementation of an EPA-accepted advance restoration plan is expected to result in attainment (303d list). A status update will be provided each 303(d) cycle to evaluate progress.

- **EPA Category 5M** – the Water Quality Standard is not attained for mercury primarily due to atmospheric deposition (303d list).

## **PART IV. GENERAL RULES OF WATER QUALITY ASSESSMENT**

305(b)/303(d) assessments seek to characterize surface waters under typical, ambient conditions. For this reason, water quality assessments are based on data that are representative of normal conditions. The assessment begins by analyzing QA/QC-approved data from DEQ ambient water quality monitoring stations, biological, sediment and fish tissue monitoring, special studies and/or other non-DEQ water quality data collected during the six-year assessment period<sup>1</sup>. This interval of time works in concert with the ambient rotating watershed monitoring program. Assessment data are compared to both numeric and narrative criteria established for Virginia’s designated uses and promulgated in its [water quality standards](#) (WQS; 9 VAC 25-260). DEQ does not interpret the impacts of a single pollution event (e.g., unusually intense wet-weather, incident response) as representative of ambient conditions if it is believed that the problem has been addressed through a permitting action or other control measure. Listing decisions will not be based on datasets that are solely targeted or biased<sup>2</sup>.

The following list of rules is to be applied uniformly, only to be modified after internal review or directive from EPA. Specific assessment procedures are outlined later in this document.

### Rule 1

Impaired waters are defined as those with exceedances of recurring or human health-related water quality standards as documented by quality assured monitoring data. Predictive data generally refers to computer-generated modeling data and may be used for assessment purposes on a case-by-case basis. Impairments are generally determined from exceedances of the numeric/narrative water quality standards, using the guidelines described in Part V of this manual.

Previous EPA guidance allows for the use of an exceedance rate of >10.5% of the total samples analyzed to establish impairment using conventional parameters (i.e., dissolved oxygen, pH and temperature.) This “allowable” exceedance rate takes into account equipment failure and/or human error. Single samples (n = 1) will be considered insufficient information for assessment. A single exceedance of the WQS for conventional parameters is also considered insufficient justification for

---

<sup>1</sup> Unless otherwise noted in Part V. Assessment Methodology.

<sup>2</sup> DEQ may assess targeted datasets collected to investigate probable stressors for existing benthic impairments (i.e. toxics). This may lead to the identification of other assessment units with impaired aquatic life uses. These impairments may be addressed under the TMDL(s) developed to address the initial benthic impairment.

303(d) listing. At least two exceedances and > 10.5% of the total samples are required for a water to be listed as impaired.

## Rule 2

Waters where restrictions are placed on the shellfishing and fish consumption uses by the Virginia Department of Health (VDH) are determined to not be meeting these designated uses ([9 VAC 25-260-10 A.](#)) and are listed as impaired, unless the designated use has been administratively removed. Uses are administratively removed in the presence of a permitted discharge outfall and any associated VDH safety zone, where the salinity regimes are not conducive for productive harvest, or where a consumption advisory that does not restrict the designated use has been issued.

## Rule 3

*Escherichia coli* (freshwater) and enterococci (saltwater and transition zone) data will be assessed for the recreation designated use. These indicators replaced fecal coliform bacteria in 2006. Any waters previously listed for fecal coliform will remain as impaired until appropriate bacteria data are available and assessed.

Virginia adopted EPA's 2012 nationally recommended primary contact recreational water quality criteria in 2019. The criteria are comprised of a Statistical Threshold Value (STV), which can be exceeded no more than 10% of the time over any period up to 90 days, and a geometric mean value, which cannot be exceeded over any period up to 90 days. For determining primary contact recreation use support, a 90-day assessment period should be used. These periods will be assessed on a rolling basis (i.e., determined from each sample date and adding 89 days.) Each 90-day period that is assessed should be evaluated using a dataset containing at least one sample that is not used in a preceding or subsequent 90-day period. The STV is 410 counts per 100 ml for *E. coli* in freshwater and 130 counts per 100 ml for enterococci in saltwater and transition zone. The geometric mean criteria is 126 counts per 100 ml for *E. coli* in freshwater and 35 counts per 100 ml for enterococci in saltwater and transition zone. See [9 VAC 25-260-140-C](#) for freshwater, saltwater, and transition zone delineation. However, water quality data attributed to an unusually intense wet-weather event such as a hurricane or tropical storm may be excluded from geomean calculation because of the technical, science-based rationale that these data are not representative of normal conditions or that they don't represent conditions used to develop the criteria.

For each assessment unit, primary contact recreation use assessments are based on bacteria monitoring data collected during the two most recent calendar years of the six-year assessment window. For the 2024 IR cycle this would mean only bacteria data collected between January 1, 2021 and December 31, 2022 will be assessed for recreation use support. The focus on the most recent data collected aligns with the DEQ bacteria monitoring strategy as outlined in the [Water Quality Monitoring Strategy](#), updated in February 2022. The strategy outlines a high frequency monitoring network of approximately 100 sites over a two-year period prioritizing primary contact recreation and public access, areas where TMDL implementation plans have been established as well as other criteria for consideration. DEQ has elected to focus on the two most recent years of monitoring data because the 2012 nationally recommended criteria do not have a recommended return interval of excursions of the geometric mean criteria. Additionally, an evaluation of more recent data ensures that the primary contact recreation use attainment will be determined from a range of representative environmental conditions rather than from extreme wet-weather events that may occur sporadically within the six-year assessment window.

Ten or more samples in a 90-day period are required to calculate a geometric mean. No exceedances of the geometric mean and less than a 10 percent exceedance rate of the STV will result in a ‘fully supporting’ determination.

A decision of impairment will be made when there are one or more 90-day periods characterized by either; a) a geometric mean exceedance, b) a STV exceedance rate greater than 10 percent (when the sample size is greater than or equal to 10) and/or c) typically two or more STV exceedances in a small dataset ( $n < 10$ ), due to the uncertainty with small datasets, but this scenario will be evaluated on a case by case basis.

Bacteria densities reported as both Colony Forming Units (CFU) and Most Probable Number (MPN) shall be assessed against the numeric values in [9VAC25-260-170-A](#), pursuant EPA’s approval of the methods specified in [40 CFR Part136.3](#). Approved test methods that report either unit shall be used for assessment.

#### Rule 4

Conventional parameter data generated by probabilistic monitoring (ProbMon) networks will be used to create a general overview of those waters and to direct targeted monitoring in the future. For most ProbMon stations, only one data point per parameter will be available, providing insufficient information for determination of impairment. For ProbMon stations with two data points for conventional parameters, assessment will be the same as any station with two or more data points.

Benthic data will be compared to the Virginia Stream Condition Index (VSCI) or Virginia Coastal Plain Macroinvertebrate Index (VCPMI) and assessed accordingly.

A single “grab sample” exceedance of human health or aquatic life toxic criteria is assessed as insufficient information and follow-up monitoring should be conducted as resources allow to determine if the water is impaired. A single chronic or acute exceedance of a 30-day semi-permeable membrane device (SPMD) sample for a toxic parameter associated with aquatic life and wildlife use is considered insufficient information.

#### Rule 5

To be eligible for assessment, a continuous monitoring dataset must cover at least 30 days (consecutive or otherwise), except in the assessment of maximum hourly temperature change criteria, which may be assessed on a dataset spanning no less than 15 days. The continuous monitoring dataset will have undergone rigorous and standardized QA/QC screening before analysis. If a continuous monitoring dataset is used to place a water on the 303(d) Impaired Waters List, then an additional continuous monitoring dataset, collected during a subsequent year, during the same month(s) as the listing dataset, must be used to delist it. See Section 5.12 for detailed assessment methodology.

#### Rule 6

When data are insufficient for the determination of use attainment but indicate possible impairment, additional monitoring should be considered. In past report cycles the parameter status “Observed Effects” was used to categorize indications in the form of single sample WQS exceedances, sediment

and fish tissue screening value exceedances, observed pollutants or signs of water quality degradation (i.e., fish kills) lacking specific standards, or lower quality data that point to possible impairment (e.g., high bacteria counts on a Coliscan<sup>®</sup> plate). To better define these scenarios where data are available, but no numeric criteria or assessment thresholds developed for this guidance manual apply, a new Category 3E has been added to Virginia's IR in the 2024 cycle. These waters are of concern to the state, but no water quality standard exists. Data show exceedances of a state screening value, failure of a toxicity test or have been identified as a most probable stressor during development of an EPA approved TMDL. This rule applies to conventional and toxic parameters (water column, sediment, nutrient, and fish tissue) as well as biological monitoring. Waters with single WQS exceedances or lower quality data that warrant follow-up monitoring would be Category 3B or Category 3C if non-agency data.

## Rule 7

Waters that are suspected to be impaired due to naturally occurring, non-anthropogenic conditions will be classified as Category 5C (possibly needing a TMDL) of the Integrated Report. Examples of natural impairments include low DO and/or pH in slow-flowing waters or high temperature from thermal springs. If EPA agrees with DEQ's assessment that the exceedances are a result of natural conditions, the parameter will then be listed in Category 4C (impaired but not needing a TMDL). For waters in Category 5C or 4C, the water quality standards will be reviewed and possibly updated during the next triennial review to reflect variations caused by natural conditions for these waters. Once appropriate water quality standards are in place, data will be reviewed again to determine whether these waters meet or exceed designated uses. It may be necessary to conduct a TMDL study or Use Attainability Analysis prior to standards modification in order to determine and/or verify the appropriate criteria based on natural pollutant loadings.

Dissolved oxygen should not be listed as an impairment cause in Class VII waters lacking human-induced pollutant sources, per [9VAC25-260-50](#). When available, other data—such as fish community composition, habitat assessment, benthic macroinvertebrate composition, etc. —should be evaluated against the narrative criterion to determine use support.

## Rule 8

Waters that have been assessed as impaired (Category 4 or 5) will continue to be tracked in the DEQ Comprehensive Environmental Data System (CEDS), the Assessment, Total Maximum Daily Load (TMDL) Tracking and Implementation System (ATTAINS) and the monitoring station list, whether they have recent monitoring data or not. These waters will retain the results of previous assessments for all impaired designated uses. Waters classified as Category 5 will carry this designation until a TMDL is developed, in which time the water will move to Category 4. Category 4 or 5 waters will carry the impaired designation until additional monitoring data reveal the waters are no longer impaired. (Justification must be provided to EPA before removing an impaired water from the 303(d) Impaired Waters List.) In contrast, fully supporting (Category 2) waters can only retain their status for two additional assessment cycles (based on data up to 4 years outside the current assessment data window) with no new data collected. After two assessment cycles with no data in the assessment data window, the water will be classified as “indeterminate”, and the outdated supporting data will be removed from the monitoring station list. The water will remain as “indeterminate” until new data is collected and assessed. Furthermore, there should not be a change in the assessment determination made on partial datasets that remain in the assessment data window from previous assessment cycles. The initial

determination based on the original dataset should remain in place and follow this rule until new data is collected.

#### Rule 9

In previous assessment cycles, for effluent limited waters, if the VPDES permit has been issued with a scheduled compliance date that extends beyond the next 303(d) listing or permit cycle, the water would be listed as Category 5E. If the compliance date falls within the next listing cycle or within the current permit cycle, whichever is longer, the water would be listed in Category 4B. The evaluation of effluent limited waters was removed during the 2022 assessment cycle. See Part VII for additional information.

#### Rule 10

Duplicate and/or split samples collected for QA/QC purposes will not be used in the assessment. The primary sample (S1) will be assessed against the appropriate standard and the duplicate/split sample (S2) will be used only to document lab analysis quality control.

#### Rule 11

Sampling stations that happen to be located within a permitted mixing zone, primarily via probabilistic monitoring, will not be individually assessed for aquatic life use. They will be included with the overall probabilistic assessment. Any other stations that inadvertently were located in mixing zones will not be assessed individually for aquatic life use as the use is exempt in mixing zones.

#### Rule 12

A review of stockable and some natural trout waters currently listed as impaired has revealed that many of these impairments are due to erroneous segment boundaries or natural conditions. Both issues were addressed as part of Virginia's most recent review of its water quality standards. For the 2024 assessment, these waters will be categorized as Category 2A if specifically addressed via amendments to the water quality standards regulation and currently meeting new criteria or 4C (impaired due to natural conditions and not needing a TMDL) as long as supporting documentation is provided by the Virginia Department of Wildlife Resources (DWR) and endangered species are not being adversely affected. This should prevent and/or correct the misclassification of these segments, pending any amendments during Triennial Review. Once these standard reviews are completed and EPA approved, these waters will be re-evaluated and classified accordingly.

#### Rule 13

Data associated with multi-probe meters are to be rounded to one decimal place to best reflect the accuracy of the probes.

#### Rule 14

Nested impairments are those waters that are located within a watershed where there is an existing TMDL for the listing parameter. A TMDL requires a specific loading reduction for all waters within the watershed whether they currently meet the standard or not. Once the TMDL is approved by EPA, any waters within the watershed that are subsequently assessed as impaired for the parameter targeted by the

TMDL will be considered Category 4A (impaired but has a TMDL). Nested impairments are normally bacteria related to recreation and shellfish consumption uses, but may also include benthic impairments (aquatic life use) depending on the stressors involved.

#### Rule 15

Division of Consolidated Laboratories (DCLS) has determined that total dissolved ammonia (STORET Parameter Code 00608) and total ammonia (STORET Parameter Code 00610) are essentially the same parameter. Thus, where only dissolved ammonia data are available, these will be used to assess against the total ammonia criteria. Where both data are available, total ammonia should be used to assess the criteria, unless the total ammonia value is remark coded as less than the detection limit and the dissolved ammonia value is not. In which case the dissolved ammonia should be used to assess the criteria.

#### Rule 16

Shellfish waters where restrictions or prohibitions are due solely to the location of a discharge outfall and associated buffer zone or where the use is deemed too limited to harvest due to low salinity or other natural reasons—and not due to water quality exceedances—will not be included in the 303(d) list. In these cases, assessment is not conducted as the shellfish designated use has been administratively removed through the issuance of a discharge permit or prohibition on harvesting and there is no instream data indicating impairment.

#### Rule 17

Uncensored values should be used when evaluating data for the water quality assessment. Two exceptions include: bacteria parameters, where the Lower Detection Limit reported from the lab will be used for analysis and benthic chlorophyll-a, where a model-predicted value will be used to represent sampling events with less than 10% filamentous algae cover observed.

#### Rule 18

All instances of rounding in the assessment process will use the “round to even” approach to ensure consistency. The memo describing the approach is included as [Appendix I](#).

#### Rule 19

Per [9VAC25-260-50](#), dissolved oxygen, temperature and pH data collected in a waterbody where the projected flow was below 7Q10 on the sample date will not be assessed.

#### Rule 20

Parameters with established water quality criteria that were identified as most probable stressors and had a TMDL equation assigned during development of EPA approved TMDLs will be listed in Category 4 along with the original cause of impairment. When the original cause of impairment shows improvement and will be proposed for delisting, both the original cause parameter and the probable stressor parameter can be delisted concurrently. No additional TMDL will be required when most probable stressor parameters are added to Category 4.



## **PART V. ASSESSMENT METHODOLOGY**

### *5.1 Monitoring Station Siting and Delineation*

DEQ has a vast network of active Ambient Water Quality Monitoring (AWQM) stations and a number of biological stations statewide. The AWQM stations are generally monitored bimonthly while the biological stations are normally monitored twice a year (usually in the spring and fall). Monitoring programs can be designed based on a “targeted” (conventional) approach or a “probability based” random selection approach or a combination of the two. Each monitoring program design has its advantages and disadvantages. Historically, most of DEQ’s monitoring strategy has been based on the conventional approach. Many of the stations were located in proximity to (above and below) Virginia Pollutant Discharge Elimination System (VPDES) facility outfalls. During this reporting cycle, DEQ has continued to use a rotating watershed approach where stations are sited for two years of bimonthly sampling within a selected major river basin. The number of stations per watershed is based on the drainage area of the watershed and the Department of Conservation and Recreation (DCR) “nonpoint source potential” rating of the watershed. In order to provide consistency between the regional planning staff and to get an accurate number of assessed stream miles in Virginia, the following stream delineation guidelines are the primary considerations used in the assessment unit (segment) size decisions. However, in certain cases, best professional judgment of the regional staff may be used if the delineation results are contrary to these guidelines. Where appropriate, documentation of these decisions should be included in the segment narrative.

1. Typically, no more than 10 miles of free-flowing stream should be assessed by conventional pollutant data from one ambient monitoring station. Miles assessed for a toxic pollutant or biological impairment may vary from the miles assessed for conventional parameters.
2. One monitoring station should not be used to assess an entire watershed unless land use, source, and habitat are relatively homogeneous.
3. When determining the miles assessed for a free-flowing monitoring station, the following items need to be considered:
  - a) WQ Standards use designations (i.e. classes and/or special standards)
  - b) point and/or nonpoint source input to the stream or its tributaries,
  - c) watershed characteristics such as land use,
  - d) local habitat characteristics such riparian vegetation, stream banks, substrate, slope, or channel morphology,
  - e) entry of a large tributary or diversion, or
  - f) hydrologic features such as channelization or dams.

4. For non-Chesapeake Bay Program tidal and estuarine stations, EPA guidance suggests using a 4-mile radius for open water stations; a 2-mile radius for sheltered bay stations, and a 0.5 mile radius for highly sheltered bay stations<sup>3</sup>.
5. Segment delineation will be performed using the USGS National Hydrography Dataset (NHD) coverage or other appropriate GIS dataset.
6. Spatial coverage for estuarine probabilistic monitoring stations should be identified in conjunction with the development of the monitoring plan and coordinated by regional monitoring and assessment staff and/or the Chesapeake Bay Program monitoring coordinator and Bay monitoring staff. Estuarine B-IBI data will be assessed according to the methodology described in Appendix G.
7. If the mixing zone of a VPDES-permitted facility exists in an impaired segment, the parameter-specific length of the mixing zone is specifically understood to be separate from the impaired segment, even though the boundaries of the segment and/or its description may show the impairment as continuous.
8. Probabilistic stations in free-flowing waters will not be delineated into 303(d) segments unless they are characterized by toxics data, biological data, or more than one measurement of a conventional parameter.
9. Assessment units with more than one monitoring station should not be assessed based on pooled results from individual stations, unless specifically described in assessment methodology (i.e., data collected in the same assessment unit in Section 187 lakes/reservoirs). Where two stations' assessment determinations differ within an assessment unit, in most cases, the unit should be split to accurately reflect conditions in the area.

## 5.2 Evaluation of Designated Uses

The 305(b) process assesses a total of six primary designated uses, as appropriate for a particular waterbody, based on Water Quality Standards: wildlife, aquatic life, recreation, fish consumption, shellfishing, and public water supply uses. The Chesapeake Bay criteria, adopted in 2005, have sub-divided the aquatic life use into several distinct sub-uses. Following are details relating to the assessment of the six designated uses of Virginia's waters.

### 1. **Wildlife Use:**

Wildlife use includes the propagation, growth, and protection of a balanced, indigenous population of wildlife. Support of wildlife use is determined by assessing the toxic standards for aquatic life found in 9 VAC-25-260-140 B. These criteria were developed to protect aquatic life as well as wildlife. For toxic pollutant assessment, waters where there are two or more samples and no exceedances of aquatic life criteria within most of the 3-year periods for which there are a sufficient number of samples within the assessment window, using grab samples or SPMD data, are considered fully supporting for wildlife use. Waters where there are two or more exceedances of the same WQS aquatic life toxic criteria in most of the 3-year periods for which there are a

---

<sup>3</sup> [https://www.epa.gov/sites/default/files/2015-09/documents/guidelines\\_for\\_preparation\\_of\\_the\\_comprehensive\\_state\\_water\\_quality\\_assessments\\_305b\\_reports\\_and\\_electronic\\_updates\\_1997\\_supplement-volume2.pdf](https://www.epa.gov/sites/default/files/2015-09/documents/guidelines_for_preparation_of_the_comprehensive_state_water_quality_assessments_305b_reports_and_electronic_updates_1997_supplement-volume2.pdf)  
*Final Guidance for 2024 IR Assessment Methodology*

sufficient number of samples within the assessment window, using either grab samples or SPMD data, are considered impaired for the wildlife use.

For toxic pollutant assessments, both chronic and acute criteria can be assessed whenever sufficient data are available as applicable. Chronic criteria are to be assessed when multiple grab samples are collected within two separate four-day periods within a three-year period, or when there are two or more separate 30-day SPMD deployments within a three-year period. Two samples (either grab or SPMD) taken within three consecutive years are sufficient to assess acute criteria.

For ammonia in free-flowing and tidal waters, acute criteria are a one-hour average concentration not to be exceeded more than once every three years on the average, and chronic criteria are 30-day average concentrations not to be exceeded more than once every three years on the average. In addition, the four-day average concentration of total ammonia nitrogen (in mg N/L) shall not exceed 2.5 times the chronic criterion within a 30-day period more than once every three years on the average in free-flowing streams. See [9VAC-25-260-155](#) for additional information on calculating the ammonia criteria and applicable aquatic life considerations.

## **2. Aquatic Life Use:**

Aquatic life use includes the propagation, growth, and protection of a balanced indigenous population of aquatic life (including game and marketable fish) which may be expected to inhabit the waters.

Support of aquatic life use can be determined by the assessment of conventional parameters (dissolved oxygen, pH and temperature), toxic pollutants in the water column (relative to water quality standards), toxic pollutant analysis of sediments, toxicity testing, nutrient analysis and/or the biological assessment of benthic communities. All available data, relative to aquatic life use, shall be considered to determine if the aquatic life use is being supported. This assessment includes the sub-categories of aquatic life use associated with the Chesapeake Bay criteria.

Conventional parameters are assessed using the “Percent Method”. A 10.5% exceedance threshold is used for determining full support or impairment for conventional pollutants in datasets greater than or equal to 10 samples. An exceedance rate that is > 10.5% with at least two exceedances is normally considered impaired. An exceedance rate ≤ 10.5% is considered fully supporting. A single exceedance in a small dataset (2-9 samples) is usually considered insufficient, due to the uncertainty around small datasets, but this scenario will be evaluated on a case-by-case basis.

For dissolved oxygen, the instantaneous minimum standard is used to assess exceedances unless continuous monitoring data are available to assess the daily average. See Section 5.7 for assessment methodology for lakes and reservoirs. See Section 5.12 for assessment methodology for continuous monitoring. Dissolved oxygen in the Chesapeake Bay and its tributaries is assessed according to the method outlined in Section 5.3.

For free-flowing stream benthic macroinvertebrate assessments, data for the overall assessment period is rated as non-impaired when the Virginia Stream Condition Index (VSCI) or the Virginia Coastal Plain Macroinvertebrate Index (VCPMI) scores are at or above their respective impairment thresholds. (60 for the VSCI or 40 for the VCPMI). See Section 5.4 for additional methodology for biological assessments.

A project to refine the estuarine biological assessment methodology (B-IBI) was completed in 2006 and subsequently used in assessment decisions. The same methodology will be used again for 2024. See Section 5.3 and Appendix G for more information.

For toxic pollutant assessments, both chronic and acute criteria can be assessed whenever sufficient data are available as applicable. Chronic criteria are to be assessed when multiple grab samples are collected within two separate four-day periods within a three-year period, or when there are two or more separate 30-day SPMD deployments within a three-year period. Two samples (either grab or SPMD) taken within three consecutive years are sufficient to assess acute criteria. Waters where there are two or more exceedances of the same WQS aquatic life toxic criteria in most of the 3-year periods for which there are a sufficient number of samples within the assessment window, using either grab samples or SPMD data, are considered impaired for the aquatic life use.

For ammonia in free-flowing and tidal waters, acute criteria are a one-hour average concentration not to be exceeded more than once every three years on the average, and chronic criteria are 30-day average concentrations not to be exceeded more than once every three years on the average. In addition, the four-day average concentration of total ammonia nitrogen (in mg N/L) shall not exceed 2.5 times the chronic criterion within a 30-day period more than once every three years on the average in free-flowing streams. See [9VAC-25-260-155](#) for additional information on calculating the ammonia criteria and applicable aquatic life considerations.

### **3. Fish Consumption Use:**

Fish consumption use includes the propagation, growth and protection of a balanced population of aquatic life including game and marketable fish. Human health is also a primary consideration with regards to fish consumption use. Support of this use is determined using three separate criteria.

First, support or lack thereof, is based on human health related advisories and/or restrictions issued by the Virginia Department of Health (VDH) based on monitoring data. The fish consumption use is determined to be impaired when the public is advised by VDH that fish consumption is prohibited for the general population or an advisory that certain fish species should not be consumed by the general population or sub-populations at greater risk, such as children and/or pregnant women. A water may be delisted once monitoring data supports lifting all fish consumption advisories within an assessment unit.

Second, a fish tissue sample (either a single specimen sample or composite sample) is compared to WQS criterion-based tissue values (TVs) and tissue screening values (TSVs) for toxic pollutants. No exceedances of a particular TV listed in Appendix E-1 results in a fully supporting assessment of the water for the fish consumption designated use, if there are no concurrent VDH fish consumption advisories. Two or more sample exceedances of a particular TV listed in Appendix E-1 results in an impaired assessment of the water for the fish consumption designated use. A single sample above a TV or TSV in the assessment window results in an insufficient information assessment and should be prioritized for follow-up monitoring.

Third, support of the fish consumption use is determined by comparison of water column or semi-permeable membrane device analytes to the human health criteria in public water supplies and other surface waters, as listed in the WQS ([9 VAC-25-260-140 B](#)). Criteria for human health metals should be assessed using total recoverable concentrations.

#### **4. Shellfishing Use:**

Shellfishing use includes the propagation, growth and protection of a balanced population of aquatic life including marketable shellfish.

Use support is based on the determination of restrictions or condemnations on the harvesting and marketability of shellfish resources made by the VDH-Division of Shellfish Sanitation (DSS) as of the most recent condemnation list (December 2022) associated with the reporting period. DSS has the statutory authority to determine shellfish harvesting and marketability status. DSS uses four classifications for describing the status of shellfish waters:

Open:	Growing areas from which shellfish may be taken for direct marketing at all times.
Conditionally Condemned, Seasonally Restricted:	Growing areas where the water quality may be affected by seasonal or sporadic use of boat docks or harbor facilities are considered conditionally approved. Normally, this would occur during the boating season (April 30 through October 31).
Restricted:	Growing areas where a sanitary survey indicates a limited degree of pollution which makes it unsafe to market shellfish for direct marketing. Shellfish from such areas may be marketed after purifying or relaying activities in accordance with certain VDH-DSS requirements.
Prohibited, Prohibited-Nonproductive:	Growing areas where the harvesting of shellfish from these areas for direct marketing, relaying, or depuration is prohibited. The sanitary survey may indicate dangerous numbers pathogenic microorganisms or other contaminants that might reach that area. Additionally, prohibited areas due to administrative closures.

Specific information regarding DSS assessment methodology and the listing/delisting flowchart for shellfish waters can be found in Appendix C of this guidance document. For the 305(b)/303(d) Integrated Report, listing and delisting will be based on instream data assessed for the reporting period. However, as the TMDL begins development, if new or more recent data shows the shellfish water is no longer impaired, a petition for delisting will be crafted and submitted to EPA for their approval by the Watershed Program (TMDL) staff.

#### **5. Recreation/Swimming Use:**

Recreation use assessment includes swimming and other primary and secondary water contact recreation uses such as water skiing and pleasure boating. Support or lack thereof of the recreation use is based on comparison of primary contact recreational criteria and/or VDH closures or advisories due to excessive bacteria, the presence of harmful algal blooms (HABs), or nuisance algal growth.

*Bacteria*

Virginia's adoption of the EPA 2012 nationally recommended primary contact recreational water quality criteria has modified the assessment methodology for bacteria data. For each assessment unit, primary contact recreation use assessments should be based on bacteria monitoring data collected within the two most recent calendar years of the six-year assessment window.

A decision of impairment will be made when there are one or more 90-day periods characterized by either; a) a geometric mean exceedance, b) a STV exceedance rate greater than 10 percent (when the sample size is greater than or equal to 10) and/or c) typically two or more STV exceedances in a small dataset ( $n < 10$ ), due to the uncertainty with small datasets, but this scenario will be evaluated on a case by case basis.

A decision of impairment is also determined if one or more VDH beach closure(s) and/or two or more advisories > 1 week duration with medium or high probability of recurrence (based on bacteria monitoring data) occurs within the current assessment window. A single short-term (< 1 week) VDH beach closure/advisory (based on bacteria monitoring data) with low probability of recurrence will result in a 'insufficient information' determination indicating follow-up monitoring is needed.

No exceedances of the geometric mean and 10 percent or less exceedance rate of the STV in any 90-day periods will result in a fully supporting determination.

For additional assessment information on bacteria monitoring in lakes/reservoirs, see Section 5.7 Lakes/Reservoirs Assessment.

### *Harmful Algal Blooms (HABs)*

The Virginia Department of Health (VDH) is the agency responsible for issuing notices and swimming advisories due to the potential or confirmed presence of HABs. Advisories are issued based on VDH's "Guidance for Cyanobacteria Bloom Recreational Advisory Management" (VDH HAB Guidance) which is posted to the VDH website: <https://www.vdh.virginia.gov/waterborne-hazards-control/harmful-algal-blooms/>. VDH uses DEQ's monitoring data to inform advisory determinations. Impairment status for the recreation use due to HABs for a waterbody is based upon presence or absence of a VDH swimming advisory and the applicable monitoring data collected to confirm the HAB, all conducted in accordance with VDH HAB Guidance and DEQ Monitoring Protocols.

A waterbody would be assessed as fully supporting for the recreation use if a VDH swimming advisory was issued in the two most recent years of the assessment window but was lifted after the minimum required follow-up sampling providing evidence which indicate there are no persistent HABs.

A waterbody would be assessed as impaired for the recreation use if there is an issued VDH swimming advisory and the HAB is confirmed through the initial follow-up sampling events in the two most recent calendar years of the assessment window. The regional assessor should delineate the impairment using the boundaries included in the VDH advisory.

A waterbody impaired for the recreational use due to HABs may be delisted and assessed as Fully Supporting if, within the 6-year assessment window, there are no VDH swimming advisories or there are VDH swimming advisories that are lifted after the minimum required follow-up sampling providing evidence which indicate there are no persistent HABs.

## *Water Quality Impacts Due to Algal Growth*

Since the early 2010s, DEQ has fielded concerns over excessive filamentous algae in certain sections of the Shenandoah River basin. Citizen complaints have centered on the negative impact that filamentous algae blooms have on stream aesthetics and recreational activities such as swimming, wading, kayaking, and fishing. In response, DEQ conducted a special study during 2016-2019 with two goals: 1) To develop a scientifically based, defensible, and reproducible field method for quantifying filamentous algae growth and 2) to develop a body of information that could be used to determine benthic chlorophyll-a thresholds for nuisance filamentous algae growth. DEQ has chosen to develop criteria for benthic chlorophyll-a, because it is a quantitative estimate of algal biomass.

The benthic chlorophyll-a criteria apply in certain wadeable portions of the mainstem sections of the Shenandoah River, North Fork Shenandoah River and South Fork Shenandoah River. The boundaries are set forth in Virginia's Water Quality Standards 9VAC25-260-310 (ii). The recreation use is assessed using either a two-month median or seasonal median for benthic Chlorophyll-a. A determination of persistent nuisance filamentous algae impeding the recreation use should be made when exceedances of either of the specified benthic chlorophyll-a concentration thresholds occur in more than one recreation season (May 1 to October 31) in three years.

### **Assessment Methodology**

To allow for the appropriate calculation of the two-month median and seasonal median, a value of 34 mg/m<sup>2</sup> will be used as the conservative left-censored value to represent monitoring events where less than 10% cover was observed and no benthic chlorophyll-a sample was taken (per DEQ monitoring protocols). This value reflects that the site was monitored but the collection of a benthic chlorophyll-a sample was not warranted because little to no algae was present.<sup>4</sup>

The following scenarios indicate an impairment of the recreation use due to persistent nuisance filamentous algae:

- where the median of at least two samples taken in two consecutive months exceeds 150 mg/m<sup>2</sup> in more than one year within any three-year-period in the assessment window, or
- a median of at least three samples collected in separate months in one season above the seasonal criterion of 100 mg/m<sup>2</sup> in more than one year within any three-year-period in the assessment window.

A waterbody will be considered fully supporting if the assessment dataset shows there are no three-year periods in the assessment window for which an exceedance of either the two-month median or seasonal median criteria occurred in more than one recreational season. This scenario would also constitute a delisting of a recreation impairment due to benthic chlorophyll-a.

Filamentous algae data or information submitted to DEQ for evaluation that is collected by citizen or non-agency organizations on wadable waterbodies and that do not meet agency QA/QC protocols, will be considered Insufficient Information and prioritized for follow-up monitoring (Category 3C). A single sample within an assessment unit collected by DEQ in response to user complaints on wadable waterbodies within the segments identified in 9VAC25-260-310 (ii) will be considered Insufficient Information and considered for

---

<sup>4</sup> The value of 34 mg/m<sup>2</sup> is derived from an exponential regression model relating percent algal coverage to chlorophyll-a concentration. It is equal to the upper 95% confidence limit of the predicted chlorophyll-a concentration at 10% algal coverage. This conservative estimate is informed by the entire Shenandoah benthic chlorophyll-a dataset. This value will be recalculated for each assessment cycle based on all available % cover and benthic chlorophyll-a data. The word 'sample' in this section refers to both the censored values and lab reported values used in assessment determinations.

### *Final Guidance for 2024 IR*

#### *Assessment Methodology*

future monthly monitoring as regional resources allow (Category 3B). Filamentous algae data<sup>5</sup> collected by DEQ in response to user complaints on wadable waterbodies outside of segments identified in 9VAC25-260-310 (ii) will be considered Insufficient Information and considered for follow-up monitoring (Category 3E). These waterbodies may be considered for inclusion in 9VAC25-260-310 (ii) in the future.

**6. Public Water Supply Use:**

Waters that are used for public drinking water supply (PWS) are identified in the WQS and are protected by additional health-related standards that are applicable only to these waters. Taste and odor criteria to maintain acceptable taste, odor or aesthetic quality of drinking water apply at the only at the drinking water intake. Support or lack thereof of this use is based on a comparison of water column data to applicable public water supply criteria, as well as VDH closures or advisories due to excessive pollutant(s) in treated water or samples from a distribution system. Criteria for human health metals should be assessed using total recoverable concentrations.

Impairment is determined if the median of two or more samples of human health criteria are observed in PWS designated waters within the assessment window or one or more VDH public water supply source closures due to contamination are issued within the two most recent years of the assessment window, with a medium to high probability that the contamination will recur.

A single exceedance of human health criteria in PWS designated waters within the assessment window or a single VDH closure with low probability of recurrence within the two most recent years of the assessment window would result in an ‘insufficient information’ determination indicating follow-up monitoring is needed.

If an exceedance of a taste and odor criteria is observed within the PWS designated area, follow up monitoring at the closest drinking water intake should occur as regional office resources allow to make an assessment determination. No exceedances of human health criteria in PWS waters with two or more samples within the assessment window will result in a fully supporting assessment determination.

Table 1 and 2 summarize the kinds of information required to establish designated use support.

**Table 1. Designated Use Matrix**

<b>DESIGNATED USE</b>	<b>USE DESCRIPTION/INDICATORS</b>
<b>Aquatic Life Use, Chesapeake Bay sub-uses</b>	Description: The propagation, growth, and protection of a balanced indigenous population of aquatic life that may be expected to inhabit a waterbody
	Indicators: Dissolved oxygen, pH, temperature, chlorophyll a*, nutrients*, water column and sediment toxics, toxicity tests, benthics, submerged aquatic vegetation
<b>Fish Consumption Use</b>	Description: Game and marketable fish species that are safe for human health
	Indicators: VDH notices, fish tissue toxics, water column toxics
<b>Shellfishing Use</b>	Description: Marketable shellfish (clams, oysters, mussels) that are safe for human health
	Indicators: VDH notices
<b>Recreation (Swimming) Use</b>	Description: Swimming, boating, and other recreational activities
	Indicators: VDH notices for bacteria and harmful algal blooms, bacteria, benthic chlorophyll-a**

<sup>5</sup> Filamentous algae data include visual bank observations, the results of FILBEN analysis from DCLS (which includes both chl a and chl b, pheophyton, and AFDM) and biomass (wet-wrung weight) collected per DEQ’s filamentous algae QAPP and SOP.



DESIGNATED USE	USE DESCRIPTION/INDICATORS
Public Water Supply Use	Description: Drinking water safe for human health
	Indicators: VDH notices, water column toxics
Wildlife Use	Description: The propagation, growth, and protection of associated wildlife
	Indicators: Water column toxics

\*Chlorophyll *a* and nutrients (total phosphorus) are assessed only in the lakes listed in Section 187 of the WQS. Chlorophyll *a* is also assessed in the tidal James River.

\*\*Benthic Chlorophyll-*a* is only assessed in sections of the Shenandoah River, North Fork Shenandoah River and South Fork Shenandoah River per 9VAC 25-260-310.ii.

**Table 2. Designated Use Assessment Methodology**

Designated Use	Parameter/Data Type	Fully Supporting	Insufficient Information (prioritized for follow-up monitoring)	Impaired
Aquatic Life Wildlife (toxics only)	Conventional <sup>1</sup>	<ul style="list-style-type: none"> <li>Exceedance rate <math>\leq</math> 10.5% for field parameters</li> <li>Median lacustrine TP<sup>2</sup> below criterion</li> <li>90<sup>th</sup> percentile lacustrine chlorophyll <i>a</i> below criterion</li> </ul>	<ul style="list-style-type: none"> <li>Level II<sup>3</sup> data with an exceedance rate <math>&gt;</math> 10.5%</li> <li>Single exceedance in small dataset (samples <math>&lt;</math>10)</li> </ul>	<ul style="list-style-type: none"> <li>Exceedance rate <math>&gt;</math> 10.5% for field parameters, with a minimum of two exceedances</li> <li>Median lacustrine TP above criterion</li> <li>90<sup>th</sup> percentile lacustrine chlorophyll <i>a</i> above criterion</li> </ul>
	Biological	Benthic index scores $\geq$ impairment threshold	<ul style="list-style-type: none"> <li>Level II<sup>3</sup> data suggest degraded community</li> <li>Benthic index score conflicts with biologist's best professional judgment</li> </ul>	Benthic index score $<$ impairment threshold
	Toxics	<ul style="list-style-type: none"> <li>No exceedances of the same chronic or acute aquatic life criteria in a 3-year period</li> </ul>	<ul style="list-style-type: none"> <li>A single exceedance of chronic aquatic life use criteria using temporally aggregated water column grab samples in a 3-year period or one SPMD sample exceedance of chronic aquatic life use criteria in a 3-year period</li> <li>Single water column grab or SPMD sample exceedance of acute aquatic life use criteria in a 3-year period</li> <li>One or more toxicity test failures or sediment screening value exceedances (aquatic life only)</li> </ul>	Two or more grab or SPMD exceedances of the same chronic or acute aquatic life criteria in a 3-year period
Recreation	Bacteria	<ul style="list-style-type: none"> <li>No geometric mean exceedances and STV exceedance rate <math>\leq</math> 10% in a 90-day period</li> </ul>	<ul style="list-style-type: none"> <li>Level II<sup>3</sup> data with one or more geometric mean exceedances or STV exceedance rate <math>&gt;</math> 10% in a 90-day period</li> <li>Single STV exceedance in one or more 90-day periods with less than 10 samples due to uncertainty with small datasets</li> </ul>	<ul style="list-style-type: none"> <li>One or more geometric mean exceedance(s) in a 90-day period</li> <li>STV exceedance rate <math>&gt;</math> 10% in a 90-day period, usually with a minimum of two exceedances</li> </ul>
	Benthic Chlorophyll- <i>a</i> <sup>6</sup>	No three-year periods in the assessment window for which an exceedance of either the two-month median or seasonal median criteria occurred in more than one recreational season	<ul style="list-style-type: none"> <li>A single sample collected in response to user complaints on segments identified in the Water Quality Standards</li> <li>Data collected on waterbodies outside of the segments identified in the Water Quality Standards</li> </ul>	<ul style="list-style-type: none"> <li>The median of at least two samples taken in two consecutive months exceeds 150 mg/m<sup>2</sup> in more than one year within any three-year-period in the assessment window</li> <li>A median of three samples collected in separate months in one season above the seasonal criterion of 100 mg/m<sup>2</sup> in more than one year within any three-year-period in the assessment window.</li> </ul>

Designated Use	Parameter/Data Type	Fully Supporting	Insufficient Information (prioritized for follow-up monitoring)	Impaired
	VDH notice (Bacteria)	No closures/advisories and bacteria data meet criteria.	A single short-term (< 1 week) VDH closure/advisory with low probability of recurrence, based on bacteria data	One or more closure(s) and/or two or more advisories > 1 week duration with medium or high probability of recurrence, based on bacteria data
	VDH notice (HABs)	A VDH advisory that is lifted after the minimum follow-up sampling events in the two most recent years of the assessment window.		A VDH advisory that persists past the minimum follow-up sampling events in the two most recent years of the assessment window.
Shellfishing	VDH notice	Approved shellfish harvest waters or area classified as “conditionally condemned or seasonally restricted”		Areas classified as “restricted” or “prohibited”—excluding VPDES <sup>5</sup> outfalls and administrations closures where no instream data are available
Fish Consumption	Toxics	<ul style="list-style-type: none"> <li>No exceedances of fish tissue TVs or human health criteria where two or more samples have been collected.</li> </ul>	<ul style="list-style-type: none"> <li>Single exceedance of a human health criterion using grab sample or SPMD data</li> <li>Single sample exceedance of a tissue value or tissue screening value</li> </ul>	<ul style="list-style-type: none"> <li>The median of two or more water samples exceed a human health criterion using grab sample or SPMD data</li> <li>Two or more sample exceedances of a tissue value</li> </ul>
	VDH notice	No advisories and fish tissue data meet toxics criteria.	A VDH advisory which does not limit consumption is in effect, based on fish tissue data	A VDH advisory or restriction limiting or prohibiting consumption, based on fish tissue data
Public Water Supply	Toxics	<ul style="list-style-type: none"> <li>No exceedances of PWS WQS where two or more water column samples have been collected.</li> </ul>	A single exceedance of PWS WQS using grab or SPMD data	The median of two or more samples exceeds a PWS WQS using grab samples or SPMD data.
	VDH notice	No closures	A single VDH closure with low probability of recurrence in the two most recent years of the assessment window.	One or more closures with medium or high probability of recurrence in the two most recent years of the assessment window.

<sup>1</sup>Refer to [Section 5.3](#) for methodology specific to Chesapeake Bay criteria. Refer to [Section 5.7](#) for methodology specific to lakes and reservoirs. Refer to [Section 5.12](#) for methodology specific to continuous monitoring data.

<sup>2</sup>TP = total phosphorus concentration. Along with lacustrine chlorophyll a, only data from the most recent two years are aggregated. See [Section 5.7](#) for methodology specific to lakes/reservoirs.

<sup>3</sup> Level II data are lower-quality data submitted to DEQ from other sources. See [Part VI](#) for more information.

<sup>4</sup>SPMD = semi-permeable membrane device (an instrument that passively samples ambient toxics over some length of time)

<sup>5</sup>VPDES=Virginia Pollution Discharge Elimination System

<sup>6</sup>The benthic chlorophyll-a criteria only apply to certain sections of the Shenandoah River, North Fork Shenandoah River and South Fork Shenandoah River. See 9VAC 25-260-310.ii for boundary applicability.

### 5.3 Chesapeake Bay Assessment

In addition to assessment of criteria for state-wide aquatic life designated uses as described elsewhere in this document, the Chesapeake Bay and its tidal tributaries will be assessed for: 1) sub-categories of aquatic life use specific to the Chesapeake Bay estuarine system, and 2) the general narrative standard for aquatic life use through assessment of benthic invertebrate community condition. The following describes the aquatic life use sub-categories, applicable criteria, assessment process, and segmentation issues, as well as the Assessment, Total Maximum Daily Load (TMDL) Tracking and Implementation System (ATTAINS) and Integrated Reporting issues. Bay-specific criteria for dissolved oxygen, chlorophyll *a*, and submerged aquatic vegetation/water clarity are detailed in [9 VAC25-260-185](#).

- **Migratory Fish Spawning and Nursery Designated Use:** waters in the Chesapeake Bay and its tidal tributaries that protect the survival, growth and propagation of the early life stages of a balanced, indigenous population of anadromous, semi-anadromous, catadromous and tidal-fresh resident fish species inhabiting spawning and nursery grounds. This designated use extends from the end of tidal waters to the downriver end of spawning and nursery habitats that have been determined through a composite of all targeted anadromous and semi-anadromous fish species' spawning and nursery habitats (see boundaries in U.S. Environmental Protection Agency. 2004. *Technical Support Document for Identification of Chesapeake Bay Designated Uses and Attainability 2004 Addendum*. Chesapeake Bay Program Office, Annapolis, Maryland. This designated use extends horizontally from the shoreline of the body of water to the adjacent shoreline and extends down through the water column to the bottom water-sediment interface. This use applies February 1 through May 31 and applies in addition to the open-water use described in this subsection.
- **Shallow-Water Submerged Aquatic Vegetation Designated Use:** waters in the Chesapeake Bay and its tidal tributaries that support the survival, growth and propagation of submerged aquatic vegetation (rooted, underwater bay grasses). This use applies April 1 through October 31 in tidal-fresh, oligohaline and mesohaline Chesapeake Bay Program segments, and March 1 through November 30 in polyhaline Chesapeake Bay Program segments and applies in addition to the open-water use described in this subsection.
- **Open-Water Aquatic Life Designated Use:** waters in the Chesapeake Bay and its tidal tributaries that protect the survival, growth and propagation of a balanced, indigenous population of aquatic life inhabiting open-water habitats. This designated use applies year-round horizontally from the shoreline at mean low water, to the adjacent shoreline but the vertical boundaries change seasonally. October 1 - May 31, the open water aquatic life use extends through the water column to the bottom water-sediment interface. June 1 - September 30, if a pycnocline is present and, in combination with bottom bathymetry and water column circulation patterns, presents a barrier to oxygen replenishment of deeper waters, this designated use extends down into the water column only as far as the upper boundary of the pycnocline. June 1- September 30, if a pycnocline is present but other physical circulation patterns (such as influx of oxygen rich oceanic bottom waters) provide for oxygen replenishment of deeper waters, the open-water aquatic life designated use extends down into the bottom water-sediment interface (see boundaries in U.S. Environmental Protection Agency. 2004. *Technical Support Document for Identification of Chesapeake Bay Designated Uses and Attainability 2004 Addendum*. Chesapeake Bay Program Office, Annapolis, Maryland. This designated use includes the migratory fish spawning and nursery and shallow-water submerged aquatic vegetation uses.

- **Deep-Water Aquatic Life Designated Use:** waters in the Chesapeake Bay and its tidal tributaries that protect the survival, growth and propagation of a balanced, indigenous population of aquatic life inhabiting deep-water habitats. This designated use applies to the tidally influenced waters located between the upper and lower boundaries of the pycnocline where, in combination with bottom bathymetry (depth, contour & shape) and water circulation patterns, a pycnocline is present and presents a barrier to oxygen replenishment of deeper waters. In some areas, the deep-water designated use extends from the upper boundary of the pycnocline down to the bottom water-sediment interface (see boundaries in U.S. Environmental Protection Agency. 2004. *Technical Support Document for Identification of Chesapeake Bay Designated Uses and Attainability 2004 Addendum*. Chesapeake Bay Program Office, Annapolis, Maryland.) This use applies June 1 - September 30.
- **Deep-Channel Seasonal Refuge Designated Use:** waters in the Chesapeake Bay and its tidal tributaries that protect the survival of a balanced, indigenous population of aquatic life inhabiting deep-channel habitats. This designated use applies to the tidally influenced waters at depths greater than the lower boundary of the pycnocline in areas where, in combination with bottom bathymetry and water circulation patterns, the pycnocline presents a barrier to oxygen replenishment of deeper waters (see boundaries in U.S. Environmental Protection Agency. 2004. *Technical Support Document for Identification of Chesapeake Bay Designated Uses and Attainability 2004 Addendum*. Chesapeake Bay Program Office, Annapolis, Maryland.) This use applies June 1 through September 30.

### Assessment Process

Full details of the assessment processes are described in USEPA, *Ambient Water Quality Criteria for Dissolved Oxygen, Water Clarity, and Chlorophyll a for the Chesapeake Bay and Tidal Tributaries*, EPA 903-R-03-002, April 2003 and the 2004 (EPA 903-R-002 October 2004) and 2007 (CBP/TRS 285-07, EPA 903-R-07-003), 2007 (CBP/TRS 288/07, EPA 903-R-07-005), 2008 (CBP/TRS 290-08, EPA 903-R-08-001), 2010 (CBP/TRS 301-10, EPA 903-R-10-002), and 2017 (CBP/TRS 320-17, EPA 903-R-17-002) addenda. A very general summarization of key aspects of the process follows.

The assessment period for DO and water clarity shall be the most recent three consecutive years within the data window. When three consecutive years of data are not available, three years within the most recent data assessment window must be available and used for the assessment.

Attainment of the dissolved oxygen criteria shall be assessed through comparison of a cumulative frequency distribution of criteria exceedances to the applicable reference curve for each designated use. A first step in the process involves spatial interpolation and extrapolation of data collected at individual fixed locations to project water quality conditions throughout the segment. A subsequent step involves development of cumulative frequency distribution (CFD) of criteria exceedances combining both spatial and temporal domains for each segment-designated use combination. A final step is to compare this CFD of criteria exceedances against a reference CFD of allowable exceedances to determine if the criteria are attained.

The revised methodology for DO is described in the 2010 addendum. For DO, the algorithm used for the calculation of the pycnocline has been adjusted, and a 10% reference curve is being used for the assessment of the Open Water and Deep Channel designated uses.

The Migratory Fish Spawning and Nursery use criteria are assessed using continuous monitoring DO datasets. More on the process can be found in [Section 5.12](#).

For the Shallow Water Submerged Aquatic Vegetation use criteria, if the submerged aquatic vegetation (SAV) acreage criteria are met in any individual Chesapeake Bay Program segment, then the shallow-water submerged aquatic vegetation use is met in that segment. If the SAV acres are not met, then the water clarity criteria shall be examined with either a CFD methodology or a “water clarity acres” methodology. If sufficient water clarity is available to support SAV growth through either of these alternatives, then the Shallow Water Submerged Aquatic Vegetation use is met regardless of the number of acres of SAV in that segment.

### **James River Chlorophyll**

Chlorophyll criteria are only applicable to the mainstem portion of the tidal James River and shall be assessed over the six-year assessment window. For a James River segment to be assessed as fully supporting, it must meet all applicable chlorophyll criteria. Furthermore, segments assessed as “fully supporting” must have chlorophyll data for a minimum of five spring and summer seasons. A segment with more than two exceedances of the same seasonal mean criterion or a greater than 10% exceedance rate of an applicable short-duration criterion shall be assessed as impaired. For a segment that is attaining its applicable short-duration criterion and has two consecutive exceedances of the same seasonal mean criterion, additional lines of evidence must be evaluated before assessing the segment as fully supporting. The occurrence of at least one of the following in the segment of concern during either of the “exceedance” seasons shall trigger an “impaired” categorization:

1. VDH HAB advisory issued by the Division of Shellfish Safety
2. A fish kill documented by DEQ coinciding with chlorophyll concentrations greater than the magnitude of the applicable short-duration chlorophyll-a criterion or a HAB event documented and confirmed by the HAB Task Force.
3. More than 10% of dissolved oxygen samples are below the Open Water instantaneous minimum criteria (4.3 mg/l for water temperature greater than 29°C and 3.2 mg/l for water temperature less or equal to 29°C).
4. More than 10% of pH samples are less than 6.0 or greater than 9.0.

All chlorophyll data collected at one meter or less in a segment in the James River should be aggregated spatially by date using the Bay Interpolator Grid. A median of same-day chlorophyll samples should be calculated for each grid cell. For segments JMSOH, JMSMH, and JMSPH, the median of grid values should be calculated to represent the chlorophyll expression of a segment over a specific date, and the median of same-month chlorophyll values should be calculated to represent the chlorophyll expression of a segment over that month. The seasonal geometric mean shall be calculated from the monthly chlorophyll values for a segment.

For segment JMSTF2, chlorophyll data collected in the "upper zone" (from the upstream boundary at the fall line to approximately river mile 95 (N37° 23' 15.27" / W77° 18' 45.05" to N37° 23' 19.31" / W77° 18' 54.03")) should be pooled, in the manner described above, separately from chlorophyll data collected in the "lower zone" (from river mile 95 to the downstream boundary of JMSTF2). The seasonal geometric mean for each of these zones should be calculated from their respective monthly chlorophyll values. To calculate the seasonal segment-wide geometric mean, an area-weighted average of the zonal geometric means should be calculated using the following equation:

Upper Zone Geometric Mean x 0.41 + Lower Zone Geometric Mean x 0.59

For segment JMSTF1, chlorophyll data collected in the "upper zone" (from the upstream boundary of JMSTF1 to approximately river mile 67 (N37° 17' 46.21" / W77° 7' 9.55" to N37° 18' 58.94" / W77° 6' 57.14")) should be pooled, in the manner described in subdivision bb (2) of this section, separately from chlorophyll data collected

*Final Guidance for 2024 IR  
Assessment Methodology*

in the "lower zone" (between river mile 67 to the downstream boundary of JMSTF1). The seasonal geometric mean for each of these zones should be calculated from their respective monthly chlorophyll values. To calculate the seasonal segment-wide geometric mean, an area-weighted average of the zonal geometric means should be calculated using the following equation:

Upper Zone Geometric Mean x 0.49 + Lower Zone Geometric Mean x 0.51

Chapter II of the 2017 addendum (EPA 903-R-17-002 , CBP/TRS 320-17) describes the assessment approach for short-duration Bay DO criteria. For Bay assessment units that are categorized "4D" when all assessed DO criteria are attained, the minimum data requirements and decision rules described in this document must be met to qualify these assessment unit(s) for category 2. DO criteria will be assessed wherever there is available data that meet these requirements.

### **Assessment Units**

The Chesapeake Bay program segmentation scheme (*Chesapeake Bay Program Analytical Segmentation Scheme-Revisions, Decisions and Rationales: 1983 -2003*, CBP/TRS 268/04. Chesapeake Bay Program, Annapolis, Maryland) shall be used as the assessment unit to determine attainment of the criteria in this section for each designated use. The spatial boundaries of each aquatic life use subcategory within each of these CBP segment are described in the *Technical Support Document for Identification of Chesapeake Bay Designated Uses and Attainability 2004 Addendum*. Chesapeake Bay Program Office, Annapolis, Maryland. Assessment results for each CBP segment/designated use will determine the Integrated Report listing category of all waterbodies (i.e. all ATTAINS Assessment Units) geographically within that CBP segment/designated use. For example, the listing category of all tidal Onancock Creek assessment units will be determined by the appropriate designated use attainment of CBP segment CB7PH. In this example, it is likely that only open water and shallow water uses of CB7PH extend into Onancock Creek.

### **Assessment, Total Maximum Daily Load (TMDL) Tracking and Implementation System (ATTAINS) Reporting Units**

The Assessment, Total Maximum Daily Load (TMDL) Tracking and Implementation System (ATTAINS) is used to track assessment data for all designated uses in distinct geographically defined waterbodies across the state. ATTAINS assessment unit spatial boundaries are defined by many factors including the spatial distribution of available data to assess for designated uses. There may be several ATTAINS assessment units included in each Chesapeake Bay Program segment.

ATTAINS can only accept estuarine assessment units defined by surface areas (i.e. square miles). The complete water column within that assessment unit is assigned to a single overall aquatic life use attainment. Each individual Bay segment assessment unit may have deep channel, deep water, and open water sub-categories of aquatic life designated use (that may only account for a portion of the total volume/area of the ATTAINS assessment unit).

Each ATTAINS assessment reporting unit will be designated as having the aquatic life use and sub-use status according to the appropriate CBP segment/Aquatic life sub-designated use assessment. The rules to be applied are:

- a) Open water designated extends from “shoreline to shoreline” within each CBP segment and thus all ATTAINS reporting units located within each CBP segment are reported as having “open water” aquatic life use attainment consistent with the CBP segment attainment of open water criteria.
- b) Deep water and deep channel designated use spatial boundaries within each CBP segment are spatially constrained as smaller areas within the larger CBP segments (see *Technical Support Document for Identification of Chesapeake Bay Designated Uses and Attainability 2004 Addendum*. Chesapeake Bay Program Office, Annapolis, Maryland.) Thus the deep water or deep channel designated use status for each CBP segment will apply only to ATTAINS reporting units which contain a “deep water” designated use area. The two-dimensional (i.e. square miles) size of each CBP segment encompassing the impaired deep water use will be reported as the actual impaired area in ATTAINS, unless only a very small portion of deep water/deep channel is present in the segment.
- c) The Shallow Water Submerged Aquatic vegetation designated use applies only out to a maximum of 2 meter contour. Each ATTAINS unit having this sub-use present in some portion will designate the sub-use as attained or not. However, the actual size of the impaired use will be tracked outside of the ATTAINS system and reported in the Integrated Report as being only the size of area within the two meter contour.
- d) The general standard aquatic life use status of the ATTAINS unit will be assigned to the “worst case” status of aquatic life sub-use within that ATTAINS assessment unit (e.g. an ATTAINS reporting unit containing an open water use which meets its associated criteria and a deep water use which fails its associated criteria will be categorized as failing the general aquatic life use). Other criteria applicable to the general standard for aquatic life use (e.g. for benthic communities, toxics, or “weight of evidence” etc...) will also determine the overall aquatic life use attainment. If the general aquatic life use is impaired only due to a smaller area of aquatic life sub-use, then only the area (i.e. square mileage) of the sub-use is reported as impaired for general aquatic life use.

### **Impact of Chesapeake Bay TMDL**

The Chesapeake Bay TMDL, developed by EPA in cooperation with Bay state partners, was approved by EPA on December 31, 2010. This TMDL focuses on reductions to nutrients (nitrogen and phosphorus) and sediment inputs into Bay waters (including major tributaries). Improvements in DO and water clarity, which are indicators for the Aquatic Life and Shallow Water Submerged Aquatic Vegetation designated uses, are anticipated after the implementation of this TMDL.

All Bay waters that are on the 303(d) list for dissolved oxygen, SAV, and chlorophyll *a* impairments should be in Category 4, and any waters newly impaired for these parameters should also get this designation. Chesapeake Bay and tidal tributary assessment units, as described in [9 VAC 25-260-185](#) (d), that were listed for dissolved oxygen by EPA in the 1999 consent decree will continue to remain in Category 4 until all applicable criteria are attained (e.g. any 7-day mean or instantaneous criteria must be assessed and attained as well as the 30-day criteria).

Category 4D should be used to classify those waters listed for dissolved oxygen by EPA that are found to be meeting all *assessed* dissolved oxygen criteria. For instance, if a water meets the 30-day mean criterion for the Open Water sub-use, but the 7-day mean and instantaneous minimum criteria were not assessed, the Open Water sub-use should be categorized as “4D” in the Assessment Database. In the absence of other aquatic life use impairments (pH, benthics, etc.), the aquatic life use for this water would be assessed as “4D” as well. Refer to Part III for the full description of Category 4D.

## 5.4 *Biological Assessments*

Biological monitoring of streams and rivers using benthic macroinvertebrates is an integral component of the water quality monitoring program in the Commonwealth of Virginia. Biological monitoring allows the Virginia DEQ to assess the ecological condition of streams and rivers. Benthic macroinvertebrate surveys are used to determine if the waterbodies meet their designated aquatic life uses.

### **The Virginia Stream Condition Index (VSCI)**

In 2000, the United States Environmental Protection Agency (USEPA) contracted TetraTech to develop a multi-metric macroinvertebrate index for Virginia. This index contains eight core metrics that when calculated into one number is known as the Virginia Stream Condition Index (VSCI). TetraTech developed the VSCI using Virginia's existing biomonitoring database, which contained a significant amount of upstream (reference) control sites for use with the USEPA's Rapid Bioassessment Protocols.

Using an independent probabilistic database (sample n=350) with data collected from 2001-2004, Virginia has validated the VSCI using a spatially diverse (ecoregionally and stream size) data set free of pseudoreplication. These probabilistic data sets have allowed DEQ to narrow data gaps and test the proposed VSCI against many classification variables, which include season, stream size, ecoregion, bioregion, river basin, regional office, and sampling technique. The VSCI validation study was designed to incorporate suggestions provided through public comment from the Academic Advisory Committee (AAC), the USEPA and the regulated community.

The validation study using probabilistic biological data has confirmed that the VSCI works well to discriminate between sites with acceptable water quality and habitat versus sites with degraded water quality and habitat. A VSCI impairment threshold score of 60 was determined from statistical analyses of the original TetraTech report and the DEQ validation study. The VSCI validation study and the aquatic life use assessment guidance using the VSCI was reviewed by the USEPA. The validation study "Using Probabilistic Monitoring Data to Validate the Non-Coastal Virginia Stream Condition Index" can be found on DEQ's Probabilistic Monitoring website.

### **The Virginia Coastal Plain Macroinvertebrate Index (VCPMI)**

In the late 1990s, the United States Environmental Protection Agency (USEPA) coordinated a six-state monitoring effort to develop a multi-metric macroinvertebrate index that included Virginia's coastal plain. This index contained five metrics that when calculated into one number is known as the Coastal Plain Macroinvertebrate Index (CPMI). This index was adopted by DEQ in the early 2000's to make aquatic life use impairment determinations in the coastal plain of Virginia. Virginia biologists recommended validation of the index and initiated a special study.

Over the past decade DEQ compiled a new database of coastal plain macroinvertebrate data, which includes significantly more Virginia reference samples than the original CPMI study. Virginia has created the new VCPMI using a spatially diverse (ecoregionally and stream size) dataset free of pseudoreplication. The VCPMI replaces metrics that did not work well in Virginia's coastal plain and has correctly calibrated each metric's best standard values. The VCPMI study has confirmed that the VCPMI works well to discriminate between sites with acceptable water quality and habitat versus sites with degraded water quality and habitat. The impairment threshold score of 40 was determined from statistical analyses conducted during the VCPMI study. The VCPMI study and the aquatic life use assessment guidance using the VCPMI was reviewed by the USEPA.



The VCPMI technical report, “The Virginia Coastal Plain Macroinvertebrate Index”, can be found on the DEQ’s Probabilistic Monitoring website.

### **Free-flowing Aquatic Life Use Determination**

The DEQ uses the VSCI for non-coastal streams for biological assessment as well as the Virginia Coastal Plain Macroinvertebrate Index (VCPMI) for coastal plain streams. Assessment rankings, based on a single VSCI or VCPMI bioassessment, are the result of the data evaluation and reduction of numerous measurements and observations conducted during the biomonitoring survey. Bioassessment measures the response of the biological community to all perturbations it has experienced. A single, properly conducted VSCI or VCPMI bioassessment is not a “single data-point” analogous to a single dissolved oxygen (DO) measurement or bacteria sample. Non-coastal streams with VSCI scores  $\geq 60$  or coastal plain streams with VCPMI scores  $\geq 40$  will be assessed as “fully supporting for aquatic life use”. VSCI scores  $< 60$  and CPMI scores  $< 40$  will result in streams being listed as “impaired”.

If the biologist has observed natural conditions, such as recent drought or flooding, etc., that could be responsible for a ranking below the impairment threshold, they should note the lack of confidence in the survey and the stream will be listed as “insufficient information” if no other samples are available until further analysis can be conducted.

The regional biologists should review the biological assessments for the assessment cycle and they should make a final biological assessment ranking based on these data. Since biomonitoring surveys are records of the condition of the community at the time of the survey, the most recent bioassessment should be the most accurate indicator of stream ecological health. An attempt to average the data would weaken the ability to accurately predict current conditions. In cases where biological assessment rankings fall above and below the impairment threshold over multiple sampling events, more weight should be given to the most recent bioassessment. In cases where only one biomonitoring survey was conducted, a stream may be assessed for aquatic life use based on a single VSCI or VCPMI score. A standardized fact sheet, as found in Appendix B of this manual, has been developed to help the regional biologists review and assess the data for the assessment cycle. The fact sheet allows for consideration of supplemental information about the watershed that is important in making the final assessment decision.

### **Estuarine Aquatic Life Assessment**

In cooperation with EPA Region III and the State of Maryland, DEQ has developed an assessment methodology for estuarine benthic community biological (B-IBI) data. This methodology assures Bay-wide consistency in determinations of estuarine benthic impairments and requires a sample size  $\geq 10$  for statistical purposes. In order to assist with meeting the sample size requirement, a six-year data window is used. This corresponds with the data window used for the assessment of other non-Chesapeake Bay criteria data.

The methodology incorporates uncertainty in the reference condition and is based on the confidence limit and bootstrap simulation concept described in Alden et al. (2002). Bootstrap simulation (Efron and Tibshirani 1998) will be applied to incorporate uncertainty in reference conditions as well as sampling variability in the assessment data. For each habitat, a threshold based on percentiles in an unimpaired reference data set will be applied (i.e. 5<sup>th</sup> percentile). This threshold is not intended to serve as criteria for classifying individual B-IBI scores, rather it will be used to categorize the segment as impaired or not based on the proportion of samples below the threshold and the variance associated with this estimate.

The impairment assessment for each segment is based on the proportion of samples below the threshold with the variance in this proportion estimated by simulation. In each simulation run, a subset of the reference “unimpaired” data for each habitat is selected at random, and the threshold is determined (i.e., the B-IBI score at the 5<sup>th</sup> percentile of the un-impaired dataset). A random subset of the assessment data is compared to the threshold value to estimate the proportion of sites below the threshold. By repeating this process over and over again (2000 runs) we estimate the variance in the proportion of sites below the threshold from the bootstrap estimates. For this analysis, it is assumed that each reference ‘un-impaired’ data set (by habitat) is a representative sample from a “super population” of reference sites.

The assessment result for each benthic segment (i.e. % of area with IBI score below 5<sup>th</sup> percentile threshold) is then statistically compared ( $p < 0.05$ ) with the percentage that would be expected even if the segment is unimpaired. This percentage under “un-impaired” conditions is assumed to be 5%.

A benthic segment will be classified as having insufficient information (Category 3B) when the number of sites sampled during the six-year data window is less than 10. A segment will also be classified as Category 3B when the analysis suggests non-impairment but the difference between the upper and lower 95% confidence limits equals or exceeds 0.5 and the average BIBI score is less than 2.7.

In addition to an assessment of impairment, a discriminant analysis tool (benthic diagnostic tool) is under development and could be used to identify sources of stress affecting benthic community condition in the Chesapeake Bay (Dauer et al. 2002). The results could distinguish stress due to contaminants versus stress due to other factors (e.g., low dissolved oxygen, or unknown). This tool could be used to identify which impaired segments have high probability of sediment contamination. Separately from the discriminant tool, the B-IBI metric scoring could also be used to identify (1) insufficient abundance patterns consistent with a low dissolved oxygen effect and (2) excessive abundance patterns consistent with eutrophication effects in the absence of low dissolved oxygen events. The combined use of these causal analyses could be used to assign causes for benthic impairments to either 1) Sediment chemical contaminants 2) Low dissolved oxygen 3) Eutrophication or 4) Unknown.

The spatial assessment unit for determining attainment of the general standard for aquatic life use using benthic community data will be the same as used in the 2018 assessment report. These criteria assessment units are described in “*Chesapeake Bay Program Analytical Segmentation Scheme-Revisions, Decisions and Rationales: 1983 -2003*, CBP/TRS 268/04. Chesapeake Bay Program, Annapolis, Maryland” with the additional caveat that minor tidal tributaries are considered separate benthic assessment segments.

Assignment of aquatic life use status, as determined by benthic assessments to ATTAINS reporting waterbodies, will be the same as described previously for the Bay criteria assessments found in Section 5.3. Each ATTAINS reporting unit will be assigned the general aquatic life use status of the benthic assessment segment in which it is geographically located.

#### References:

- Alden, R.W. III. 1992. Uncertainty and sediment quality assessments: Confidence limits for the Triad. *Environmental Toxicology and Chemistry* 11:645-651.
- Alden, R.W. III, D.M. Dauer, J.A. Ranasinghe, L.C. Scott, and R.J. Llansó. 2002. Statistical verification of the Chesapeake Bay Benthic Index of Biotic Integrity. *Environmetrics* 13:473-498.
- Dauer, D.M., M.F. Lane, and R.J. Llansó. 2002. Development of diagnostic approaches to determine sources of anthropogenic stress affecting benthic community condition in the Chesapeake Bay.

- Report submitted to the USEPA Chesapeake Bay Program Office, Annapolis, Maryland, by Old Dominion University Department of Biological Sciences, Norfolk, Virginia. 65 pp.
- Efron, B. and R. Tibshirani. 1998. *An Introduction to the Bootstrap*. Chapman & Hall/CRC.
- Llansó, R.J., J.H. Vølstad, and D.M. Dauer. 2003. *Decision Process for Identification of Estuarine Benthic Impairments*. Final Report submitted to Maryland Department of Natural Resources, Tidewater Ecosystem Assessments, Annapolis, Maryland, by Versar, Inc., Columbia, Maryland.

## 5.5 *Toxics Assessment*

### **Fish Consumption Use**

The Fish Tissue Monitoring Program (FTM) collects fish tissue samples from designated monitoring stations for contaminant analysis. FTM staff provide data to water quality assessment staff for each sampling year in the assessment data window. Older fish tissue data may be included in assessment determinations when deemed appropriate. Fish tissue samples are assessed according to Section 5.2.3 of this manual and the protocols outlined below.

Fish tissue samples are collected via protocols described in the Quality Assurance/Quality Control Project Plan for the Fish Tissue and Sediment Contaminants Monitoring Program (August 1998). Fish tissue data collected at stations during routine monitoring throughout Virginia represent Tier 1 monitoring data. Tier 1 sampling stations may include freshwater, brackish or saltwater locations selected on a rotational river basin approach among the major river basins or sub-basins in Virginia. The samples collected at each site are dependent on what species are available. Collections at freshwater stations usually include 3-6 tissue samples, either as single species edible filets or composite tissue samples (usually 5-10 individuals of the same species per composite) consisting of fish species that are typically consumed by humans. Samples usually include at least one bottom feeder (e.g. catfish), which may be highly exposed to chemically contaminated sediments compared to other species, and two to four upper and middle trophic level feeders (e.g. bass and sunfish species, respectively.), which may be exposed to chemical contaminants via biomagnification. Tier 1 monitoring data are meant to identify sites where concentrations of contaminants in the edible portions of commonly consumed fish indicate a potential health risk to humans.

If Tier 1 results reveal potential problems, a more intensive Tier 2 study is initiated by the FTM staff to determine the magnitude, geographical extent, and potential sources of contamination in the fish. The need for a Tier 2 study takes into consideration the severity of the potential concern and is initiated as soon after the discovery of a potential problem as resources allow. Generally, if additional information is requested by the Virginia Department of Health (VDH) for determining the need for fish consumption advisories, a follow-up monitoring effort is initiated the year after the discovery of the potential problem. If limited resources prevent this, the water body will be sampled more intensely as soon as resources allow and/or during the next scheduled monitoring event in the affected river basin.

Currently, most fish tissue monitoring is focused on the development of PCB TMDLs throughout the Commonwealth.

Analytical results for fish tissue are expressed in wet-weight and are compared to WQS Tissue Values (TVs) and Tissue Screening Values (TSVs) for the toxic pollutants using EPA risk assessment techniques for non-carcinogen and carcinogen effects. WQS human health calculations use the 10<sup>-5</sup> risk level adopted

by the State Water Control Board and exposure factors<sup>6</sup> and relative source contribution factors recommended by EPA for the general U.S. population. These same values were used to calculate the human health water quality criteria found in [9 VAC 25-260-140 B](#). Also included in the calculation are toxicological data pertinent to human health effects. A reference dose (RfD) is used for non-carcinogen toxic effects and a cancer oral slope factor is used for carcinogen effects. TVs are based on the same toxicological data (and body weight, fish consumption, RfD or cancer risk level, and relative source contribution factors) that form the basis for the water quality criteria listed in [9 VAC 25-260-140 B](#), under the column labeled "Human Health, All Other Surface Waters". These water quality criteria are water column concentrations that are based on a specific fish tissue concentration, which were calculated to represent a safe or acceptable minimal human health risk level. The water quality criteria are designed to prevent the fish from bioconcentrating the toxic contaminants to levels greater than these fish tissue concentrations. The TV concentrations listed in Appendix E-1 represent the same fish tissue concentrations that are the basis for the water quality criteria listed in [9 VAC 25-260-140 B](#) and may be considered the fish tissue concentration equivalent of those water quality criteria. Appendix E-1 contains TVs for all chemicals for which Virginia has adopted water quality criteria. However, many of the chemicals listed in Appendix E-1 do not bioaccumulate and are not often found in fish tissue and have been included for completeness. All TVs are rounded to two significant digits.

Appendix E-2 also lists TSVs for additional toxic chemicals for which Virginia has not adopted human health water quality criteria that are designed to prevent excessive bioaccumulation in fish tissue (those criteria listed under " Human Health, All Other Surface Waters" in [9 VAC 25-260-140 B](#)). It includes chemicals recommended for monitoring by EPA or of special interest to DEQ as well as some chemicals that are based on recent changes to toxicological data and/or exposure assumptions that are different from those used to calculate the water quality criteria found in [9 VAC 25-260-140 B](#). The TSVs are updated using available data from the EPA IRIS database and/or recommendations from EPA or VDH before each assessment effort.

If there are no WQS TV exceedances in fish tissue samples in the assessment window, the water body should be assessed as fully supporting for the fish consumption use. If a fish tissue sample exceeds a single WQS TV or exceeds a TSV in the assessment window, the water body should be assessed as insufficient information and prioritized for follow-up monitoring. If the TV for the same toxic pollutant is exceeded in two or more samples from the same site in the assessment window, the water is considered impaired. Data from all Tier 1 and Tier 2 monitoring studies are evaluated by DEQ as well as provided to the VDH for their consideration of the need for establishing fish consumption advisories. DEQ and VDH have signed a Memorandum of Agreement (MOA) that describes how the agencies exchange information regarding the results of all Tier 1 and Tier 2 fish tissue monitoring. If VDH issues a fishing ban or advisory, limiting consumption, the segment should be designated impaired for fish consumption use based on the advisory. The results of the Tier 2 study should be clearly communicated in the Integrated Report narrative.

Additionally, DEQ uses the Virginia WQS for human health in surface waters to assess the fish consumption use in public water supplies as well as all other surface waters ([9 VAC 25-260-140 B](#)). If the median of two or more water samples from the same site collected in the assessment window exceed the WQS, the water is considered impaired. If the median of multiple water samples does not exceed the WQS in the assessment window, the water body should be assessed as fully supporting for the fish consumption use. If a site is represented by only one toxic pollutant sample in the assessment window and it exceeds the

---

<sup>6</sup> All fish tissue values in Appendix E-1 and E-2 assume the average body weight of 80 kg and a fish consumption rate of 22.0 grams/day.

WQS, the water body should be assessed as insufficient information and prioritized for follow-up monitoring.

In waters not designated as PWS, the criteria in the “All Other Surface Waters” column of [9 VAC 25-260-140B](#) should be used to assess the fish consumption use.

### **Public Water Supply Use**

DEQ uses the Virginia WQS for human health in public water supplies ([9 VAC 25-260-140 B](#)) to assess the public water supply use in surface waters designated as PWS per [9 VAC 25-260-380](#). The applicable criteria should be assessed throughout the extent of the designated areas. Taste and odor criteria to maintain acceptable taste, odor or aesthetic quality of drinking water apply at the drinking water intake.

Virginia will declare waters impaired for the public water supply use if a PWS WQS is exceeded by the median of two or more samples from the same site in the assessment window. If there are no PWS WQS exceedances in the assessment window in a dataset with 2 or more samples, the water body should be assessed as fully supporting for the public water supply use. If a single toxic pollutant sample exceeds a PWS WQS in the assessment window the water body should be assessed as insufficient information and prioritized for follow-up monitoring.

If an exceedance of a taste and odor criteria is observed within the PWS designated area, follow up monitoring at the closest drinking water intake should occur as regional office resources allow to make an assessment determination.

In waters designated as PWS, the criteria in the PWS column of [9 VAC 25-260-140B](#) should be used to assess both fish consumption and PWS uses, except for pollutants which do not have a criterion in the “All Other Surface Waters” column. For these pollutants, exceedances of the PWS criteria only indicate impairment of the PWS use.

### **Aquatic Life/Wildlife Use**

#### *Sediment Analysis*

Similar to the sediment monitoring and analysis conducted by FTM, the regional offices will assess the AWQM sediment data. For freshwater sediments above the fall-line and in tidal fresh zones, as described in the WQS, the consensus-based Probable Effects Concentrations (PEC; MacDonald et al. 2000) should be applied. Estuarine sediment contaminant data collected during scheduled AWQM monitoring should be compared to National Oceanic and Atmospheric Administration (NOAA *Screening Quick Reference Tables* (SQuiRT) Tables 1999) for effects-range-median (ER-M) SVs for sediment. Transition zones should be assessed against the more stringent of the two screening values. One or more exceedances of an ER-M/PEC value results in an insufficient information status for aquatic life use support. In these cases, additional biological monitoring should be scheduled to assess actual aquatic life use support. For National Coastal Assessment, a “weight of evidence” approach using sediment toxicity and sediment chemistry will be used to determine aquatic life designated use. See Appendix G for additional information. All metals contaminant screening values found in Appendix F have been converted to parts per million (ppm) for consistency.

## *Water Column Analysis*

For overall freshwater toxics evaluation, DEQ uses the Virginia WQS for human health in surface waters, other than public water supplies ([9 VAC 25-260-140 B](#)). These same values are used to assess the fish consumption use in public water supplies as well as all other surface waters. (Please note the criteria for human health in public water supplies will be used to assess the drinking water use in PWSs only). When assessing the aquatic life and wildlife use support for toxic contaminants, compliance should be based on meeting the aquatic life WQS found in [9 VAC 25-260-140 B](#). See [Section 5.2](#) for additional information.

Virginia will declare waters impaired for aquatic life use if 1) an acute criterion is exceeded two or more times in most of the three-year periods, based on either grab samples or samples collected with a 30-day semi-permeable membrane device (SPMD) or if 2) a chronic criterion is exceeded two or more times in most of the three-year periods for which there are a sufficient number of samples within the assessment window, based on either multiple grab samples collected within two separate four-day periods or multiple samples collected with a 30-day semi-permeable membrane device (SPMD)". The three-year period should be represented by three consecutive calendar years (i.e., samples are taken between January 1 in the first year to December 31 in the third year).

## *Weight-of-Evidence Assessment in Estuarine Waters*

The "Weight-of-Evidence" (WOE) approach that DEQ currently uses for its general evaluation and assessment of the designated Aquatic Life Use (ALU) for estuarine benthic communities has evolved from a previously more limited application of the "Sediment Quality Triad" concept (SQT – Appendix G Figure 1). The SQT concept was originally conceived and applied for the evaluation of the presence and effects of toxic contaminants in marine sediments (Long and Chapman, 1985). It was further applied by Chapman et al. (1986, 1987), and has continued to be one of the preferred approaches for the evaluation of toxics in marine and estuarine benthic environments (Chapman, 1992; Chapman et al., 1997; McGee et al., 2001). The Interstate Chesapeake Bay Program (CBP) employed SQT evaluations along with other methods to produce a Bay-wide toxics characterization in 1999 (US EPA, 1999) that identified (1) "Regions of Concern – areas with probable adverse effects," (2) "Areas of Emphasis – areas with potential adverse effects," (3) "Areas with Low Probability for Adverse Effects", and (4) "Areas with Insufficient or Inconclusive Data" relative to toxics contamination in Bay waters. Maps of more recent characterizations (2006, 2008, and 2009) can be found at: <http://www.chesapeakebay.net/maps.aspx?menuitem=15230>.

See Appendix G for more details on the WOE approach and the SQT, as well as references cited here.

## *5.6 Naturally Low DO and pH Evaluation in Swamp Waters*

The natural conditions process is currently under revision. The agency is working in partnership with an Academic Advisory Committee to develop a new classification process for swamp waters. This new process will differ from past protocols in that the steps of classifying natural conditions and evaluating anthropogenic disturbance will be separated. This will allow for the correct assignment of waters to Class VII regardless of the level of anthropogenic disturbance in the system and surrounding watershed.

## *5.7 Lakes/Reservoirs Assessment*

The current agency guidance on the monitoring of targeted lakes and reservoirs is found in the Department Guidance Memo No. 09-2005 "Monitoring of Lakes and Reservoirs" (December 2020). The guidance provides [Final Guidance for 2024 IR Assessment Methodology](#)

information on how DEQ prioritizes the many lakes and reservoirs in the Commonwealth for monitoring. This prioritization allows the Department to focus on the most important lakes as they relate to designated uses. The most recent updates to this guidance document removed all assessment methodology references. Section 5.7 provides guidance on how to conduct assessments on all lakes and reservoirs, whether they are considered ‘significant’ or not.

GM09-2005 defines “significant lakes and reservoirs”:

*“A significant lake/reservoir is defined as: a publicly accessible lake/reservoir that is a public water supply and/or 100 acres or more in size **and** is included in 9 VAC 25-260-187 list of reservoirs with nutrient criteria.”*

A list of the current “significant lakes” is included in Appendix H of this guidance. Since 2007, these are the man-made lakes and reservoirs identified under the nutrient standards for lakes and reservoirs in [9 VAC 25-260-187](#), and the two natural lakes, Mountain Lake and Lake Drummond, which have been assigned special standards for nutrients ([9 VAC 25-260-310](#)).

Publicly accessible means direct access to the water from public property during normal work hours.

The significant lakes designation includes the federally owned lakes which meet these criteria, but all other federally owned lakes would be excluded from the agency lakes monitoring program. There are additional lakes/reservoirs that should also be considered, as resources allow, for monitoring and assessment that are not listed in 9 VAC 25-260-187 but do meet one of the other two characteristics above. Although nutrient criteria do not apply to these, other criteria should be assessed as available data will allow.

At least one of these two criteria must be met for a lake or reservoir to be assessed as impaired:


1. A lake/reservoir has exceedances of numerical WQS or assessment threshold published in this guidance, as observed during multiple sampling events (see Table 4), or
2. a VDH Advisory or Closure during the assessment window occurred in the lake/reservoir.

### Determination of Thermal Stratification

Determination of thermal stratification is needed to determine how to apply DO and pH criteria and trophic state indices (TSI) where applicable. During periods of thermal stratification of significant lakes/reservoirs, attainment of the minimum dissolved oxygen and pH criteria are only assessed in the epilimnion at monitoring stations where stratification is observed. Dissolved oxygen and pH criteria do not apply below the epilimnion during thermal stratification. Determination of thermal stratification in non-significant lakes/reservoirs is needed to determine if nutrient impacts through the TSI evaluation is needed. All dissolved oxygen and pH values recorded during sampling events on non-significant lakes/reservoirs should be assessed.

The usual procedure to determine thermal stratification is to plot temperature vs. depth to determine where a slope occurs in the temperature/dissolved oxygen profile measurement. In deeper lakes or reservoirs, it may be possible to use the classical textbook definition of lake stratification developed for natural lakes if the temperature profile meets the following definition for a thermocline: the temperature decrease reaches 1 degree C or more per meter of descent. The first instance of a 1 degree C change per meter of descent from the surface of the lake or reservoir should represent the bottom of the epilimnion. Under the classical procedure, if the bottom of the epilimnion cannot be delineated (*i.e.* the temperature decrease is less than 1 degree C for each meter of descent), then it is not considered to be thermally stratified. However, many lakes/reservoirs are

shallow, often less than twenty feet deep, and exhibit stratification but do not meet the “classical” 1 degree C per meter of descent definition of stratification. In such situations, staff will need to use best professional judgment to determine whether a lake/reservoir is exhibiting stratification. Examples of annual temperature cycles in stratified lakes are given in Figure 1.



(WDNR)

(University of Illinois)

Figure 1. Annual temperature cycles in stratified lakes, Wisconsin DNR (WDNR) and University of Illinois

## Assessment of Significant Lakes/Reservoirs

### Nutrient Evaluation

Nutrients including chlorophyll *a* and total phosphorus are assessed for aquatic life use. Total phosphorus is assessed in addition to chlorophyll *a* if there is documented use of algaecides any time during the monitoring period. Observations regarding nuisance algal, plant growth, or discolored water are assessed using the general standard; the nutrient criteria for lakes and reservoirs did not modify these existing criteria.

The nutrient criteria for the man-made lakes and reservoirs listed in [9 VAC 25-260-187](#) of the WQS only apply in the top 1 meter of the lacustrine zone. If chlorophyll *a* or total phosphorus are collected outside the lacustrine zone, the data will not be used to assess of the aquatic life use in other areas of the lake/reservoir. “Lacustrine” means the zone within a lake or reservoir that corresponds to non-flowing lake-like conditions within reservoirs that are generally deeper than 3 meters (10 feet). In small, shallow lakes and reservoirs where the classic lacustrine, transition and riverine zones are not pronounced, the lacustrine zone and nutrient criteria in 9 VAC 25-260-187 apply in a representative area of the water body. The other two zones within a deeper reservoir are riverine (flowing, river-like conditions) and transitional (transition from river to lake conditions). [Guidance Memo 09-2005](#) provides general characteristics of the riverine, transitional and lacustrine zones in lakes and reservoirs. The document also provides guidelines for identifying these zones within a lake/reservoir for assessment purposes. Assessment staff should consult with regional monitoring staff to confirm the location of lake monitoring stations as it relates to these zones.

This assessment procedure for nutrients in significant lakes and reservoirs replaces the combined TP/DO TSI approach used in previous assessments for nutrient assessment related to assessing natural low DO conditions. However, the TSI approach will continue to be used to determine natural conditions for non-significant lakes or reservoirs if DO problems have been documented.

Final Guidance for 2024 IR  
*Assessment Methodology*



A master copy of owner and/or manager contact information of significant lakes will be maintained by DEQ assessment staff. A request will be sent to these contacts at the beginning of each assessment cycle to determine the use of algaecides during the lake monitoring period. Any DEQ monitoring staff observations of algaecide applications during their monitoring runs on a lake or reservoir shall also be confirmed with these contacts. (The intent is to use both chlorophyll *a* and total phosphorus when algaecides are applied within any zone of the reservoir.)

The 9 VAC 25-260-187 states, “The 90th percentile of chlorophyll data collected at one meter or less within the lacustrine portion of the man-made lake or reservoir between April 1 and October 31 (considered a lake monitoring year) shall not exceed the chlorophyll *a* criterion for that waterbody in *each* of the two most recent monitoring years within the assessment window. For a waterbody that received algaecide treatment, the median of the total phosphorus data collected at one meter or less within the lacustrine portion of the man-made lake or reservoir between April 1 and October 31 shall not exceed the total phosphorus criterion in each of the two most recent years that total phosphorus data are available.” The aquatic life (fishery) use of any lake assessment unit is considered impaired for nutrients if the criterion for either chlorophyll *a* or total phosphorus is exceeded at a station or pooled stations in that unit in each of the two most recent monitoring years within the assessment window.

For lake or reservoir assessment units with multiple stations or multiple samples collected in a month, chlorophyll-*a* and total phosphorous data should be pooled by calculating the median of same-month observations from April to October of any given year in a relatively homogenous unit. The 90th percentile of monthly chlorophyll-*a* medians is the value used to compare to the chlorophyll-*a* criterion for a particular lake/reservoir, consistent with the criterion development. The median of monthly TP medians should be used to assess against the applicable total phosphorous criterion.

Assessment is based on the two most recent monitoring years that data are available within the assessment window. Each year must have data for at least six of the seven months of required monitoring to be considered a valid year. If sampled for fewer than six of the seven sampling months within a calendar year, the lake or reservoir is placed in Category 3 (insufficient data) for assessment purposes and sampling is conducted in the next calendar year, based on regional prioritization and available resources. A third sampling year may be needed if the two previous years result in differing assessment statuses. This monitoring should be done as soon as resources allow.

*Assessment for aquatic life (fishery) use for nutrients in the two natural lakes:*

Assessments of the two natural lakes in the special standards section will follow the guidelines above for chlorophyll *a* and total phosphorus except that orthophosphate-P rather than total phosphorus applies to Mountain Lake.

#### Process for confirmation of use impairments when nutrient criteria are exceeded

9 VAC 25-260-187.C. includes a process for confirmation of the aquatic life (fishery) use impairments via a consultation with the Virginia Department of Wildlife Resources (DWR, formerly Virginia Department of Game and Inland Fisheries) when the nutrient criteria in 9 VAC 25-260-187.B are exceeded. Where the nutrient criteria are exceeded but the designated uses of the water body are being attained, the water will still be considered impaired in accordance with § 62.1-44.19.5 of the Code of Virginia until site-specific criteria are

adopted and become effective in order to remove the lake/reservoir from the impaired waters list. Site-specific criteria would be developed using the process described in § 62.1-44.15 of the Code of Virginia.

*Consultation Process with DWR:* When DEQ determines that the applicable nutrient criteria in Section 187 of the water quality standards regulation for a specific lake/reservoir are exceeded, the DEQ central office water quality standards program coordinator, or his/her designee, may contact the DWR Regional Fisheries Manager regarding the status of the fishery in determining whether or not the designated use for that water body is currently being attained. Since the nutrient criteria for lakes/reservoirs were developed using water chemistry data from those lakes/reservoirs where DWR biologists rated the fishery as good or excellent, documentation should support this level of fishery. Appropriate documentation for confirmation of maintenance of the designated fishery use would include the most recent information available on:

- catch per unit effort of specific size classes of managed fish populations
- population size structure
- observations or records regarding changes in fishing use or
- general trends in community structure.

If the above documentation requirements are met for DWR owned and managed lakes during the consultation process, the nutrient impairment observed during the assessment cycle should be classified as Category 5C. Any other impairments in these lakes should be classified as Category 5A.

### Dissolved Oxygen Evaluation

The dissolved oxygen criteria are based on the appropriate criteria established for that class of waters in Section [9 VAC 25-260-50](#). Dissolved oxygen (DO) information is used for assessment of aquatic life use.

- Assessment for aquatic life use of lakes and reservoirs for the dissolved oxygen criterion:

The 10.5% rule is applicable to assessments for the minimum dissolved oxygen criterion in all lakes and reservoirs year-round. For lakes/reservoirs listed in 9 VAC 25-260-187, dissolved oxygen samples taken for all months, at all stations within a given lake or reservoir, are assessed only in the epilimnion if the water body is thermally stratified. If not stratified, dissolved oxygen should be assessed throughout the water column. Typically, two or more exceedances and >10.5% exceedance of data collected at an individual station are required before a water body is listed as impaired for the minimum dissolved oxygen criterion (4 mg/l for most freshwater lakes and reservoirs) under § 62.1-44.19:5 and 7 of the Code of Virginia.

In most cases, a single monitoring station should represent a lake/reservoir assessment unit. In cases where there are multiple stations in an assessment unit and it is determined that the water in that unit is relatively homogenous and not influenced by tributary contribution, the independent station assessment results may be pooled to determine DO exceedance rates in that AU.

In cases where the applicable nutrient criteria are met for the man-made lakes/reservoirs listed in 9 VAC 25-260-187 but there are seasonal exceedances of the dissolved oxygen criterion due to fall overturn that result in the lake/reservoir assessment unit to be listed as impaired, a Category 4C rationale may be developed for EPA review. The rationale should include;

- maps of the lake/reservoir assessment unit and surrounding land-use to evaluate any potential sources in the watershed, and

- dissolved oxygen/temperature data profiles from at least two monitoring years in line with the Fall overturn phenomena.

In cases where the applicable nutrient criteria are not met, but there are seasonal exceedances of the dissolved oxygen criterion due to fall overturn that result in the lake/reservoir assessment unit to be listed as impaired, the lake or reservoir should be classified as Category 5C and recommended for a WQS review due to seasonal DO fluctuations.

### pH Evaluation

The pH criterion ranges are based on the appropriate criteria established for that class of waters in section [9 VAC 25-260-50](#). pH information is used for assessment of aquatic life use.

- Assessment for aquatic life use of lakes and reservoirs for the pH criterion range:

The 10.5% rule is applicable to assessments for the pH criterion range in all lakes and reservoirs year-round. For lakes/reservoirs listed in 9 VAC 25-260-187, pH samples taken for all months, at all stations within a given lake or reservoir, are assessed only in the epilimnion if the water body is thermally stratified. If not stratified, pH should be assessed throughout the water column. Typically, two or more exceedances and >10.5% exceedance of data collected at an individual station are required before a water body is listed as impaired for pH criterion range (6.0-9.0 for most freshwater lakes and reservoirs) under § 62.1-44.19:5 and 7 of the Code of Virginia.

In most cases, a single monitoring station should represent a lake/reservoir assessment unit. In cases where there are multiple stations in an assessment unit and it is determined that the water in that unit is relatively homogenous and not influenced by tributary contribution, the independent station assessment results may be pooled to determine pH exceedance rates in that AU.

In cases where the applicable nutrient criteria are met for the man-made lakes/reservoirs listed in 9 VAC 25-260-187 but the pH criterion range is not met, the lake or reservoir should be classified as Category 5C and recommended for a WQS review due to natural pH fluctuations. In lakes that are not listed in 9 VAC 25-260-187 and no nutrient problem is detected through TSI calculations, the waterbody would be listed as impaired (Category 5C), as well.

### **Assessment of Lakes/Reservoirs Not Listed in 9 VAC 25-260-187**

#### Nutrient Evaluation using Trophic State Index (TSI)

TSI equations for secchi depth (SD), chlorophyll *a* (CA), and total phosphorus (TP) will be calculated in stratified lakes using aggregated station data from mid-June through mid-September (at 0.3 m for TP and CA) and will be used to determine if DO problems in lakes and reservoirs not listed in 9 VAC 25-260-187 are natural (Category 4C).

A trophic state index value of 60 or greater for any one of the 3 indices will indicate that nutrient enrichment from anthropogenic sources are adversely interfering, directly or indirectly, with the designated uses. A TSI value of 60 corresponds to a CA concentration of 20 ug/l, a SD of 1 meter, and a TP concentration of 48 ug/l. The TSI equations:

$$TSI(SD) = 10(6 - (\ln SD / \ln 2))$$

*Final Guidance for 2024 IR  
Assessment Methodology*

$$TSI(CA) = 10(6 - ((2.04 - 0.68 \ln CA) / (\ln 2)))$$

$$TSI(TP) = 10(6 - ((\ln (48 / TP)) / (\ln 2)))$$

SD = meters

CA = ug/

TP = ug/l

The following rules apply:

1. Do not calculate a chlorophyll *a* TSI in lakes that are treated with algaecides.
2. The chlorophyll *a* TSI will normally be the preferred indicator in untreated lakes.
3. Assume that typical Virginia freshwater lakes and reservoirs are phosphorus limited.
4. Do not use the secchi depth index in the assessment if it is much larger than the CA and TP indices in the same assessment unit (this indicates prevalence of inorganic matter).
5. The appropriate TSIs should be calculated based on all summer sample data collected in the segment using the spreadsheet that has been developed for easier data processing.

For each monitoring station, if one or more of the TSIs  $\geq 60$ , the lake/reservoir not listed in 9 VAC 25-260-187 will be assessed as impaired partially due to one or more pollutants from anthropogenic sources. The assessment unit or entire lake/reservoir will be placed in Category 5A for TMDL development.

For each monitoring station, if each of the TSIs  $< 60$ , the lake/reservoir will be assessed as impaired due to pollution from natural sources and placed in Category 4C. A TMDL is not needed for the assessment unit represented by the monitoring station(s) and appropriate DO criteria will be developed for the hypolimnion.

If multiple stations are sampled on the lake/reservoir or within a lake/reservoir assessment unit, the individual TSI equations should be calculated at each station and then averaged (using a median or arithmetic mean) to determine the values for the waterbody.

**Table 3. Trophic status in ATTAINS**

<b>Trophic Index Trophic State</b>	<b>Carlson Trophic State Index</b>	<b>IR Category</b>
Hypereutrophic	80 – 100	5A
Eutrophic	60 – Less than 80	5A
Mesotrophic	40 – Less than 60	4C
Oligotrophic	0 – Less than 40	4C
Unknown	Insufficient Data	3A

### Dissolved Oxygen/pH Evaluation

Lakes/Reservoirs not listed in 9 VAC 25-260-187 should have all DO and pH samples collected during lake/reservoir sampling events assessed, regardless of thermal stratification determination. Two or more exceedances and  $>10.5\%$  exceedance of data collected at an individual station are required before a water body is listed as impaired for the minimum dissolved oxygen criterion (4 mg/l for most freshwater lakes and reservoirs) under § 62.1-44.19:5 and 7 of the Code of Virginia. Typically, two or more exceedances and  $>10.5\%$  exceedance of data collected at an individual station are required before a water body is listed as impaired for pH criterion range (6.0-9.0 for most freshwater lakes and reservoirs) under § 62.1-44.19:5 and 7 of the Code of Virginia.

In most cases, a single monitoring station should represent a lake/reservoir assessment unit. In cases where there are multiple stations in an assessment unit and it is determined that the water in that unit is relatively

### Final Guidance for 2024 IR

#### Assessment Methodology

homogenous and not influenced by tributary contribution, the independent station assessment results may be pooled to determine DO and pH exceedance rates in that AU.

### **Use of Citizen or Other External Data**

In order to use citizen data in assessments for nutrient impairments: 1) the collector must provide documentation that the data meet QA/QC requirements per [Virginia's Citizen Monitoring Methods Manual](#) for chlorophyll *a* and total phosphorus (orthophosphate-P for Mountain Lake), 2) the location of the sampling was within the lacustrine portion of the reservoir and outside the littoral (near shore) zone and 3) sampling corresponds with the lake monitoring year requirements.

Dissolved Oxygen and pH data must be collected throughout the water column and depth profile data (including temperature data) must meet the QA/QC requirements to be used for assessments of lakes/reservoirs. Surface only dissolved oxygen, pH and temperature samples will not be included in lake/reservoir assessment determinations.

### **Other Parameters Assessed in all Lakes/Reservoirs**

#### Bacteria Evaluation

The bacteria criteria are based on the appropriate criteria established for that class of waters in section [9VAC25-260-170](#). Bacteria information is used for assessment of the recreation use. Bacteria assessment methodologies are discussed in previous sections of this guidance and summarized in Table 4.

When individual stations (whether sampled by DEQ or non-agency/citizen groups) are located in the same assessment unit and they are sampled on the same day, the median of same-day bacteria measurements should be calculated.

#### Temperature Evaluation

Temperature data should be assessed throughout the water column using the “Percent Method” or 10.5% rule.

In most cases, a single monitoring station should represent a lake/reservoir assessment unit. In cases where there are multiple stations in an assessment unit and it is determined that the water in that unit is relatively homogenous and not influenced by tributary contribution, the independent station assessment results may be pooled to determine temperature exceedance rates in that AU.

#### Ammonia Evaluation

Ammonia is collected in the top 1 meter of lakes and reservoirs. Both chronic and acute water quality criteria are used to assess the aquatic life and wildlife uses. Calculations used to assess can be found in [9VAC25-260-155](#) as well as applicable aquatic life considerations. Two or more exceedances of the same acute or chronic toxic criteria in a 3-year period constitutes an impairment in a lake/reservoir assessment unit.

#### Nitrate Evaluation

Nitrate as N is collected in the top 1 meter of lakes and reservoirs to assess the public water supply use. If the median of two or more samples exceeds the Nitrate criteria within the assessment window it constitutes an impairment in a lake/reservoir assessment unit.

**Table 4. Lake/Reservoir Designated Use Assessment Summary**

Designated Use	Parameter	Lake/Station Type <sup>(2)</sup>	Depth	Monitoring Year	No. of Monitoring Years	Assessment	Pool Data?
Aquatic Life	Chlorophyll a	Significant Lacustrine	Collected in top 1m	Must have 6 out of 7 samples between Apr-Oct for valid monitoring year	2 most recent or 2 out of 3 if results conflict	90th percentile of all months within 1 monitoring year <a href="#">9 VAC 25-260-187</a>	Yes - calculate the median of same-month observations from April to October of any given year in a relatively homogenous lake AU, then assess.
	Total Phosphorus <sup>(1)</sup>	Significant Lacustrine (where there is documented use of algaecides)				Median of all months within 1 monitoring year <a href="#">9 VAC 25-260-187</a>	
	Trophic State Index (TSI) Secchi Depth, Chlorophyll a, Total Phosphorus	Non-Significant Stratified Lakes/Months Only	Chl a & TP collected in top 1m	Mid-June through Mid-September	All	TSI > 60 for any one of the 3 parameters will indicate impairment (Category 5)	Calculate TSI equations at each station and then average (using a median or arithmetic mean) to determine the values for the waterbody.
	Dissolved Oxygen	Significant All	If stratified: Epilimnion If not stratified: All depths	Year-round	All	Typically, two or more exceedances and >10.5% exceedance of total samples results in impairment	May pool exceedance rates for multiple stations in a relatively homogenous lake AU, not influenced by tributary contribution
		Non-significant All	All depths				
	pH	Significant All	If stratified: Epilimnion If not stratified: All depths	Year-round	All	Typically, two or more exceedances and >10.5% exceedance of total samples results in impairment	May pool exceedance rates for multiple stations in a relatively homogenous lake AU, not influenced by tributary contribution
		Non-significant All	All depths				
Temperature	All	All depths	Year-round	All	Typically, two or more exceedances and >10.5% exceedance of total samples results in impairment	May pool exceedance rates for multiple stations in a relatively homogenous lake AU, not influenced by tributary contribution	
Recreation	Bacteria (Freshwater-E.coli)	All	Surface Sample (0.3m)	All	All	n ≥ 10, one or more geometric mean exceedance(s) in a 90-day period or STV exceedance rate > 10% in a 90-day period results in impairment	Calculate the median of same-day bacteria measurements then assess, if within a relatively homogenous lake AU.
Wildlife	Ammonia	All	Surface Sample (0.3m)	All	All	Two or more exceedances of the same acute or chronic toxic criteria in a 3-year period	No
Public Water Supply	Nitrates	All	Surface Sample (0.3m)	All	All	The median of two or more samples exceeds the criteria in the assessment window	No

Notes:

- 1) Orthophosphate-P rather than total phosphorus applies to Mountain Lake for nutrient evaluation.
- 2) ‘Significant’ refers to lakes/reservoirs listed in 9 VAC 25-260-187 and the two natural lakes, Mountain Lake and Lake Drummond. ‘Non-significant’ refers to lakes/reservoirs not listed in 9 VAC 25-260-187.

## 5.8 Nonpoint Source (NPS) Assessment

Past versions of this guidance described a NPS Assessment process performed by the Virginia Department of Environmental Quality in collaboration with the Virginia Department of Conservation and Recreation (DCR). The assessment consisted of calculations of net loadings of the NPS pollutants - nitrogen, phosphorous, and sediment - per hydrologic unit as well as evaluations of NPS-related measures in these units. The watershed-based loadings are still calculated by DCR to support TMDL implementation project prioritization as part of the [Virginia Nonpoint Source \(NPS\) Pollution Management Program](#). Moving forward, the Nonpoint Source Assessment Chapter of the Integrated Report will summarize Virginia's coordination of nonpoint source pollution control programs based on annual reporting through [EPA's Grant Reporting and Tracking System](#).

## 5.9 Coastal Assessment

Virginia has 120 miles of Atlantic Ocean coastline and approximately 2,500 square miles of estuary. This resource has a prominent place in Virginia's history and culture. It is valued for its commercial fishing, wildlife, sporting, and recreational opportunities, as well as its commercial values in shipping and industry. In the 1970's adverse trends in water quality and living resources were noted and prompted creation of the Federal-Interstate Chesapeake Bay Program (CBP). The coastal assessment is conducted in the same manner as the estuarine assessments previously. Additionally, the federal BEACH program, implemented by VDH, has enabled the collection of recreational use data during the swimming season. Assessment of this data has been incorporated into the Integrated Report.

## 5.10 Wetlands Assessment

### Background

Impacts to tidal wetlands, including vegetated tidal wetlands and non-vegetated shoreline between mean low and mean high water, are regulated under the Virginia Tidal Wetlands Act (Title 28.2, Chapter 13 of the Code of Virginia) enacted in 1972 and revised in 1982. The Virginia Marine Resources Commission (VMRC) is the regulating authority for the tidal wetlands laws while localities in Tidewater Virginia have the option to regulate their own tidal wetlands through citizen Wetlands Boards with oversight from VMRC. The Virginia Department of Environmental Quality (DEQ) is responsible for providing Section 401 Certification of Clean Water Act for Section 404 federal permits for tidal and nontidal wetlands and water withdrawals, through the Virginia Water Protection Permit (VWPP) Program, first developed in 1992.

The VWP permit program is administered by DEQ's Office of Wetlands & Stream Protection, and derives its regulatory authority from both the Clean Water Act (§401) and State Water Control Law (§62.1-44.20 of the Code of Virginia), found at <http://water.epa.gov/lawsregs/guidance/wetlands/sec401.cfm> and <http://leg1.state.va.us/cgi-bin/legp504.exe?000+cod+62.1-44.15C20>, respectively

Before July 1, 2000, applicants seeking a Clean Water Act § 404 permit (<http://water.epa.gov/lawsregs/guidance/wetlands/sec404.cfm>) from the U.S. Army Corps of Engineers (the Corps) for the discharge of dredged or fill materials in wetlands or waters of the United States were also required to submit an application to DEQ for a permit or waiver under § 401 Certification. In 2000, Virginia passed a Nontidal Wetlands Act that amended Title 62.1 of the Code of Virginia relating to wetlands. The Nontidal Wetlands Act mandates that the Commonwealth implement a nontidal wetlands regulatory program to achieve no net loss of existing wetland acreage and function, and to develop voluntary and incentive based

### Final Guidance for 2024 IR Assessment Methodology

programs to achieve a net resource gain in wetlands. Amendments to the Virginia Water Protection (VWP) permit program, fully implemented in October 2001, provided additional state jurisdiction and required a state permit for the following activities in a wetland: excavation, filling or dumping, activities in a wetland that cause drainage or otherwise significantly alter or degrade existing wetland acreage or function, and permanent flooding or impounding.

Amendments to the VWPP program in July 2007 provided clarifications of state jurisdiction of small water withdrawals, incorporated several provisions of the Local and Regional Water Supply Planning regulation, and addressed water supply permitting and surface water withdrawal concerns. Further amendments to the VWPP program in December 2008 provided exclusion of certain in-stream fills for water supply on agricultural properties.

Further reducing duplication of permitting between State and Federal agencies while ensuring minimal individual and cumulative consequences to wetland and stream resources, the Corps issued a State Program General Permit (SPGP) and suspended a few of the Nationwide Permits. The projects must be below applicable wetland and stream impact thresholds and meet all other limitations and conditions of the SPGP. A type of Section 404 permit issued to the Commonwealth of Virginia. If a project meets the eligibility criteria and conditions of the permit, then coverage under the SPGP will be issued by DEQ in conjunction with a Virginia Water Protection permit.

22-SPGP-RCIR authorizes the discharge of dredged or fill material in non-tidal waters of the United States (WOTUS), including wetlands, associated with residential, commercial, institutional, and recreational development (RCIR) projects within the geographical limits of the Commonwealth of Virginia and under the regulatory jurisdiction of the U.S. Army Corps of Engineers, Norfolk District (Corps or Norfolk District). These projects must not cause the loss of greater than 1 acre of WOTUS, must have no more than minimal individual and cumulative impacts to WOTUS and must have avoided impacts to WOTUS to the greatest extent practicable.

22-SPGP-LT authorizes the discharge of dredged or fill material in non-tidal WOTUS, including wetlands, associated with linear transportation (LT) projects within the geographical limits of the Commonwealth of Virginia and under the regulatory jurisdiction of the Corps. These projects must not cause the loss of greater than 1/2 acre of WOTUS, must have no more than minimal individual and cumulative impacts to WOTUS and must have avoided impacts to WOTUS to the greatest extent practicable.

This SPGP is not available for use in the Commonwealth of Virginia subwatersheds shared with the State of Tennessee. These watersheds are identified in Appendix A of the 22-SPGP-RCIR and 22-SPGP-LT documents. Applicants may instead apply to the Corps of Engineers for Nationwide Permit or Regional General Permit verifications or a Standard Individual Permit. More information is available on DEQ's website at <https://www.deq.virginia.gov/permits-regulations/permits/water/wetlands-streams>.

The permit process for both tidal and nontidal wetland activities relies on a Joint Permit Application (JPA) which receives independent and concurrent review by local wetlands boards, VMRC, DEQ and the U.S. Army Corps of Engineers (Corps), as appropriate.

By statute and by regulation, Virginia adopted the same definition of wetlands as the federal definition, and requires that wetlands be defined in the field using the Corps' 1987 Manual. Specifically, wetlands are defined as "those areas that are inundated or saturated by surface or groundwater at a frequency and duration sufficient to support, and that under normal circumstances do support, a prevalence of vegetation typically adapted for life



in saturated soil conditions. Wetlands generally include swamps, marshes, bogs, and similar areas." Wetlands are part of state waters, which are defined as "all water, on the surface and under the ground, wholly or partially within or bordering the Commonwealth or within its jurisdiction, including wetlands." The Corps has created Regional Supplements in an effort to address regional wetland characteristics and improve the accuracy and efficiency of wetland-delineation procedures. The Regional Supplements that apply to Virginia are: U.S. Army Corps of Engineers (2010) "Interim Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Eastern Mountains and Piedmont Region," ERDC/EL TR-10-9, U.S. Army Engineer Research and Development Center, Vicksburg, MS and U.S. Army Corps of Engineers (2010). "Regional Supplement to the Corps of Engineers Wetland Delineation Manual - Atlantic and Gulf Coastal Plain Region (Version 2.0)," ERDC/EL TR-10-20, U.S. Army Engineer Research and Development Center, Vicksburg, MS.

Tidal wetlands are defined to include tidally influenced areas within Tidewater Virginia contiguous to mean low water extending landward to an elevation 1 1/2 times the mean tide range at a site and upon which is growing certain listed plant species. They also include "nonvegetated wetlands" which include unvegetated lands between mean low water and mean high water tides.

Section 62.1-44.15:21 of the Code of Virginia specifies that the state utilize the Corps' Wetlands Delineation Manual (Technical Report Y-87-1, January 1987, Final Report) as the approved method for delineating wetlands, and that the state shall adopt appropriate guidance and regulations to ensure consistency with the Corps' implementation of delineation practices.

### Purpose

Two key aspects of the VADEQ non-tidal wetlands<sup>7</sup> program consist of ensuring that (1) there is no net loss of wetland acreage and function through permitted impacts, and (2) there is a net gain in wetland resources through voluntary programs. In order to accomplish these goals, it is critical to first know the status of wetland resources in Virginia, in terms of location and extent of wetlands in each watershed, and have a general knowledge of the quality of these wetland resources. Secondly, the functions of wetland resources impacted through the Virginia Water Protection (VWP) permitting program must be accurately evaluated to determine those functions to be replaced through compensatory mitigation. It is also important to assess the degree to which the required compensatory mitigation is performing in relation to those impacted functions. Characterizing and evaluating wetlands quality is a third key aspect of this monitoring program. Using this information, the agency can then track changes in wetland acreage and quality within the Commonwealth, target problematic watersheds, and help determine the effectiveness of compensatory mitigation to replace lost wetland acreage and function.

A monitoring and assessment program is defined as the establishment and operation of appropriate devices, methods, systems and procedures necessary to monitor, compile, and analyze data on the condition of wetlands (adapted from the United States Environmental Protection (EPA) Agency's "Elements of a State Water Monitoring and Assessment Program", March 2003). Monitoring is the systematic observation and recording of current and changing conditions, while assessment is the use of that data to evaluate or appraise wetlands to support decision-making and planning processes. Wetlands can be characterized both by their condition and by

---

<sup>7</sup> DEQ administers the wetland monitoring and assessment program in the Office of Wetlands and Stream Protection, and is the legal authority for the protection of non-tidal wetlands. The Code of Virginia designates the authority for tidal wetlands protection to the Virginia Marine Resources Commission (VMRC). See the VMRC Habitat Management WebPages at: <http://www.mrc.virginia.gov/regulations/hm-permits.shtm>.

functions. Wetland condition is the current state as compared to reference standards for physical, chemical, and biological characteristics, while functions represent the processes that characterize wetland ecosystems.

The overarching goal of Virginia's wetland monitoring and assessment strategy was to develop a long-term implementation plan for a wetland monitoring and assessment program that protects the physical, chemical, and biological integrity of the Commonwealth's water resources, including wetlands. In order to accomplish this goal, it is critical to first know the status of wetland resources in Virginia, in terms of location and extent of wetlands in each watershed and have a general knowledge of the quality of these wetland resources. Secondly, the functions of wetland resources impacted through VWP permitting program must be accurately evaluated to determine those functions to be replaced through compensatory mitigation. It is also important to assess the degree to which the required compensatory mitigation is performing in relation to those impacted functions.

Since 2003, the overall wetland monitoring and assessment strategy has been to establish baseline conditions in various broad contexts, such as land use, watershed, and wetland type. This information can then be used to guide management decisions regarding wetland restoration efforts, programmatic compensatory mitigation, and integration with overall WQ Standards. This strategy provides the ultimate framework for an ongoing assessment of the status of the Commonwealth's wetland resources and the success of both wetland regulatory and voluntary programs. The wetlands monitoring strategy will be coordinated with Virginia's comprehensive water quality monitoring program strategy. The monitoring objectives are designed to support regulatory decision-making, allow reporting of wetland conditions, and provide information for policy development.

The wetland monitoring program will also meet the Clean Water Act objectives for water monitoring programs by addressing the quality of the Commonwealth's wetlands and their condition as part of the overall condition assessment of state waters.

### Wetlands Assessment Approach

Virginia has developed an integrated three-tiered approach to wetland sampling and analysis. Comprehensive coverage of all mapped wetlands is achieved with a GIS based analysis of remotely sensed information (Level I analysis). These data are summarized on the basis of small watersheds or hydrologic units. It provides a first order evaluation of the condition and functional capacity of wetlands based on their landscape position.

The second level assessment (Level II) is intended for use in a statistically selected sub-sample of the watershed wetland population and involves a more sophisticated analysis of remotely sensed information and a site visit for verification and additional data collection. The third level assessment (Level III) involves very detailed analysis of wetland performance of specific functions (i.e., habitat and water quality). This involves extensive sampling of a limited number of sites, specifically chosen to allow validation of the conceptual model of wetland function that underlies the Level I and Level II assessments.

A critical part of the overall monitoring and assessment strategy is effective validation and calibration of the underlying models. The Level III assessments are designed to specifically evaluate performance of functions in wetlands under varying degrees of stress, as indicated by the Levels I and II protocols (for additional detail see [wetcat\\_havens-et-al-2018 \(vims.edu\)](#)).

### Wetlands Monitoring Program Development

The DEQ wetlands program, in coordination with the overall DEQ water quality monitoring program, has developed a ten-year plan for wetland monitoring and assessment in Virginia. This work is being accomplished

*Final Guidance for 2024 IR  
Assessment Methodology*

as work products under EPA State Wetland Development Grants CD-983380-01, CD 983815-01, BG 983924-4, and BG-983925-01, BG-98392502, BG-98392503, BG-98392504-0, BG-98392504-4, BG-98392505-0 and BG98392505-9 to the Department of Environmental Quality. The development of this strategy follows the EPA October 2002 draft document “[Elements of a Wetland Monitoring and Assessment Program Checklist](#),” EPA May 2006 “Application of Elements of a State Water Monitoring and Assessment Program for Wetlands” (a supplement to the 2003 EPA document) and includes discussion of the following ‘Ten Essential Elements of a State Water Monitoring and Assessment Program’ ([USEPA, March 2003](#)):


1. Monitoring Program Strategy
2. Monitoring Objectives  
Information derived from monitoring will be used to:
  - Report ambient wetland conditions in Virginia's Clean Water Act (CWA) Section 305(b) reports;
  - Assist in the evaluation of environmental impacts of proposed impacts to wetlands during permit review as part of Virginia's regulatory program;
  - Evaluate the performance of wetland restoration and compensatory wetland mitigation in replacing wetland acreage and function; and
  - Evaluate the cumulative impacts of wetland loss and restoration in watersheds relative to ambient ecological conditions.
3. Monitoring Design
4. Core and Supplemental Water Quality Indicators
5. Quality Assurance
6. Data Management
7. Data Analysis/Assessment  
Examples of different wetland quality data analyses may include:
  - Comparison of wetland quality within a watershed and between watersheds
  - Comparison of wetland quality within a locality and between different localities
  - Comparison of wetland quality within a watershed or locality over time
  - Comparison of wetland quality between wetland types
  - Correlation of wetland type and specific stressor
  - Comparison of wetland quality within and between hydrogeomorphic (HGM) classes
  - Comparison of wetland quality within a specific wetland over time
8. Reporting
9. Programmatic Evaluation
10. General Support and Infrastructure Planning

Virginia’s wetland monitoring and assessment program is being implemented through a cooperative agreement between DEQ and the Center for Coastal Resources Management at the Virginia Institute of Marine Science (CCRM) using funds awarded through EPA’s Wetland Program Development Grants to continue these efforts.

Parameters used in the assessment reflect information from published literature, with consideration of on-going work being conducted through the Mid-Atlantic Wetland Workgroup (MAWWG), regarding each parameter’s validity, usefulness, and utility for field data collection.

*Final Guidance for 2024 IR  
Assessment Methodology*

The protocol for the wetland monitoring and assessment developed in Virginia consists of a multi-tiered sampling design coupled with methods for regulatory updates and field office data delivery (see Figure 2 below). Each assessment level informs the other levels, and is essential in development of the final assessment protocol.



**Figure 2. Multi-tiered sampling design of wetlands**

The elements of Virginia’s wetland monitoring and assessment program are listed in Table 5 below.

<b>Table 5 - Wetland Monitoring and Assessment Program Elements</b>	
Monitoring Strategy	Establish baseline condition of nontidal wetlands by broad category scalable from individual wetland to small watershed to physiographic province to entire State. Guide management decisions regarding restoration, compensation, and regulation of wetlands.
Monitoring Objectives	Support regulatory decision-making. Report wetland condition. Guide policy development. Evaluate cumulative impacts of wetland loss. Evaluate wetland restoration and compensatory mitigation effectiveness.
Survey Design	Three-Tiered: Sample Frame = all NWI wetlands Enhanced GIS analysis (census) – Level I. Probability-based sampling for field assessment of anthropogenic stressors – Level II. Intensive study of biological endpoints (birds, amphibians, water quality) along stressor gradient – Level III+


Assessment Indicators and Methods	<p>Level I: land use adjacent, within 200m, and within 1000m of wetland, wetland size, type, hydroperiod, proximity to other wetlands, road type, road density, and road alignment.</p> <p>Level II: Field assessment of anthropogenic stressors within 30m of wetland assessment point and within 100m of wetland assessment point.</p> <p>Level III: Population and community structure metrics for birds and amphibians. Water quality modification metrics.</p>
Quality Assurance	An EPA-approved Quality Management Plan coupled with the Center Quality Assurance Plan used to prevent random and systematic errors. Techniques include direct electronic field data assimilation to prevent transcription error as well as random return site visits and redundant QA assessment loops.

The strategy continues to develop a complete wetland monitoring and quality assessment in Virginia’s Coastal Plain, Piedmont, and Ridge and Valley physiographic provinces in Virginia. The strategy developed in Phase 1 provides the framework for the ongoing assessment of the status of the Commonwealth’s wetland resources and performance measures for both the wetland regulatory and voluntary programs. Products from this strategy directly support Goal 4 of EPA’s Strategic Plan to provide “...additional focus on assessment of wetland condition” and the National Priority of “wetlands monitoring and assessment.”

One of the advantages of the Virginia protocol for monitoring and assessment of nontidal wetlands is the opportunity to develop a comprehensive assessment of the functional condition of all mapped wetlands whenever there is updated land cover information. This information is particularly useful for evaluating the performance of the regulatory program. It is also useful for indicating cumulative impacts to wetland resources arising from development activities that do not directly impact wetlands. This information can help to raise awareness of consequences and motivate essential change in general land use management and planning that affects lands outside wetland jurisdictional boundaries. Linking decisions in these areas to wetlands policy will be essential to attainment of the no net loss goal.


In a pilot study that used the analysis of wetland condition change, the water quality data was analyzed for Virginia’s coastal plain. By developing catchment areas for the various water quality monitoring stations, the primary objective of this task was to search for relationships between water quality condition recorded at DEQ water quality stations and the condition of wetlands in the contributing drainage.

To test wetland water quality condition scores, Virginia Department of Environmental Quality coastal plain water quality stations (n=99) were used to determine possible trends between wetland water quality condition scores and in-stream water quality metrics (E. coli, fecal coliform, total nitrate nitrogen, DO, pH, and turbidity). Contributing drainage areas were developed for water quality stations using the same protocol for development of individual wetland drainage areas (Figure 3). Water quality station data was compared to contributing drainage wetland water quality condition scores for multiple years (1996, 2001, and 2006).




**Figure 3. Wetland water quality stress condition within the contributing drainage to a Virginia Department of Environmental Quality water quality station.**

While there were no obvious trends between wetland water quality condition score and average DO, pH, and turbidity, there were trends in total nitrate nitrogen, fecal coliform levels, and E. coli levels. As shown in Figure 4, the higher the wetland water quality condition score in the contributing drainage the lower the levels of nitrate, fecal coliforms, and E. coli suggesting a relationship between those water quality parameters and wetland condition.



**Figure 4. Comparison of wetland water quality condition scores (mode)**




**Figure 5. Comparison of in-stream water quality parameters (mean)**

## GIS Wetland Data Viewer

Coordination with VIMS and DEQ staff to design and implement procedures to facilitate the routine application of inventory and monitoring data for regulatory decisions on wetland permits is ongoing and is based on stakeholder input. The data collected has been compiled into a wetland data viewer created by CCRM with substantial input from DEQ. The goal is to automate the processing of database information through GIS necessary to support DEQ's regulatory decision-making, allow reporting of wetland condition, and provide information for policy development.

Stakeholders are continually canvassed for suggestions of additions of data sets and GIS layers that allows Virginia to continue to enhance the GIS-based wetland data viewer for use by regulatory agencies and the general public (see Figure 6). By having a statistically-validated tool that measures wetland quality as a function of habitat and water quality parameters, our permit staff are able to make better permit decisions relative to potential cumulative impacts. Further, we are also able to measure how well we are protecting the function of our more vulnerable wetlands (i.e. isolated wetlands, vernal pools, Atlantic white cedar swamps), by comparing the condition of wetland habitat and water quality parameters, as a function of the assessment scoring over time.

The interactive Wetland Dataviewer, also identified as the Wetland Condition Assessment Tool (WetCAT), allows users to access wetland condition in designated buffers or by HUC unit and allows analysis of linear projects to meet the specific needs of the Virginia Department of Transportation for NEPA review, mitigation requirements, and analyzing linear corridors for cumulative wetland impacts. WetCat received the Governor's Technology Award and the Environmental Council of States Award.



**Figure 6. WetCAT Data Viewer**



The wetland data viewer illustrated above is currently available for general use on DEQ's website at <https://www.deq.virginia.gov/water/wetlands-streams/wetcat> and at [http://cmap2.vims.edu/WetCAT/WetCAT\\_Viewer/WetCAT\\_VA\\_2D.html](http://cmap2.vims.edu/WetCAT/WetCAT_Viewer/WetCAT_VA_2D.html). Development of mechanisms for formatting desk-top delivery of assessment material for permit review are ongoing.

Additional information regarding DEQ's Wetland Monitoring and Assessment Strategy can be found at <https://www.deq.virginia.gov/water/wetlands-streams/monitoring-assessment-strategy>.

### *5.11 Freshwater Probabilistic Assessment*

The freshwater probabilistic monitoring program is designed to allow Virginia DEQ make estimates of water quality with known confidence for 100% of Virginia's freshwater stream miles. Freshwater probabilistic monitoring is not designed to make segment/assessment unit decisions. However, a small number of parameters collected at probabilistic sites can be used to determine use support for that site.

The following parameters will be used for assessment unit impairment decisions:

- 1) Dissolved Oxygen (if 2 out of 2 do not meet standard for stream class)
- 2) pH (if 2 out of 2 do not meet standard for stream class)
- 3) Temperature (if 2 out of 2 exceed standard for stream class)
- 4) Virginia Stream Condition Index – Using the guidance set forth in the freshwater benthic assessment guidance.

### *5.12 Continuous Monitoring Assessment Methodology*

Continuous monitoring, in which multiple observations are collected during a 24-hour period at a relatively high frequency, can provide for a more comprehensive assessment of water quality than what more traditional discrete or "grab sample" monitoring provides because it generates more accurate descriptive statistics and can reveal daily, weekly, monthly, and seasonal variability. High-frequency data collection allows for a more accurate calculation of the frequency and duration of excursions as well. This is especially true for conventional field parameters (i.e., dissolved oxygen concentration and saturation, conductance, pH, temperature, and turbidity). Traditional sampling regimes (semi-monthly, monthly, bimonthly, or quarterly) can only provide a snapshot of conditions, only allowing evaluation of parameter magnitudes and a very rough estimate of excursion frequencies. Another advantage of continuous monitoring is that it monitors environmental conditions at times when field staff rarely sample, such as during nighttime or early morning hours.

Although these are significant benefits of continuous monitoring, the large datasets generated by such monitoring can be a challenge for assessment. It is considered appropriate to apply a 10.5% rule to grab sample datasets, which tend to be relatively small, but applying that rule to a continuous monitoring dataset, which can contain as many as tens of thousands of observations, could result in a water being assessed as attaining the standard for a parameter that it may be actually impaired for. Thus, using continuous monitoring data for listing and delisting waters requires caution and thoughtfulness. The following rules were crafted with this in mind:

#### Rule 1

A continuous monitoring dataset that is eligible for assessment must cover at least thirty 24-hour periods (with the exception of data being assessed for maximum temperature exceedances, which must cover at least fifteen 24-hour periods). This allows for an informative characterization of a water during the

critical period (May to September) when exceedances of conventional field parameters are most expected.

#### Rule 2

The continuous monitoring dataset will have undergone rigorous and standardized QA/QC screening before analysis. Every 24-hour period with at least 75% of its observations deemed as valid should be assessed and counted as a single sample. Grab samples must be collected during the run that a continuous monitor is deployed.

#### Rule 3

Daily averages are the mean of all valid observations (including grab samples from the same station) collected during a 24-hour period. An excursion of the DO daily average is defined as a mean calculated from all valid data collected during a 24-hour period (midnight-to-midnight) that is below the appropriate daily average criterion for a given water.

#### Rule 4

A 24-hour period exceeds minimum and maximum instantaneous criteria when > 10.5% of its observations exceed the criteria. Any two such days, even if consecutive, would count as two separate exceedances. Water temperature should be evaluated for exceeding increases as described in Section 9VAC25-260-60 of the Water Quality Standards. Exceedances recorded during the continuous monitoring run should be combined with grab samples within the assessment data window. A 10.5% rule should then be applied to the combined data set.

#### Rule 5

For water temperature standards specifying a maximum hourly change (9VAC25-260-70), a 10.5% rule should be applied to the total number of monitored hours where data meet QA/QC (including hours of the first and last days of deployment.) The frequency of hourly change exceedances should be determined by dividing the number of hourly change exceedances into the total number of clock hours monitored. An additional continuous monitoring dataset, collected during a subsequent year during the same month(s) as the listing dataset, must be used for delisting.

#### Rule 6

If a continuous monitoring dataset is used to list a water on the 303(d) Impaired Waters List, then an additional continuous monitoring dataset, collected during a subsequent year, during the same month(s) as the listing dataset, must be used to delist it. A water that was previously listed using grab samples may be delisted using continuous monitoring data collected for at least 30 days, during a subsequent year and during the same month(s) when exceedances were previously found.

#### Rule 7

Continuous dissolved oxygen data collected in Chesapeake Bay waters (mainstem and tributaries) should only be assessed against the Open Water 7-day mean and instantaneous minimum criteria and Migratory Fish Spawning 7-day mean and instantaneous minimum criteria. A continuous DO dataset

that indicates an applicable 7-day mean criterion was exceeded in more than 10% of the observed 7-day periods in a three-year assessment period should be considered evidence of impairment. A 7-day DO mean should be calculated from observations taken over at least five days, and each day must have been monitored for at least 75% of the diurnal cycle. Per Rule 2-A of the 2017 EPA addendum (EPA 903-R-17-002, CBP/TRS 320-17), a continuous DO dataset that indicates there were more than two consecutive “violating days” of the applicable instantaneous minimum criterion in a three-year assessment period should be considered evidence of impairment. A “violating day” is defined as one where the DO concentration was below the applicable instantaneous minimum criterion for at least 10% of the time over that day. To support evidence of criteria attainment, the continuous monitoring dataset should span at least one summer season (June-September).

#### SCENARIO # 1:

A monitor was deployed July 31 at noon and run continuously through September 1 (noon) at a station. Five grab samples were collected at that station during the same year as the monitor’s deployment (during February, April, June, July, and November); no other data exist in the assessment window for this station. No excursion of the minimum DO criterion is detected in the grab samples, while four 24-hour periods in the continuous monitoring dataset have >10.5% of their total observations not meeting the minimum DO criterion.

##### *Assessment*

- The sample size is 36 (31 continuous monitor “samples” + 5 grab samples). The first and last 24-hour periods observed by the monitor should not be used for assessment, since at least 75% of the diurnal cycle was not recorded by the monitor on these two days.
- The exceedance rate is 11.1% and is therefore excessive. Accordingly, the water fails to meet the water quality standard for DO and should be placed on the 303(d) Impaired Waters list for this parameter.
- To delist this water, a continuous monitor must be set up for the same length of time as the original run, during the same month (August). Grab samples should be collected during other months of the year to maintain “temporal representativeness”.

#### SCENARIO #2:

A monitor was deployed April 1 at noon and run continuously through August 31 (noon). Three grab samples were collected at that station during the same year as the monitor’s deployment (February, October, December), and ten were collected two years previously. None of the newer grab samples exceed any standard, but twelve 24-hour periods, the majority clustered in the summer months, have >10.5% of their observations not meeting the minimum DO criterion. The older dataset contained two excursions of the DO minimum criterion, and these excursions were also found during the summer. The water had therefore been placed on the 303(d) Impaired Waters list during the previous cycle.

##### *Assessment*

- The sample size is 164 (151 continuous monitoring samples + 3 new grab samples + 10 older grab samples).
- The exceedance rate is not technically excessive (8.5%), as defined by the 10.5% rule. However, there is evidence that the water experiences hypoxia during the summer. Before considering to delist the water, the assessor should address the following questions:

- a) Do the excursions observed in the continuous monitoring dataset correspond temporally to those found in the older dataset used to list the water?
- b) What is the average duration of the excursions? It would not be wise to delist a water characterized by long durations of excursions—particularly for excursions of the DO minimum.
- c) What is the temporal frequency of the excursions? Are the exceeding 24-hour periods mostly consecutive, or are they spaced relatively far apart (potentially allowing for aquatic life recovery if the excursions are not too severe)?
- d) Were hydrological and/or weather conditions similar between the current dataset and the older dataset?
- e) Were there specific documented practices put into place that have improved water quality over the two-year period? (refer to Appendix D for more details)
- f) Are excursions observed in the grab samples collected during the continuous monitoring run?

Note that this is not an exhaustive list of considerations. To resolve situations such as the one described above, the assessor may need to rely on best professional judgment rather than following a strict interpretation of the 10.5% rule.

## **PART VI. PROCEDURES FOR CITIZEN AND NON-AGENCY DATA**

For the purposes of this guidance document, a citizen water quality monitoring program, or “citizen monitoring,” is defined as water quality monitoring which uses volunteers to collect the data. Some of these programs are run by local governments, Soil and Water Conservation Districts, citizen organizations, community organizations or colleges. Generally, K-12 school monitoring is conducted for educational purposes and does not fall under citizen monitoring unless working in cooperation with existing citizen monitoring efforts. Citizen monitoring is not defined as monitoring conducted by all entities external to DEQ, such as colleges and local governments, unless volunteers are used in their efforts.

DEQ does routinely receive water quality data from non-citizen volunteer sources such as local governments, universities, and other non-state or federal sources. The review and assessment of non-agency data is done using the same QA/QC review as with citizen monitoring data.

In 1997, Water Quality Monitoring, Information and Restoration Act (WQMIRA) was passed by the Virginia General Assembly. This bill charged DEQ with monitoring and assessing all the waters within the Commonwealth. During this same General Assembly session, the position of citizen monitoring coordinator was added into the operating budget of DEQ. The primary duties of the citizen monitoring coordinator were providing guidance and support to citizen water quality monitoring groups in the development of monitoring programs and quality assurance project plans. In addition, the citizen monitoring coordinator facilitated communication among citizen groups and other state agencies, sponsoring citizen monitoring seminars, promoting the use of citizen water quality data in a manner consistent with the data use goals of the organization and encouraging additional citizen monitoring efforts. In 2002, the Virginia General Assembly passed legislation that established the Virginia Citizen Water Quality Monitoring Program in the Code of Virginia (§62.1-44.19:11).

In 2004, the citizen monitoring coordinator position evolved into the role of water quality data liaison. This was done to centralize the task of requesting any and all available data collected outside of DEQ for inclusion into water quality assessment reports and follow up monitoring by DEQ. The duties and responsibilities of the

*Final Guidance for 2024 IR  
Assessment Methodology*

former position regarding citizen monitoring data submissions and working with the citizen monitoring community have been maintained and expanded to include all other non-DEQ potential sources of water quality data.

#### **Assessment Process:**

The process of assessing water quality data submitted to DEQ involves staff from both the central office and the regional offices. In order to include any citizen or non-agency monitoring data in the biennial 305(b)/303(d) Integrated Water Quality Assessment Report (Integrated Report), it must be received and evaluated by central office staff before the published data solicitation deadline. By adhering to the tasks outlined below, the agency can ensure that all qualifying monitoring data is properly assessed.

#### **Submitting Data for Evaluation:**

1. All water quality data provided to DEQ from citizen and non-agency organizations should be uploaded through the Virginia Data Explorer before the published data solicitation deadline. The DEQ water quality data liaison and the appropriate QA/QC staff in the Water Monitoring and Assessment (WMA) Program will review all standard operating procedures (SOPs), QA/QC plans or Quality Assurance Project Plans (QAPPs) for each citizen/non-agency monitoring group submitting water quality data.

For citizen/non-agency chemical and bacteria monitoring programs, the liaison will work with the WMA quality assurance (QA) coordinator. The liaison, QA coordinator, and the Biological Monitoring Program Coordinator will review all supporting documentation for benthic macroinvertebrate citizen/non-agency monitoring programs. Based upon the review of all procedures, the appropriate use of the data will be determined based on a three-tiered system.

2. The designation of DEQ tiered uses of data will be determined based upon the review of all procedures in conjunction with the organization submitting the water quality data. Any changes in QA/QC and/or SOP methods and/or any additions or deletions of current monitoring sites should be brought to the attention of the data liaison and the appropriate QA/QC staff in the Water Monitoring and Assessment (WMA) Program.

Since 2007, DEQ has provided a [data use authorization form](#) to monitoring groups. Because not all non-DEQ organizations may wish to have their data used for water quality assessment reports, this form allows DEQ to meet their wishes. This authorization form cannot be used to upgrade the use of lower tiered data for a higher tiered purpose. Such an example would be a data submitter requesting DEQ to assess their data for Level III (use for 303(d) listing/delisting of impaired waters) based on Level II or Level I quality data.

#### **Central Office Assessment Tasks:**

1. The QA coordinator, with the help of the liaison, will provide a copy of all Level II and III citizen and non-agency monitoring data received during a given assessment cycle to the regions. The format of the data provided to the regions will be as follows:
  - a. Data will be in electronic spreadsheet format compatible with programs used by the regional assessors.
  - b. Level II and III data will be combined with columns denoting the applicable QA status and assessment use for that data point.

- c. All data not meeting QA/QC requirements or otherwise not relevant for assessment will be omitted by the QA Coordinator. However, an unedited master copy of all data submitted will be maintained. Examples of excluded data could include benthic macroinvertebrate samples collected outside the spring and fall season or lake monitoring performed outside of the April-October timeframe.
  - d. At a minimum, all citizen and non-agency monitoring sites submitted to the regions for assessment will contain the following metadata:
    - i. Name of waterbody monitored
    - ii. Latitude and Longitude information
    - iii. Physical description of the site (i.e. At Route 646 bridge crossing)
    - iv. Previous IR cycle assigned DEQ Station IDs
    - v. Assessment Region Code
2. The liaison and QA coordinator will review data collected without SOPs and QAPPs plans. This data will be acknowledged in the applicable river basin evaluation as appropriate.
  3. In coordination with the liaison, the appropriate WQM QA/QC staff and the assessment coordinator, each regional office should provide any appropriate final editing of the citizen and non-agency monitoring assessment levels. The data quality level assigned by the QA coordinator or appropriate QA/QC staff will reflect the QA measures included for the dataset (eg. QAPP, SOP, calibration and QA sampling conducted), but may not reflect the physical location of the site. Regional assessment staff should review the site locations to ensure that the assigned data level and the resulting assessment decisions are appropriate, or if the data should be downgraded or excluded. Final decisions to exclude or downgrade data due to sampling location should be reviewed and approved by the QA coordinator. A narrative explanation of the reasoning for site exclusion, along with any supporting data should be retained by the QA coordinator along with the original dataset and provided to the citizen/non-agency monitoring group for review and correction.
  4. After the release of the final biennial Integrated Report, regional monitoring staff should review the station list results with assessment staff to consider including sites for follow-up monitoring as appropriate to their regional monitoring plan.
  5. With the help of the liaison, the QA coordinator will provide all data approved by DEQ for use in the Integrated Report as tables to be saved on the internal Water Quality Assessment SharePoint site. These data will also be available by request.

**Regional Office Assessment Tasks:**

1. All approved conventional parameter data should be summarized by major watershed and characterized according to the procedures and considerations in Part V of this manual.
2. For benthic macroinvertebrate monitoring programs used by citizen and non-agency monitoring organizations, data will be assessed based on the criteria outlined in *Guidance Memo No. 06-2010, Guidelines for DEQ Review and Approval of Biological Monitoring QAPPs*.
  - a. For organizations that complete the requirements outlined in the guidance memo for Level III, DEQ staff will assess the data for the purposes of 305(b) water quality assessment and 303(d)

listing and delisting of impaired waters. If a validation study showed inconclusive correlation with DEQ benthic protocols, the corresponding scores showing inconclusive correlation will not be assessed as Level III. The data from an inconclusive validation study may be used to characterize waters as having insufficient information with or without the need for follow-up monitoring (Cat 3C or 3D).

- b. For all other methods not validated by DEQ or using DEQ protocols, biological monitoring sites characterized by citizen and non-agency organizations as “excellent,” “good” or “acceptable” should be designated as “Area of low probability for adverse conditions” (Category 3D). Biological sites periodically characterized as “fair,” “poor,” “unacceptable” or “moderate” should be designated as “Area of medium probability for adverse conditions” and listed as insufficient data and prioritized for follow-up monitoring (Category 3C). Likewise, biological sites that are consistently “poor” or “unacceptable” should be characterized as “Area of high probability for adverse conditions” and listed as insufficient data with DEQ follow up monitoring to be prioritized (Category 3C).
  - c. Citizen or Non-DEQ benthic data collected outside of the established sampling seasons will be excluded from assessment.
3. Segment lengths represented by a monitoring site should be determined using the mileage delineation guidance found in Section 5.1. Each monitoring site used in the assessment should have a unique station ID using a system similar to the DEQ station ID system. The regional office staff assigns this station ID to each citizen/non-agency monitoring site and relays the newly assigned DEQ station ID to the QA Coordinator.
4. Level III data collected at sites that complement and are comparable (i.e. chemical to chemical comparisons and biological to biological comparisons) to DEQ monitoring sites, should be included in assessment determinations. However, the final assessment of that segment will be made using the DEQ monitoring data and the data collected by the monitoring organization would be used as supplemental data. If the assessment results differ between these stations, the data will be reviewed on a case-by-case basis.
5. Level III data collected at sites that do not complement or compare (i.e. benthic to chemical comparisons) to DEQ monitored sites, should be included in assessment determinations. For example, Level III citizen benthic macroinvertebrate data shows impairment of the aquatic life use while a nearby DEQ chemical monitoring station does not directly show impairment of the aquatic life use. The assessment unit would be impaired for benthic macroinvertebrates.
6. Level II ambient and bacteria data collected at sites will undergo the similar evaluation process as used for Level III and DEQ results. Since Level II data may have some variation in quality assurance, corresponding waterbodies that indicate poor water quality will be listed as insufficient data and prioritized for follow-up monitoring (Category 3C). Waterbodies that have Level II data indicating good water quality will be listed as insufficient data with low probability for adverse conditions (Category 3D).
7. If during the regional review, a discrepancy between data from DEQ monitoring stations and data from nearby citizen/non-agency monitoring stations is believed to be suspect, the QA coordinator should be notified and effort made to rectify the discrepancy.

The QA coordinator and liaison will evaluate the potential causes for the data disparity and/or review the QAPP and the monitoring techniques of the data submitting group. After this evaluation is complete and a problem is confirmed, appropriate corrective actions will be recommended to the monitoring group for inclusion in the citizen/non-agency monitoring organization's QAPP and/or SOPs.

Until the discrepancies with the data and/or methods are fully evaluated by DEQ, the data (either for the parameter(s) of concern or for all observations) should not be used in assessment determinations. If the citizen or non-agency monitoring group does not initiate corrective action, the QAPP for that parameter and/or for the group may no longer be considered valid by DEQ, and the data will not be considered for statewide water quality assessments.

8. Regional assessment staff should use best professional judgement to determine which sampling sites are appropriate for use in assessment determinations until changes to citizen and non-agency monitoring plans can be reviewed to exclude inappropriate sampling sites. Final decisions to exclude or downgrade data due to sampling location should be reviewed and approved by the QA coordinator. A narrative explanation of the reasoning for site exclusion, along with any supporting data should be retained by the QA coordinator along with the original dataset and provided to the citizen/non-agency monitoring group for review and correction.
  - a. If during regional review, location information of Level III citizen/non-agency monitoring stations cannot be confirmed, the data will be downgraded to Level II until the location of the sampling can be confirmed. DEQ regional assessment staff should work with the QA Coordinator to document these issues as they are discovered.

### **Other State and Federal Water Quality Data**

After review and approval of monitoring and QA/QC protocols, DEQ will consider data generated by other State and Federal monitoring programs for use in the Integrated Report. DEQ accepts water quality data from several state and federal agencies including, but may not be limited to the Virginia Department of Health, Tennessee Valley Authority, National Park Service, United States Forest Service, and the United States Geological Survey.

**Virginia Department of Health (VDH)** - DEQ receives and lists areas closed by VDH for swimming and shellfish harvesting due to high bacteria levels. All Enterococcus bacteria results provided by VDH are also used along with any DEQ water quality data in assessing water quality. Any other water quality data collected by VDH and shared with DEQ will be used at the latter agency's discretion.

**Tennessee Valley Authority (TVA)** - The TVA routinely monitors for *E. coli* bacteria along TVA reservoirs in Virginia. These data are considered acceptable for assessing water quality in Virginia.

**National Park Service** - The National Park Service has several long-term monitoring programs in place at many of the national parks in Virginia. Many of the parks monitor for chemical and benthic macroinvertebrate parameters using varying methodologies or procedures. Because of this, the liaison and QA coordinator provide guidance to the regional office assessment staff in assessing data received from the parks.

**United States Forest Service (USFS)** - The USFS program collected macroinvertebrate data from numerous monitoring stations within the George Washington and Jefferson National Forests. Sampling for

*Final Guidance for 2024 IR  
Citizen and Non-Agency Data*



macroinvertebrates is conducted utilizing the same collection methodology (Plafkin et al 1989) that DEQ biologists use in the ambient biomonitoring program. Therefore, the raw data collected by the USFS should be highly comparable with DEQ data. The USFS has used the Macroinvertebrate Aggregated Index for Streams (MAIS) to assess this raw data and make an initial water quality interpretation.

The DEQ regional biologists and planners may use the data, provided to DEQ by the USFS, in the Integrated Report if they find it acceptable for assessment purposes. If the regional biologists or planners have information that conflicts with the initial USFS assessment, or for any other reason questions the USFS stream assessment, they may elect to disregard the USFS assessment results until further verification can be obtained. If the initial assessment is not used, documentation relating to this decision will need to be provided. The regional biologists may elect to reevaluate the raw data using the Virginia Stream Condition Index (VSCI) metrics to confirm consistent assessment methodology and conclusions. If differences become apparent, the regional biologists may decide not to use the assessment data in the Integrated Report until an on-site stream visit can be performed and conditions verified. Final assessment results of the USFS data should be consistent with the ambient biological assessment criteria described in Section 5.4 of this guidance. Any non-approved data will not be used directly in the assessment.

**United States Geological Survey (USGS)** - The USGS monitors several water quality stations throughout Virginia. Data collected by the USGS is considered Level III by DEQ and is used in assessing water quality including 303(d) impairment listings and delistings. Water quality parameters for which there are no established numerical criteria in Virginia’s water quality standards or developed assessment thresholds for this guidance manual are not used for the purposes of 303(d) impairment listing but can be used to assess waters for follow-up monitoring (Category 3C/3D).

## **PART VII. 303(d) LISTING/DELISTING and TMDL PRIORITY RANKING**

### *Alternative Control Waters (Category 4B/5R)*

Previous iterations of this manual included a process for listing and delisting facilities with water quality-based effluent limits in VPDES permits (Category 4B/5E). The listing methodology did not include an evaluation of water quality monitoring data reflecting ambient, instream conditions. The facility listings were based on water quality-based effluent limits in VPDES permits with compliance schedules or an enforcement action associated with such limits. Discharge monitoring report (DMR) data and/or enforcement correspondence was required to justify removal or delisting facilities from Virginia’s 303(d) list.

DEQ Water Permitting and Compliance programs regulate water resources and water pollution from point sources in Virginia. Information on water permit facilities in Virginia, including compliance and enforcement data is currently available online through the [EPA Enforcement and Compliance History Online \(ECHO\) tool](#).

### **Rule 1**

#### Category 4B – Alternative Control

EPA’s 2006 IR Guidance acknowledged that the most effective method for achieving water quality standards for some water quality impaired segments may be through controls developed and implemented prior to the TMDL development and/or implementation (referred to as a “4B alternative”). DEQ requests EPA to evaluate, on a case-by-case basis, the Commonwealth’s decisions to exclude or delist certain segment/pollutant

*Final Guidance for 2024 IR  
Citizen and Non-Agency Data*

combinations from Category 5 based on the 4B alternative. A 4B rationale will be provided to EPA in the submission of the 2024 IR which supports the Commonwealth’s conclusion that there are “other pollution control requirements” sufficiently stringent to achieve applicable water quality standards within a reasonable period of time.

**Required elements of the 4B rationale:**

Specifically, this rationale should include:

- (1) a statement of the problem causing the impairment,
- (2) a description of the proposed implementation strategy and supporting pollution controls necessary to achieve water quality standards, including the identification of point and non-point source loadings that when implemented assure the attainment of all applicable water quality standards,
- (3) an estimate or projection of the time when water quality standards will be met,
- (4) a reasonable schedule for implementing the necessary pollution controls,
- (5) a description of, and schedule for, monitoring milestones for tracking and reporting progress to EPA on the implementation of the pollution controls, and
- (6) a commitment to revise, as necessary, the implementation strategy and corresponding pollution controls if progress towards meeting water quality standards is not being shown.

**Rule 2**

Category 5R – Advance Restoration Plan

EPA’s 2016 IR Guidance acknowledges that restoration plans that are done ahead of TMDLs may be the best option to reach water quality standards faster. However, when the advance restoration plan lacks enforceable “other pollution control requirements,” the water cannot be assessed as 4B, and must remain in category 5. In EPA’s 2016 IR Guidance the national subcategory of 5-alternative (now advance) is discussed and introduced. In Virginia this is the state subcategory 5R (detailed description in Appendix D-2). When DEQ develops an advance restoration plan to a TMDL, DEQ requests EPA to review the plan. While EPA cannot approve the plan, they can review it and accept it as a 5R advance restoration plan. The six main elements of an acceptable 5R advance restoration plan are outlined in Appendix D-2. Once EPA has accepted a 5R advance restoration plan, the impaired waters that are addressed by this plan are to be assessed as state subcategory 5R.

*Delisting Rules*

**Rule 1**

Waters listed as impaired and needing a TMDL in the Integrated Report will remain on the list and tracked in subsequent Integrated Reports until:

- An EPA approved TMDL is developed for all pollutants causing impairment

**OR**

A subsequent assessment of new monitoring data (or in special cases, modeling data) results show that the water is no longer impaired and EPA approves the delisting of the water.

**Rule 2**

Documentation required by EPA for delisting previously listed impaired waters that are now restored:

**Scenario # 1:** when new data demonstrates a previously impaired waterbody is currently attaining WQS, DEQ should submit the following information to justify the delisting of this segment from the 303(d) list.

- Cause Group Code, Assessment Unit ID, Stream Name and Listed Parameter, Associated Use(s) and Initial List Cycle
- ATTAINS Rationale for the decision to remove the previously impaired segment from the 303(d) list
- Any differences between the sampling techniques should be explained

**Scenario # 2:** when new water quality modeling determines the stream is now attaining WQS, DEQ should submit the following documents to justify the removal of this segment from the 303(d) list.

- Cause Group Code, Assessment Unit ID, Stream Name and Listed Parameter, Associated Use(s) and Initial List Cycle
- ATTAINS Rationale for the decision to remove the previously impaired segment from the 303(d) list
- Submission of any new data that were used in the modeling
- A copy of the EPA approved model that was used. A summary of the differences between the new and the old models. The reasons why the stream attains WQS under the new model opposed to the former model (data, modeling assumptions, modeling applications, etc.)

**Scenario # 3:** when new management practices from point and/or nonpoint sources lead to the attainment of WQS, DEQ should submit the following documents to justify the removal of this segment from the 303(d) list.

- Cause Group Code, Assessment Unit ID, Stream Name and Listed Parameter, Associated Use(s) and Initial List Cycle
- ATTAINS Rationale for the decision to remove the previously impaired segment from the 303(d) list
- Submission of the most recent 2 years of water quality data that indicate the water is a candidate for delisting and
- A description of the new management practices and other changes that have occurred in the watershed to explain the change in water quality.

The TMDL staff should apply the Proactive Approach, as appropriate, any time a TMDL is scheduled for development. Appendix D contains additional procedural information on this approach.

**Scenario # 4:** when errors are detected in the rationale for the initial listing of the segment or WQ Standards have been modified and the segment is attaining WQ Standards, DEQ should submit the following documents to justify the removal of this segment from the 303(d) list.

- Cause Group Code, Assessment Unit ID, Stream Name and Listed Parameter, Associated Use(s) and Initial List Cycle
- ATTAINS Rationale for the decision to remove the previously impaired segment from the 303(d) list
- Documentation of the errors in the initial listing
- A copy of the data and/or modeling that demonstrates the segment attains WQS

The following statement should be noted in delist documentation where the WQ Standards have been amended and the water is now classified as a Class VII swamp water:

“The \_\_\_\_\_ watershed was reclassified as Class VII swamp water in Virginia’s EPA-approved water quality standards regulations during the 2024 cycle. Per Virginia’s Water Quality Standards (9VAC25-260-50), numeric dissolved oxygen standards only apply to Class VII waters when there is sufficient evidence the narrative criterion is not protective of aquatic life uses. To date, this Class VII water has not exhibited a need for a site-specific DO criterion, so the dissolved oxygen impairment has been removed.

In certain cases EPA may request additional documentation to justify the removal of the segment from the 303(d) list.

### **Rule 3**

A new impairment is “nested” when it is determined that the impairment has the same source/cause as a previously listed impairment within an existing TMDL. In such a case, it is assumed the new impairment is adequately addressed by the pre-existing TMDL and should thus be classified as Category 4A. Assessors should coordinate with TMDL staff to review nesting guidance for specific qualifications for nesting, procedural requirements, and appropriate documentation.

Bacteria impairments within the existing TMDL watershed or within the “tidal range” of the existing TMDL boundary can be immediately nested when land uses in the existing TMDL and newly impaired segment are comparable and all existing sources are accounted for in the TMDL. A narrative nesting memo is not necessary for these impairments.

Nesting non-bacterial impairments may be appropriate if the existing TMDL(s) addresses all appropriate stressor(s) for benthic impairments or all source(s) for other non-bacterial impairments. It is not appropriate if new applicable stressor(s) or source(s) exist.

A rationale memo describing the TMDL, the watershed, and the relevant assessments unit(s) as well as justification for the nesting should be submitted to EPA before nesting impairment(s) under the following conditions:

- Non-bacteria impairment (e.g., nesting a pH impairment under a TMDL originally addressing DO and nutrients)
- Bacteria impairment outside the boundary of a TMDL watershed or not within the “tidal range” of the existing TMDL boundary.

### **Rule 4**

Section 303(d) requires States to “establish a priority ranking” for the waters it identifies on the impaired waters list, taking into account the severity of the pollution and the uses to be made of such waters, and to establish TMDLs “in accordance with the priority ranking.” Federal regulations provide that “schedules for submissions of TMDLs shall be determined by the Regional Administrator and the State” (40 CFR 130.7(d)(1)). Other reasonable factors such as the State’s use of a rotating basin approach or commitments specified in court orders or consent decrees may also be considered when States develop priorities and schedules.

In scheduling TMDLs for development, every effort should be made to address all related impairments in a watershed at the same time. If endangered species are affected by an impairment listing, TMDL development

*Final Guidance for 2024 IR*

*303d Listing/Delisting and TMDL Priority Ranking*

should be scheduled as expeditiously as possible. If a public water supply is affected by an impairment listing, TMDL development should be scheduled as expeditiously as possible. In the absence of impacts to public water supplies or endangered species, a watershed approach should be used for TMDL development scheduling. Other factors that may impact TMDL scheduling include public interest and support, locally available funding to implement controls, or the coordination of TMDL development efforts with an adjoining state.

Starting in December 2013 as part of EPA’s 303(d) Program Vision, EPA tasked states with prioritizing impaired waters for TMDL or TMDL alternative development over the 2016-2022 six year window. A statewide prioritization strategy included evaluation of waters impaired by 1) low dissolved oxygen concentrations, 2) high or low pH levels, 3) high temperatures in cold water streams, and 4) impairments to the benthic macroinvertebrate communities. Since these impairments result in impacts to the aquatic ecosystem, DEQ sought to prioritize waters impaired for the aquatic life use where additional benefits to aquatic ecosystems could be gained. To maximize those additional benefits, DEQ conducted an analysis of small watersheds (12-digit hydrological unit) that also contained one or more of the following criteria:

1. The presence of a state and/or federally listed threatened or endangered species (spatial data provided by DWR)
2. The presence of DWR Stream Conservation Unit, which is based on:
  - The presence of a rare, threatened, or endangered species; or
  - The presence of an aquatic ecological community of high integrity

The second cycle of EPA’s 303(d) Program Vision will encompass the 2024-2032 window and again calls for the prioritization of impaired waters for TMDL or TMDL alternative development. DEQ is working towards establishing priorities for the second cycle and began by seeking initial public input in September 2020 to inform the process of identifying types of water quality impairments and other strategic measures for developing a list of priority waters. DEQ will solicit public comment for the 2024-2032 priorities prior to finalizing Vision priorities.

Because there is a two-year gap between the end of the first cycle and the beginning of the second cycle, DEQ provided EPA with a priority list of impaired waters for the 2023-2024 window (a.k.a. “bridge” priorities). The impaired waters on this list included TMDL development work that is ongoing from the first Vision cycle (2016-2022) and included some additional impairments selected using a strategy similar to that used for the selection of priorities for the first cycle as well as a strategy to increase efficiency of plan development. These bridge priorities were included in the 2022 Integrated Report, Appendix 1A, by denoting “H” under the “TMDL Development Priority” column indicating a high priority for TMDL or TMDL alternative development in the 2023-2024 window. DEQ solicited public comments on the Integrated Report in June 2022, which has since been approved by EPA.

After the TMDL and TMDL alternative priorities have been selected, the order in which TMDLs and TMDL alternatives are established might be subject to some modifications to accommodate logistical efficiencies or data availability. The process is a dynamic process, and any priority ranking may be changed if substantial factors change or become apparent during the scheduling process.

## **APPENDIX A**

### **Clean Water Act Sections**

#### **SEC. 305. WATER QUALITY INVENTORY**

- (b) (1) Each State shall prepare and submit to the Administrator by April 1, 1975, and shall bring up to date by April 1, 1976, and biennially thereafter, a report that shall include—
- (A) a description of the water quality of all navigable waters in such State during the preceding year, with appropriate supplemental descriptions as shall be required to take into account seasonal, tidal, and other variations, correlated with the quality of water required by the objective of this Act (as identified by the Administrator pursuant to criteria published under section 304(a) of this Act) and the water quality described in subparagraph (B) of this paragraph;
  - (B) an analysis of the extent to which all navigable waters of such State provide for the protection and propagation of a balanced population of shellfish, fish, and wildlife, and allow recreational activities in and on the water;
  - (C) an analysis of the extent to which the elimination of the discharge of pollutants and a level of water quality which provides for the protection and propagation of a balanced population of shellfish, fish, and wildlife and allows recreational activities in and on the water, have been or will be achieved by the requirements of this Act, together with recommendations as to additional action necessary to achieve such objectives and for what water such additional action is necessary;
  - (D) an estimate of (1) the environmental impact, (ii) the economic and social costs necessary to achieve the objective of this Act in such State, (iii) the economic and social benefits of such achievement, and (iv) an estimate of the date of such achievement; and
  - (E) a description of the nature and extent of nonpoint sources of pollutants, and recommendations as to the programs which must be undertaken to control each category of such sources, including an estimate of the costs of implementing such programs. (2) The Administrator shall transmit such State reports, together with an analysis thereof, to Congress on or before October 1, 1975, and October 1, 1976, and biennially thereafter.

#### **GRANTS FOR SEC. 106. POLLUTION CONTROL PROGRAM**

- (e) Beginning in fiscal year 1974 the Administrator shall not make any grant under this section to any State which has not provided or is not carrying out as a part of its program—
  - (1) the establishment and operation of appropriate devices, methods, systems, and procedures necessary to monitor, and to compile and analyze data on (including classification according to eutrophic condition), the quality of navigable waters and to the extent practicable, ground waters including biological monitoring; and provision for annually updating such data and including it in the report required under section 305 of this Act;

## SEC. 204 LIMITATION AND CONDITIONS

(a) Before approving grants for any projection for any treatment works under section 201(g)(1) the Administrator shall determine—

“that (A) the State in which the project is to be located (1) is implementing any required plan under section 303(e) of this Act and the proposed treatment works are in conformity with such plan, or (ii) is developing such a plan and the proposed treatment works will be in conformity with such plan, and (b) such State is in compliance with section 305(b) of this Act;”

## SEC. 314. CLEAN LAKES

(a) Each State shall prepare or establish, and submit to the Administrator for his approval—

“(A) an identification and classification according to eutrophic condition of all publicly owned lakes in such State;

“(B) a description of procedures, processes, and methods (including land use requirements), to control sources of pollution of such lakes;

“(C) a description of methods and procedures, in conjunction with appropriate Federal agencies, to restore the quality of such lakes;

“(D) methods and procedures to mitigate the harmful effects of high acidity, including innovative methods of neutralizing and restoring buffering capacity of lakes and methods of removing from lakes toxic metals and other toxic substances mobilized by high acidity;

“(E) a list and description of those publicly owned lakes in such State for which uses are known to be impaired, including those lakes which are known not to meet applicable WQ Standards or which require implementation of control programs to maintain compliance with applicable standards and those lakes in which water quality has deteriorated as a result of high acidity that may reasonably be due to acid deposition; and

“(F) an assessment of the status and trends of water quality in lakes in such State, including but not limited to, the nature and extent of pollution loading from point and nonpoint sources and the extent to which the uses of lakes is impaired as a result of such pollution, particularly with respect to toxic pollution.

“(2) SUBMISSION AS PART OF 305(b) (1) REPORT. – The information required under paragraph (1) shall be included in the report required under section 305(b) (1) of this Act, beginning with the report required under such section by April 1, 1988”.

## **APPENDIX B**

Virginia Department of Environmental Quality  
Biological Monitoring Program  
305(b) Assessment Fact Sheet

Regional Office:

Regional Biologist's Signature: \_\_\_\_\_

Review Date:

River Basin:

Stream Name and Site Location:

Station ID #:

Reference Station ID #:

Assessment Method:

VSCI

Coastal Plain (MACS)

### **Biological Assessments for the Last Six Years**

<b>Year</b>	<b>Spring score</b>	<b>Spring assessment</b>	<b>Fall score</b>	<b>Fall assessment</b>
2015				
2016				
2017				
2018				
2019				
2020	0.0		0.0	
Seasonal avg 6-yrs	0.0		0.0	
Seasonal avg last 2-yrs	0.0		0.0	
Final 6-yr average	0.0		0.0	
Final 2-yr average	0.0		0.0	

Note, because of the long, six-year time frame covered by this review and for a variety of reasons, some sites may not have been sampled during every year or season and/or an assessment ranking or score may not be available for every "cell" in the above table. The above table is intended to be a convenient method to summarize and review all the data available for the reporting period. The final assessment ranking for each site should be based on a review of all the available rankings shown in the above table and any pertinent supplemental data described below. For the purpose of Integrated Report preparation, if more recent bioassessment rankings differ significantly from earlier rankings, primary consideration should be given to the more recent assessed data. This is described in more detail of Section 5.4 of the Integrated Report Guidance Manual.

### **Supplemental Information (if applicable):**

Are any seasonal differences noted?

Summary of any comments associated with assessments.

*Final Guidance for 2024 IR*

*Appendix B*



Have any factors been observed in watershed that may be affecting the benthic community? Have there been any recent changes in activity in the watershed that may have affected the more recent bioassessments. Are these changes likely to affect the benthic community for a short or long-term basis?

**Final Assessment Rating:**

## **APPENDIX C**

### **Classification of Virginia's Shellfish Growing Areas**

Robert E. Croonenberghs, PhD

The Division of Shellfish Sanitation (DSS) follows the requirements of the National Shellfish Sanitation Program (NSSP), which is regulated by the U.S. Food and Drug Administration. The NSSP classification uses the shoreline survey as its primary tool for classifying shellfish growing waters. Fecal coliform concentrations in seawater samples collected in the immediate vicinity of the shellfish beds function to verify the findings of the shoreline surveys, and to define the border between approved and condemned (unapproved) waters.

DSS uses the shoreline survey to locate as many sources of pollution as possible on the watersheds of shellfish growing areas. DSS conducts a property-by-property inspection of the onsite sanitary waste disposal facilities of many properties on un-sewered sections of watersheds, and investigates other sources of pollution such as wastewater treatment facilities (WWTF), marinas, livestock operations, landfills, etc. The information is compiled into a written report with a map showing the location of the sources of real or potential pollution found, and sends it to the various state agencies that are responsible for regulating these concerns and the city or county. The local health departments (LHDs) of the Virginia Department of Health (VDH) play a major role in the process by obtaining correction of the onsite sanitary waste disposal problems. Most of the Division's shoreline survey effort is focused on locating potential fecal contamination, and in this manner we prevent significant amounts of human pathogens from getting into shellfish waters. We believe that this is a primary reason why we have not had a confirmed shellfish-borne disease outbreak due to Virginia-grown shellfish since the early 1960's. VDH is reducing the input of these pathogens to back yards, waterways, unofficial swimming areas and shellfish waters. The shoreline survey work is the foundation of the shellfish growing area classification program.

In addition to the shoreline survey, the NSSP requires that DSS collect seawater samples in the growing areas as part of the classification procedure. States must use the most recent 30 samples, collected randomly with respect to weather (scheduled one month in advance), to classify a station. The two-part standard for fecal coliforms in waters for direct shellfish harvest to market is a geometric mean no greater than 14 MPN fecal coliforms/100 ml and an estimated 90<sup>th</sup> percentile no greater than 31. Exceeding either number requires closure of that station.

To a lesser degree, the Division collects shellfish samples from sentinel growing areas and has them analyzed for heavy metals and chlorinated hydrocarbons (pesticides and PCBs). Such toxic substances in shellfish are not a public health threat in Virginia's waters, with the potential exception of the Southern Branch of the Elizabeth River and perhaps Little Creek, both of which are located in the Hampton Roads area.

Thus, classification based on fecal pollution is a multi-layered and multi-step process. Initially one uses the shoreline survey to determine if there are any actual or potential sources of fresh fecal pollution to the growing area. If so, then the area cannot be used for the direct harvest of shellfish for marketing. Hampton Roads is an example. Most of Hampton Roads is permanently closed, due to the tremendous amount of shipping and the concern of contamination from treated sewage outfalls and runoff from the urban watershed. However, microbiological results are generally acceptable.

Another example of actual or potential pollution that requires closure is a discharge, such as from a WWTF or the potential discharge from boats in marinas. DSS uses relatively simple computer models developed by VIMS, which employ fairly sophisticated mathematics, to determine the size of buffer zones around these sources. These models use inputs of fecal coliforms (estimated from sewage treatment facility outfall volumes or factors related to the number and size of boats in marinas), die-off factors, and readily available tidal current and channel configuration information. Buffer zones around marinas are typically only in effect during the warmer boating months (April 1 -

*Final Guidance for 2024 IR*

*Appendix C*

October 31), whereas those around WWTF are in effect all year. Once these buffer zones are determined, they do not change in size unless the capacity of the WWTF or the marina changes.

Our third layer of classification, and our most common in Virginia, consists of evaluating areas that are not affected by urban runoff or significant wastewater discharges. One must evaluate the watershed for the potential impacts of known failing onsite sanitary waste facilities to estimate whether their input could be of such a magnitude as to require closure, even if the water quality data is acceptable. If the impact from these failing systems does not appear to pose an undue threat, then the water quality data can be used to verify whether the waters should be classified as approved or not.

Since DSS collects approximately 9-10 seawater samples annually, this means that our geometric mean typically incorporates data reaching back 2.5 to 3 years. Heavy rainfall or very high tides due to winds or moon phase can wash unusually high concentrations of fecal coliforms into shellfish growing areas that can increase the geometric mean or the 90<sup>th</sup> percentile beyond the allowed standard. As more data is collected and the unusually high concentrations fall off the trailing end of the data set, the water quality then appears to improve. This is one of the factors that can cause a continual fluctuation in the classification of the water quality at the interface between impacted upstream waters and the relatively unaffected downstream water body.

Since DSS is not a research organization, we cannot do much to determine the cause of water quality deterioration in areas. However, the Division has tried over the years to do so, and we have encouraged the Commonwealth to put resources into determining those causes. The Division has rarely found an association between obviously failing septic systems adjacent to growing areas and deteriorating water quality in large bodies of water. We have seen areas where impacts on fecal coliform concentrations in smaller bodies of water occur due to failing onsite sanitary waste disposal systems, but these seem to be rare. This should not be taken to downplay the concern from such failing onsite sanitary waste disposal systems, since even small inputs of fecal coliforms from these systems are quite likely to contain significant concentrations of human pathogens. Indeed, failing onsite sanitary waste disposal systems are one of the types of pollution sources of greatest concern with regard to the consumption of bivalve molluscan shellfish. Drainfields located in seasonally high water tables may contribute significant numbers of fecal coliforms to impact water quality, and research into this potential source is needed.


Virginia's urban suburban watersheds like the Lynnhaven River in Virginia Beach are clearly impacted by the flushing action of rapid runoff from storm drains. Other areas are much less predictable. Sometimes heavy rainfalls cause elevated counts in rural areas and sometimes they do not. While the Division used to depend upon rain gauges at airports and other widely scattered locations, it now uses NOAA Doppler predicted rainfall, which provides much improved data during spotty summertime thunderstorms. We may find that with this new data, that some areas respond more predictably to rainfall events than was apparent in the past.

In more rural areas the wildlife component of fecal coliform inputs is significant, as can be the human input. Wildlife, such as raccoons, muskrats and deer, living near the intertidal zone, can have dramatic local impacts on fecal coliform concentrations in the adjacent shellfish waters, with the attendant possibility of introducing human pathogens. New data indicates that wildfowl can have significant impacts on water quality too. Wildlife inputs of fecal material are basically accounted for by the seawater sampling data.

The Division is not seeing a steady increase in the number of acres of condemned waters in the state. Instead, what we see are fluctuations in the location of the border between acceptable and unacceptable water quality measurements moving up and down tributaries over time. Again, these fluctuations seem to be due largely to changing factors on the watershed, chance weather events (rain, high tides), changes in wildlife populations near shore or unknown factors (perhaps movement of livestock from one field to another, migratory bird flocks, or runoff from recently plowed fields that later contribute little when crops stabilize the soil).

Man does directly impact the fecal coliform counts in the waters. The headwaters of smaller streams are impacted by development due to the loss of the filtering and detention of runoff waters through upland swamps and other slow moving water areas. These natural detention areas provide the extended time element so that predators (*e.g.*, rotifers and ciliates) and sunlight can reduce the numbers of fecal coliforms and pathogenic human bacteria and viruses. When these are replaced with drainage systems the fecal coliforms and potentially present human pathogens are directly discharged into the shellfish waters.

EPA Shellfish Listing /Delisting Chart  
**Making §303(d) & §305(b) CWA Listing Decisions Based on National Shellfish Sanitation Program (NSSP) Growing Area Classifications**



\* 2007 NSSP Model Ordinance Subsection IV.@03: Growing Area Classification

## **APPENDIX D-1**

### **Incorporating the *Proactive Approach* to delisting 303(d) listed segments into the 2024 Water Quality Assessment**

For the 1998 assessment cycle, EPA changed the data analysis period for the 305(b) assessment from two to five years. Virginia's water quality assessments and the subsequent 303(d) list have since been based on a 5-year data window. In 2008 the assessment data window was expanded to six years to coincide with the two-year ambient watershed rotation monitoring schedule. The data window for 2024 is January 1, 2017 through December 31, 2022.

In August 2001, the Office of Water Quality Programs negotiated with EPA an approach, termed the *Proactive Approach*, which results in the proposed delisting of waters on the Section 303(d) list through assessment of less than six years of data. Correspondence and information related to the issue is attached to this memorandum. In short, EPA Region III has consented that Virginia can delist a segment on the 303(d) list if the following requirements are met:

- 1) For conventional parameters, no more than one of twelve samples taken over a two-year period exceeds the water quality criteria ( $\leq 10.5$  percent exceedance for larger data sets).
- 2) For biological impairment, a minimum of two consecutive samples, taken over a one to two year period, show attainment of the applicable standard.
- 3) The samples are taken at the same location (monitoring station) which demonstrated the impairment.
- 4) A rationale document is submitted to EPA justifying why the State believes the waters are achieving WQ Standards. This rationale document can consist of a description of measures taken in the watershed which are considered to be responsible for improvement of the water quality.

### **Eligibility and Water Quality Assessment**

The following procedure is to be used to consider the eligibility of, and to subsequently assess, any particular waterbody segment submitted for consideration for delisting under the *Proactive Approach*.

Locations where proactive measures are being taken to improve water quality through the TMDL or Water Quality Management Plan program such that the *Proactive Approach* is eligible for consideration are to be provided by the DEQ TMDL program. Assessment staff can recommend segments for consideration, but only those locations provided by the DEQ TMDL program as candidates for the *Proactive Approach* are to be considered for assessment under the *Proactive Approach*. Notification must be made in writing through memorandum to the affected regional assessment manager, copied to the DEQ water quality assessment coordinator, and must include the required documentation supporting consideration of the *Proactive Approach*. At a minimum, this is to include documentation of those implementation measures considered to be responsible for improvement in water quality and subsequent achievement of WQ Standards.

Regional assessment staff members are responsible for assessment of water quality in their respective regions and for the defense of their assessments. Therefore, the decision for delisting consideration is to be made by regional assessment staff based on the analysis of the proactive measures being taken, available monitoring data, any ancillary information collected, and their professional knowledge of site specific influences on water quality in the affected segment.

Where there is agreement between TMDL program and assessment staff that it is appropriate to pursue delisting based on implementation of the *Proactive Approach*, the assessment must be performed based on the

*Final Guidance for 2024 IR*  
*Appendix D-1*

requirements outlined in 1, 2 and 3 above. For a scheduled 305(b)/303(d) assessment, only the last two years of the assessment window are to be used for assessment of eligible segments. For delisting assessment at any other time, all years of the assessment window are to be used.

**Assessment Documentation and Delisting Procedure**

Assessment, Total Maximum Daily Load (TMDL) Tracking and Implementation System (ATTAINS)	A segment meeting the above criteria is considered monitored, fully supporting. The assessment comments section should include the phrase <i>Proactive Approach Assessment</i> . The <i>Proactive Approach</i> data window used must be specifically identified.
Delisting Documentation	Documentation must include the information provided by the TMDL program related to control measures implemented using the <i>Proactive Approach</i> (requirement 4, above), and the results of data analysis related to requirements 1, 2, and 3 above.
EPA Review, Approval and Public Participation	Fulfillment of EPA review and approval requirements, and fulfillment of public participation requirements for removal of waterbody segments (delisting) at EPA required 303(d) list submittal dates, is the responsibility of the Monitoring and Assessments Program. At other times, fulfillment of these requirements in an effort to delist waters not needing TMDLs is the responsibility of the TMDL program. Final documentation for segments delisted by the TMDL program staff must be provided to the regional assessment manager and copied to the DEQ 305(b) coordinator at least five months prior to any EPA required 303(d) list submittal date, if time permits.

## **APPENDIX D-2**

### ***Requirements for Category 5R Waters***

EPA specifically recommends that the Category 5R documentation describe the following six minimum elements:

- a) *The identification of the point and nonpoint sources.* For point sources, an analysis should be included to document whether they are causing or contributing to the water quality impairments. If it is determined that the point sources are causing or contributing, then a Water Quality Based Effluent Limitation (WQBEL) or Best Management Practices Approach<sup>8</sup> should be developed and implemented through NPDES permits.
- b) *The point source and nonpoint source water quality restoration activities that are expected to result in water quality improvements and restoration.* Where applicable, describe any authorities that may require water quality controls to be implemented (e.g., state or local regulations, permits, contracts and grant/funding agreements).
- c) *Cost estimates and funding commitments to implement the water quality restoration activities.* In order to provide assurance that water quality restoration can occur through the implementation of water quality restoration activities, cost estimates and secured funding sources that will be used to implement these activities should be identified.
- d) *An anticipated schedule for implementing the water quality restoration activities, including the anticipated completion date and the estimated pollutant load reductions necessary to meet water quality standards.* The schedule should outline specific activities and include a timeline of when each phase will be implemented and accomplished. The schedule can be revised and updated at each 303(d) listing cycle.
- e) *A water quality monitoring component to evaluate and track the effectiveness of the scheduled water quality restoration activities at each 303(d) listing cycle.* Baseline water quality conditions should be established in order to accurately measure water quality progress. At each 2-year 303(d) listing cycle, performance measurements, whether environmental, programmatic, or social, should be provided for each implemented water quality restoration activity to measure progress. It is understood that each water restoration activity may not result in improved water quality; however the combined restoration activities should result in improved water quality at each 303(d) listing cycle.
- f) *An anticipated date for achieving water quality standards.* Projects are expected to follow adaptive management allowing critical milestones to be adjusted as project plans and goals may change as implementation occurs. Once water quality standards have been met, the State may determine that the waterbody is appropriate to be included in category 1 or 2. If the project does not meet water quality standards by the estimated completion date, sufficient trends toward improved water quality must be shown in order to continue in the 5R program and an updated implementation schedule including revised critical milestones should be submitted to EPA. The project will continue to be reviewed every 2-year 303(d) listing cycle until water quality standards are met.

---

<sup>8</sup> EPA currently recommends point sources be addressed with WQBEL, but DEQ intends to explore how BMPs can also be effectively employed.



## APPENDIX E-1

<b>FISH TISSUE VALUES (TV)*</b>		<b><u>NON CARCINOGEN</u></b>	<b><u>CARCINOGEN</u></b>
		<b>CRITERION BASED TISSUE VALUE (TV)</b>	<b>CRITERION BASED TISSUE VALUE (TV)</b>
<b>COMPOUND</b>	<b>CAS #</b>	<b>PPB (wet-weight)</b>	<b>PPB (wet-weight)</b>
Acenaphthene	83-32-9	44,000	
Acrolein	107-02-8	360	
Acrylonitrile	107-13-1		67
Aldrin	309-00-2		2.1
Anthracene	120-12-7	220,000	
Antimony	7440-36-0	580	
Benzene	71-43-2		660
Benidine	92-87-5		0.2
Benzo(a)anthracene	56-55-3		50
Benzo(b)fluoranthene	205-99-2		50
Benzo(k)fluoranthene	207-08-9		500
Benzo(a)pyrene	50-32-8		5
Bis2-chloroethyl ether	111-44-4		33
Bis2- chloroisoproply ether	108-60-1	29,000	
Bis2- ethylhexyl Phthalate	117-81-7		2,600
Bromoform	75-25-2		8,100
Butyl benzyl phthalate	85-68-7		19,000
Carbon tetrachloride	56-23-5		520
Total Chlordane	57-74-9		100
Chlorobenzene	108-90-7	15,000	
Chlorodibromomethane	124-48-1		910
2-Chloronaphthalene	91-58-7	230,000	
Chloroform	67-66-3	7,000	
2-Chlorophenol	95-57-8	3,600	
Chrysene	218-01-9		5,000
Cyanide	57-12-5	440	
DDD	72-54-8		170
DDE	72-55-9		120
Total DDT	50-29-3		120
Dibenz(a,h)anthracene	53-70-3		5.0
1,2-Dichlorobenzene	95-50-1	220,000	
1,3-Dichlorobenzene	541-73-1	7,300	
1,4-Dichlorobenzene	106-46-7	51,000	
3,3-Dichlorobenzidine	91-94-1		81
Dichlorobromomethane	75-27-4		1,100
1,2-Dichloroethane	107-06-2		11,000
1,1-Dichloroethylene	75-35-4	36,000	
1,2-Trans-dichloroethylene	156-60-5	15,000	
2,4-Dichlorophenol	120-83-2	2,200	
1,2-Dichloropropane	78-87-5		1,000
1,3-Dichloropropene	542-75-6		300

Dieldrin	60-57-1		2.3
Diethyl phthalate	84-66-2	580,000	
2,4-Dimethylphenol	105-67-9	15,000	
Dimethyl Phthalate	131-11-3	7,300,000	
Di-n-butyl phthalate	84-74-2	360,000	
2,4-Dinitrophenol	51-28-5	1,500	
2-Methyl-4,6-dinitrophenol	534-52-1	220	
2,4-Dinitrotoluene	121-14-2		54
Dioxin	1746-01-6		0.00023
1,2-Diphenylhydrazine	122-66-7		45
Endosulfan (I and II)	115-29-7	4,400	
Endosulfan sulphate	1031-79-8	4,400	
Endrin	72-20-8	870	
Endrin aldehyde	7421-93-4	870	
Ethylbenzene	100-41-4	16,000	
Fluoranthene	206-44-0	29,000	
Fluorene	86-73-7	29,000	
Heptachlor	76-44-8		8.9
Heptachlor epoxide	1024-57-3		6.6
Hexachlorobenzene	118-74-1		36
Hexachlorobutadiene	87-68-3		910
Hexachlorocyclohexane (alpha-BHC)	319-84-6		5.8
Hexachlorocyclohexane (beta -BHC)	319-85-7		20
Hexachlorocyclohexane (gamma-BHC) (lindane)	58-89-9	8,500	
Hexachlorocyclopentadiene	77-47-4	4,400	
Hexachloroethane	67-72-1		2,500
Indeno(1,2,3-cd)pyrene	193-39-5		50
Isophorone	78-59-1		38,000
Mercury (Methyl) **	22967-92-6	300	
Methyl Bromide	74-83-9	15,000	
Methylene Chloride	75-09-2		18,000
3-Methyl-4-chlorophenol	59-50-7	73,000	
Nickel	744-00-2	73,000	
Nitrobenzene	98-95-3	1,500	
N-nitrosodimethylamine	62-75-9		0.71
N-nitrosodiphenylamine	86-30-6		7,400
N-nitrosodi-n-propylamine	621-64-7		5.2
PCB Total/congeners	1336-36-3		18
Pentachlorophenol	87-86-5		91
Phenol	108-95-2	440,000	
Pyrene	129-00-0	22,000	
Selenium	7782-49-2	18,000	
1,1,2,2-Tetrachloroethane	79-34-5		180
Tetrachloroethylene	127-18-4		17,000
Thallium	7440-28-0	49	
Toluene	108-88-3	7,100	
Toxaphene	8001-35-2		33
1,2,4-Trichlorobenzene	120-82-1		1,300
1,1,1-Trichloroethane	71-55-6	15,000,000	

1,1,2-Trichloroethane	79-00-5		640
Trichloroethylene	79-01-6		730
2,4,6-Trichlorophenol	88-06-2		3,300
Vinyl Chloride	75-01-4		24
Zinc	7440-66-6	1,100,000	

\*These fish tissue values have been calculated based on the Water Quality Standards that are associated with the latest Triennial Review criteria proposals as adopted by the State Water Control Board in October 2019. Additionally, the values reflect the current default values for fish consumption rate (22 g/day) and body weight (80 kg) recommended by USEPA.

\*\*The fish tissue criterion for methylmercury applies to fish species commonly eaten in the local waterbody and applies to most fish species in the DEQ database except bowfin or longnose gar because fish consumption surveys show that these species are rarely consumed in Virginia. Total mercury concentrations in fish tissue are assumed to equal methylmercury concentrations.

**APPENDIX E-2**

**RISK-BASED TISSUE SCREENING VALUE (TSVs) FOR FISH TISSUE DERIVED FROM REFERENCE DOSE AND ORAL SLOPE FACTORS PUBLISHED BY EPA FOR GENERAL POPULATION (ADULT)**

BODY WEIGHT (KG) 80  
 RISK LEVEL 10<sup>-5</sup>  
 CONSUMPTION RATE (KG/DAY) 0.022

Fish Tissue Screening Values (TSV)		<u>NON</u> <u>CARCINOGEN</u>	<u>CARCINOGEN</u>
		TISSUE SCREENING VALUE (TSV)	TISSUE SCREENING VALUE (TSV)
COMPOUND	CAS #	PPB (wet-weight)	PPB (wet-weight)
Aluminum	7429-90-5	3,600,000	
Arsenic (inorganic)	7440-38-2		240*
Barium	7440-39-3	730,000	
Beryllium	7440-41-7	7,300	
BHC isomers (Technical HCH)	<del>608-93-1</del> 608-73-1		20
Brominated Diphenyl ethers (BDEs)		500 (VDH)**	
Cadmium	7440-43-9	3,600	
Decabromdiphenyl ether	1163-19-5	25,000	
Hexabromodiphenyl ether	<del>36483-60-0</del> 68631-49-2	730	
Pentabromodiphenyl ether	32534-81-9	7,300	
Chromium III	16065-83-1	55,000	
Chromium VI	18540-29-9	11,000	
Chlorpyrifos	2921-88-2	1,100	
Copper	7440-50-8	36,000	
Diazinon	333-41-5	720	
Disulfoton	298-04-4	150	
Ethion	563-12-2	1,800	
Kepone	143-50-0		300 (VDH)**
Manganese	7439-96-5	510,000	
Methoxychlor	72-43-5	18,000	
Mirex	2385-85-5	730	
Oxyfluorfen	42874-03-3	11,000	
PAHs (sum PEC) ***			5.0
Silver	7440-22-4	18,000	
Terbufos	13071-79-9	180	
Tributyltin	56-35-9	1,100	
Vanadium	7440-62-2	36,000	

\*The screening value for arsenic applies to inorganic arsenic only. Organic forms of arsenic are not carcinogenic and are relatively nontoxic. There is a general consensus that 85 to 90% of arsenic found in fish tissue is organic arsenic. The screening value of 240 ug/kg total arsenic is based on the estimate that 10% of total arsenic detected in fish tissue is inorganic arsenic.

\*\* These values are based on recent changes to the toxicological data used to calculate the screening values, or recent recommendations from U.S. EPA or the Virginia Department of Health. These screening values are not based on the same toxicological data that were used to develop the existing water quality criteria.

\*\*\* Mixtures of seven polynuclear aromatic hydrocarbons (PAHs) that are classed as probable human carcinogens were assessed based on a screening value concentration of 5.0 ppb calculated as a sum potency equivalency concentration (PEC) using methods described in EPA's Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories, Vol. 1, (EPA 823-R-95-007) and Vol. 2 (EPA 823 B-00-008) using the following equation;

Final Guidance for 2024 IR  
 Appendix E-2

$$PEC = \sum (RP_i \times C_i)$$

where;

$RP_i$  = relative potency for the  $i$ th PAH

$C_i$  = concentration of the  $i$ th PAH in fish tissue)

The relative potency estimates used for these PAHs were:

Benzo(a)pyrene	1.0
Benzo(a)anthracene	0.145
Benzo(b) fluoranthene	0.167
Benzo(k)fluoranthene	0.020
Chrysene	0.0044
Dibenz(a,h)anthracene	1.11
Indeno(1,2,3-cd)pyrene	0.055

## APPENDIX F

<b>Freshwater Consensus- Based Sediment Screening Values (SVs)</b>	
<b>Analyte (Metals)</b>	<b>Consensus PEC (ppm) dry weight</b>
Arsenic	33
Cadmium	4.98
Chromium	111
Copper	149
Lead	128
Mercury	1.06
Nickel	48.6
Silver	NA
Zinc	459
<b>Analyte (Organics/Pesticides)</b>	<b>Consensus PEC (ppb) dry weight</b>
Acenaphthene	NA
Acenaphthylene	NA
Anthracene	845
Benzo-a-pyrene	1,450
Benz(a)Anthracene	1,050
Chrysene	1,290
Dibenz[a,h]Anthracene	NA
Fluoranthene	2230
Fluorene	536
Methylnaphthalene, 2-	NA
Naphthalene	561
Phenanthrene	1,170
Pyrene	1,520
LMW PAHs	NA
HMW PAHs	NA
Total PAHs ** (see footnote)	22,800
Chlordane	17.6
DDD	28
DDE	31.3
DDT	62.9
DDT, total	572
Dieldrin	61.8
Total PCBs	676
Endrin	207
Heptachlor Epoxide	16
Lindane	4.99
NA = Not Available	

<b>Estuarine NOAA-based ER-M Sediment Screening Values (SVs)</b>	
<b>Trace Elements (Metals)</b>	<b>ER-M Value ppm (dry weight)</b>
Antimony (Sb)	NA
Arsenic (As)	70
Beryllium	NA
Cadmium (Cd)	9.6
Chromium (Cr)	370
Copper (Cu)	270
Lead (Pb)	218
Manganese (Mn)	NA
Mercury (Hg)	0.71
Nickel (Ni)	51.6
Selenium (Se)	NA
Silver (Ag)	3.7
Thallium	NA
Zinc (Zn)	410

<b>Pesticides and Other Organic Substances –parts per billion dry weight</b>		
<b>CAS #</b>	<b>Substance</b>	<b>ER-M Value(dry weight) (ppb)</b>
336363	Polychlorinated Biphenyls (PCBs)	180
309002	Aldrin	NA
57749	Chlordane	6
NA	total DDT (include metabolites)	46.1
72548	DDD	20
50293	DDT	7
72559	DDE	27
60571	Dieldrin (EPA proposed criteria)	8
72208	Endrin	NA
76448	Heptachlor	NA
1024573	Heptachlor epoxide	NA
118741	Hexachlorobenzene	NA
608731	Hexachlorocyclohexane	NA
58899	Lindane	NA
2385855	Mirex	NA
108952	Phenol	NA
117817	Di (2-Ethylhexyl) Phthalate	NA
84742	N-Butyl Phthalate	NA
83329	Acenaphthene	500 LMW PAH
208968	Acenaphthylene	640 LMW PAH
120127	Anthracene	1100 LMW PAH
50328	Benzo-A-Pyrene	1600 HMW PAH
191242	Benzo [GHI] Perylene	NA HMW PAH
56553	Benz[A] Anthracene	1600 HMW PAH
218019	Chrysene	2800 HMW PAH
53703	Dibenz [A,H] Anthracene	260 HMW PAH
206440	Fluoranthene	5100 HMW PAH
86737	Fluorene	540 LMW PAH
193395	Indeno (1,2,3-CD)Pyrene	NA HMW PAH
91576	Methylnaphthalene, 2	670 LMW PAH

<b>Pesticides and Other Organic Substances –parts per billion dry weight</b>		
<b>CAS #</b>	<b>Substance</b>	<b>ER-M Value(dry weight) (ppb)</b>
91203	Naphthalene	2100 LMW PAH
85018	Phenanthrene	1500 LMW PAH
129000	Pyrene	2600 HMW PAH
NA	Low Molecular Weight (LMW)PAHs	3160
NA	High Molecular Weight (HMW) PAHs	9,600
NA	Total PAHs **(see footnote)	44,792

\*Changes or updates to any of the ER-M or PEC screening values should be updated in the assessment spreadsheet used to calculate the estuarine weight of evidence.

\*\*sum of 24 Polyaromatic hydrocarbons used in previous reports, also polynuclear aromatic hydrocarbons (PNAs)

DEQ acknowledges the use of the ER-M or PEC may be limited (for several reasons) in their ability to accurately predict biological effects. Given that DEQ continues to employ the collection of bulk sediment with chemical analysis as a cost-effective way to monitor a great number of sediment sites, these thresholds are an appropriate tool for assessing sediment data relative to its potential harm to aquatic life.

**Citation:**

**Freshwater PECs:** MacDonald, D.D., C.G. Ingersoll, T.A. Berger. 2000. Development and Evaluation of Consensus-Based Sediment Quality Guidelines for Freshwater Ecosystems. *Arch. Environ. Contam. Toxicol.* 39:20-31.


**Estuarine ER-Ms:** Buchanan, M.F. 1999 National Oceanic and Atmospheric Administration *Screening Quick Reference Tables*, NOAA HAZMAT Report 99-1 Seattle, WA, Hazardous Materials Response and Assessment Division, 12 pages.



## APPENDIX G

### Weight-of-Evidence (WOE) Aquatic Life Use Assessment in Estuarine Waters

The “Weight-of-Evidence” (WOE) approach that DEQ currently uses for its general evaluation and assessment of the designated Aquatic Life Use (ALU) for estuarine benthic communities has evolved from a previously more limited application of the “Sediment Quality Triad” concept (SQT – Figure 1). The SQT concept was originally conceived and applied for the evaluation of the presence and effects of toxic contaminants in marine sediments (Long and Chapman 1985). It was further applied by Chapman et al. (1986, 1987), and has continued to be one of the preferred approaches for the evaluation of toxics in marine and estuarine benthic environments (Chapman 1992, Chapman et al. 1997, McGee et al. 2001). The Interstate Chesapeake Bay Program (CBP) employed SQT evaluations along with other methods to produce a Bay-wide toxics characterization in 1999 (US EPA 1999) that identified (1) “Regions of Concern – areas with probable adverse effects,” (2) “Areas of Emphasis – areas with potential adverse effects,” (3) “Areas with Low Probability for Adverse Effects”, and (4) “Areas with Insufficient or Inconclusive Data” relative to toxics contamination in Bay waters. Maps of more recent characterizations (2006 through 2014) can be found by using the Search Function for “Chemical Contaminants” at: <http://www.chesapeakebay.net/maps.aspx?menuitem=15230>.



**Figure 1 - The Sediment Quality Triad (SQT = triangle A.B.C.) as originally conceived for the identification and characterization of potential toxics-induced stressors.**

The apex of the triangle, Circle A or “Benthos,” represents the condition of the benthic community, which is the primary objective of the “Aquatic Life Use” assessment, while B - “Sediment Chemistry” and C - “Sediment Toxicity” provide two lines of evidence for the evaluation of possible causes of stress due to toxic contamination. Tools for the evaluation of benthic condition (D – “Benthic Indices”) and “Additional Stressors” (E), as well as Sediment Chemistry (B), and Sediment Toxicity (C), are discussed in the text.

Subsequent to the 1999 CBP characterization, DEQ, in conjunction with researchers from the Virginia Institute of Marine Science (VIMS), used the SQT for the characterization of those Virginia Bay waters that had been identified as Class 4 (Insufficient or Inconclusive Data) in the previous CBP study, namely the tidal fresh regions of the James River, the tidal York River drainage, and Mobjack Bay (Roberts et al. 2002a, 2002b,

2003). A newer report on toxics throughout the Chesapeake Basin was published in December 2012 (EPA et al., 2012).

The original objective of such ambient toxics monitoring was primarily to perform a quick screening of the medium of interest (water, sediment, fish tissue, etc.) to determine whether toxic pollutants were present and could potentially have a negative impact on aquatic life or human health. In addition to the evaluation of potential *causes* of impact (based on Sediment Chemistry – element B), potential toxic *effects* on the biota (including individual survival, growth and/or reproduction) could be evaluated based on the results of toxicity tests (Sediment Toxicity – element C), most commonly conducted in the laboratory but at times carried out with test species maintained *in situ* (e.g., Roberts et al. 2002a, 2002b, 2003). The general welfare of the benthic community (Benthos – element A) was evaluated as a manifestation of elements B and C if they were positive (i.e., actual *observed effects* of sediment contamination). Although the potential effects of other stressors were acknowledged, they did not play a significant role in the earlier SQT evaluations.

Because sediment chemical contamination and its resultant toxicity are relatively stable through time, they are much more appropriate for characterizing probabilistic sites (that are normally only visited once) than are water quality parameters, which may vary on much shorter time scales (seasonally, daily, hourly, or minute-to-minute). The condition of the benthic infaunal community reflects long-term (and potentially chronic) effects from sediment chemical contamination as well as from a variety of other stressors.

Virginia DEQ began to apply a modified, more formal WOE assessment procedure employing the SQT in its 2006 Integrated 305(b)/303(d) Water Quality Assessment Report. DEQ’s assessment procedure, however, goes beyond the original SQT toxics-related evaluations and includes tools for the tentative evaluation of some of the additional potential stressors (E – “Additional Stressors” of Figure 1) affecting estuarine benthic communities. WOE assessment is carried out on data collected within DEQ’s Estuarine Probabilistic Monitoring (ProbMon) Program and, periodically, the National Coastal Condition Assessment (NCCA) surveys which sample the coastal Delmarva region, the Back Bay / North Landing River region, and the tidal tributaries, embayments, and mainstem of the Chesapeake Bay drainage. Because all three elements of the SQT are collected and water quality and additional sediment analyses are carried out simultaneously, the WOE procedure is able to provide an integrated assessment for individual sampling sites. General guidance for the delineation of the area represented by each site is provided in Section 5.1, “*Monitoring Station Siting and Delineation*”, Rules 4 and 6, of this Assessment Guidance Manual. Note that within the tidal portions of the Chesapeake Bay basin, the WOE assessment discussed in this section complements the probabilistic benthic assessments carried out by Virginia’s Chesapeake Bay Program (CBP). The probabilistic benthic monitoring carried out by the CBP collects benthic samples and a few measures of bottom conditions at each site (sediment type and total organic carbon [TOC] content, salinity, dissolved oxygen, etc.), but does not carry out chemical analyses or toxicity tests of sediment. Consequently, the results of CBP benthic characterizations are spatially integrated and assessments are performed only on pre-designated Bay segments that have a sufficiently large benthic sample size ( $N \geq 10$ ).

The SQT is an effects-based approach that describes the condition of the sediment and associated benthic infaunal communities relative to toxic pollutants and their effects. The three main data components that were integrated into the original SQT analysis included: (1) sediment bulk chemical concentrations, (2) sediment toxicity test results, and (3) an evaluation of benthic infaunal community condition. Rather than considering each type of characterization individually, the complementary methods integrate biological responses with chemical data (Chapman 1992) for a more scientifically defensible assessment process. Chapman (1992) provided eight possible scenarios from which conclusions could be drawn with the SQT approach. It was this concept that first served as the foundation required to implement the WOE assessment of SQT data as summarized in the evaluation matrix of Figure 2.

The objective of this guidance is to provide orientation for interpreting data generated by the traditional SQT approach, as complemented by additional lines of evidence, with added insight on how to consider “weighting” of each component. This is not to suggest that sound scientific interpretation and best professional judgment are unnecessary, but does provide some degree of standardization for the process. Conceptually, this is similar to the approach used by the Chesapeake Bay Program and its partners for its Toxics Characterization of the Bay (EPA 903-R-00-010, June 1999). The use of this guidance will provide assistance in applying “weights” to the different triad components, which are then inserted into the classic SQT matrix. The bulk chemistry results, for example, can receive additional weighting based on the magnitude of exceedance of the applicable Sediment Quality Guideline (SQG). For toxicity tests, greater weight is applied with two or more statistically significant sediment toxicity tests than is applied to a single significant test, for the affected endpoint. The type of toxicity test endpoints that exhibit statistical significance must also be given consideration, since the acute ecological consequence of not surviving would be greater on the population of a species than the rate of growth of individuals. Test results based on survival/mortality would therefore receive a greater weight.

Scenario	Chemistry	Toxicity	Benthic Community Alteration <sup>9</sup>	Taxonomic/Community Summary	Listing Category
1	Score > 0	Score > 0	0 < Score < 1	Potential for chemical contaminant-induced degradation.	3B
			Score > 1	Evidence of chemical contaminant-induced degradation	5A
2	0	0	0	No evidence of chemical contaminant-induced degradation.	2A
3	Score > 0	0	0	Chemical contaminants exhibit potential to cause degradation.	3E
4	0	Score > 0	0	Chemical contaminants or conditions (including those unmeasured) may exist that have potential to cause degradation.	3E
5	0	0	Score > 0	No evidence for chemical contaminant-induced degradation.	3B
6	Score > 0	Score > 0	0	Potential for chemical contaminant-induced degradation.	3E
7	0	Score > 0	0 < Score < 1	Potential for chemical contaminant-induced degradation.	3B
			Score > 1	Evidence for chemical contaminants or conditions (including those unmeasured) causing degradation.	5A
8	Score > 0	0	0 < Score < 1	Potential for chemical contaminant-induced degradation.	3B
			Score > 1	Evidence for chemical contaminants or conditions (including those unmeasured) causing degradation.	5A

**Figure 2 - The SQT Evaluation Matrix summarizing the eight scenarios originally described by Chapman (1992).**

The “SQT Evaluation Matrix” DEQ uses (Figure 2) has been adapted from the original (Chapman 1992) in order to incorporate additional lines of evidence and to provide numerical scores for the three classes of characterization: Chemistry, Toxicity, and Benthic Community Alteration. The column “Tentative/Possible Conclusions” summarizes Chapman’s (1992) descriptions of the eight possible scenarios with the addition of some comments on possible assessment classifications, and the last column lists the specific listing categories that may be assigned for Virginia’s 305(b)/303(d) Integrated Water Quality Assessment Report. Further

<sup>9</sup> Natural conditions may be present that cause atypical benthic communities and result in low index scores (e.g. high-flow areas with sand habitat, indicating erosional conditions). Under such conditions, the Estuarine ProbMon Coordinator may assign category 3 under scenarios 5, 7 and 8. In such cases, this recommendation should be accompanied with a narrative explanation and any accompanying data that provides support for natural degradation of the benthic community.

discussion of the matrix is provided in the text sections below related to the Microsoft Excel® “Weight-of-Evidence Assessment Workbook.”

The Weight-of-Evidence Workbook consists of 13 individual worksheets that fully document the location of the monitoring site, the complete analytical results of sediment chemical analyses, of sediment toxicity tests, and of benthic identifications and enumerations, the interpretation of those results, and the integration of all lines of evidence (including water quality, sediment quality and benthic community well-being) into a final assessment:

<b>Page No.</b>	<b>Tab Title</b>
1.	Orientation - Read Me First
2.	Summary Sheet
3.	SQT Evaluation Matrix
4.	(1) SedChem – Mean ERM Quotient Model
5.	(2) SedChem – Logistic Regression Model
6.	(3) SedChem – Equilibrium Partitioning Sediment Benchmark for PAH Mixtures
7.	(4) Sediment Toxicity
8.	(5) Benthic Infauna
9.	Sediment Chemistry Data
10.	Chemistry QA Codes
11.	Sediment Toxicity Data
12.	Benthic Data
13.	Site Map (USGS TopoQuad) and Satellite Imagery

In keeping with DEQ’s continual planning process, the WOE assessment procedure and the format of the Weight-of-Evidence Assessment workbook continue to evolve as additional lines of evidence and new assessment tools become available. Page 5 “(2) Sediment Chemistry – Logistic Regression Model” was added in 2015, for the assessment of 2014 results, following its inclusion in the 2014 draft Technical Appendix of National Coastal Condition Assessment 2010 (US EPA, 2015) by the EPA Office of Water and Office of Research and Development. The following discussions will describe the various lines of evidence considered and summarize the characterization procedures as currently employed (January 2019). More details and explicit instructions relative to each step are included within the Weight-of-Evidence Workbook.

### **Workbook Summary Sheet - Final Assessment and Comments:**

Descriptive information identifying the specific sampling site is entered into this workbook Tab prior to adding additional results of field measurements and laboratory analyses there or elsewhere in the file. As on other tabs throughout the workbook, information should be typed or pasted into fields that are highlighted in pastel green. Fields highlighted in pale yellow should not be altered in any way. Yellow fields are populated automatically by embedded calculations or direct transfers from other fields within the workbook. Entering observations or comments in undesignated fields should be avoided, since many fields contain hidden reference values for populating other cells of the workbook. Once assessment is complete, the assessor may highlight additional cells to classify results as “Good” (bright green), “Fair” (bright yellow), or “Poor” (bright red).

### **Benthic Community Characterization: Workbook Tab 8 - “(5) Benthic Infauna”**

Because the WOE assessment is applied strictly for the designated “Aquatic Life Use” (ALU), evaluation of the structure and function of the benthic community is the ultimate, most heavily weighted indicator for site characterizations. The condition of the benthic community constitutes an integrated *observed effect* (Va. Assessment Category 3B) of the existing environmental stressors, whether the individual stressors are identified

or not. If the benthic community is found to be *severely degraded*, a site may be assessed as “Impaired” for ALU even if evidence for a specific cause is lacking. Additional, conformational sampling would be required, however, prior to initiating TMDL development. In the opposing case, if the benthic community was found to be in good condition (“non-degraded” or “meeting goals”), a classification of “(5A) Impaired” would be unlikely unless chemical and/or toxicity results were exceptionally extreme.

The general objective of the weight-of-evidence (WOE) assessment methodology is to integrate multiple lines of evidence, based primarily on sediment analyses, to provide a standardized, objective evaluation of the severity and probable cause(s) of benthic degradation. Individual benthic index scores are subject to sampling error, which results from the great heterogeneity of biological communities as well as from methodological variations in sample collection, handling and analysis. When additional lines of evidence such as significant chemical contamination and/or significant acute or chronic toxic effects corroborate low benthic IBI scores, they serve to identify probable causes of degradation, and consequently to confirm the validity of low benthic scores and to justify an assessment classification of “Impaired” (5A). When low benthic scores are not corroborated by integrative chemical or toxicological measures, additional lines of evidence (e.g., low DO, high nutrient concentrations, evidence of sedimentation, or other habitat characterizations) may contribute to their interpretation. These alternative lines of evidence are of more limited value, however, in the case of single-visit probabilistic sampling.

Within tidal Chesapeake Bay waters the natural variability of benthic communities, both within and among habitat types, is recognized and included in a formal statistically-based procedure which integrates multiple ( $N \geq 10$ ) CBP probabilistic benthic IBI scores for ALU assessment of pre-defined tidal water segments. The CBP benthic IBI scores from DEQ’s Estuarine Probabilistic Monitoring Program (ProbMon) are included in this procedure. In non-Bay tidal waters, where no other benthic community evaluation is carried out, the WOE assessment places greater weight on alternative lines of evidence from the probabilistic sample, including the degree of consensus among the three benthic indices generally calculated. In coastal Delmarva waters and in the Albemarle Sound drainage (Back Bay, North Landing River), greatest weight is given to the Middle Atlantic Benthic IBI. In these waters, if the CBP Benthic IBI and/or the EMAP Virginia Province Index of Estuarine Condition corroborate evidence of severe benthic degradation, an assessment of impaired ALU may result even in the absence of supporting chemical or toxicological evidence.

The number of different benthic taxa present in a standardized sample<sup>10</sup>, their relative abundances, and knowledge of their specific ecological/functional roles provide the information for calculating numerous measures or metrics of community structure and function. Several of these metrics are used individually for a preliminary, general characterization of the benthic community while many of them are subsequently integrated into various, more objective multi-metric indices of biological integrity (Benthic IBIs) or an index of estuarine condition (IEC). Raw taxonomic data and intermediate results are provided on the “Benthic Data” Tab (page 12) of the WOE Workbook, while the integrated evaluation is summarized on Tab “(5) – Benthic Infauna” (page 8).

The values of seven individual metrics derived from the taxonomic results provide an initial qualitative evaluation of benthic condition during the WOE process. The first five are measures of taxonomic abundance

---

<sup>10</sup> The standardized benthic sample for the Estuarine Probabilistic Monitoring Program consists of a composite of two separate sediment grabs using a 6-inch Petite Ponar sediment sampler, representing a total bottom area of approximately 0.046 m<sup>2</sup>. The contents of each grab must conform to quality assurance criteria specified in the National Coastal Condition Assessment program QAPP (U.S. EPA., 2009b, 2014a) and Field Operations Manual (U.S. EPA., 2009c, 2014b) and must include at least 7.0 cm of sediment. The B-IBI results with this sampling protocol have been shown not to differ significantly from standardized samples collected with a single 8” Young sampler (Dauer and Lane, 2005).

and diversity, while the last two are the abundances of two stress-tolerant taxa that are also used in calculating an IEC, to be discussed below.

Metric	Description
1. Total Abundance -	The total number of individuals in a benthic sample; generally symbolized as “N”.
2. Total Taxa -	The total number of taxa that are identified from a sample. Depending upon the group of organisms, an individual taxon may represent a species, a genus, a family, or a higher level of identification. Usually symbolized as “S” for number of species, but “S” is maintained here as the number of taxa.
3. Shannon H' -	The Shannon-Weiner Species Diversity Index: $H' = -\sum_{i=1, S} (p_i \ln p_i)$ . This index is calculated here as it was originally expressed, using $\log_2$ . Elsewhere, however, it is often calculated using natural logs (ln) or occasionally using decimal logs ( $\log_{10}$ ). The use of “S” as defined here would make H' an index of taxonomic diversity rather than of species diversity.
4. Gleason-D -	Gleason’s Diversity Index: $D = S / \ln N$
5. Pielou-J' -	Pielou’s Index of Equitability (or Evenness): $J' = H' / H'_{\max}$ , where $H'_{\max}$ is the theoretical maximum diversity with “N” individuals divided among “S” taxa. The value of J' can vary from 0.0 to 1.0; both H' and $H'_{\max}$ must be calculated to the same logarithmic base; in this case $H'_{\max} = \log_2 (S)$
6. Tubificidae -	The numeric abundance of the family Tubificidae (Annelida, Oligochaete), a stress-tolerant taxon.
7. Spionidae -	The numeric abundance of the family Spionidae (Annelida, Polychaete), another stress-tolerant taxon.

Although the abundance of individuals and the diversity of taxa vary among habitat types - muddy vs. sandy sediment, in combination with salinity regime - within a specific habitat type higher values of metrics 1 through 5 are generally indicative of more healthy, less degraded benthic communities. A high abundance (metric 1) with few taxa (metric 2) may indicate a degraded benthic community, especially if the abundant taxa are stress-tolerant as are those of metrics 6 and 7. This would result in relatively low values for metrics 3, 4 and 5. An excessive abundance of tubificids (6) and/or spionids (7) generally indicates a highly stressed and probably degraded benthic community.

Benthic community alteration is also evaluated by integrating various individual metric scores into a single Benthic Index of Biotic Integrity (B-IBI) based on previously established and verified thresholds (e.g., Chesapeake Bay B-IBI; Weisberg et al., 1997), and then comparing the overall site B-IBI score with the defined ranges characteristic of specific habitat types (e.g., Llansó and Dauer, 2003 for the Chesapeake Bay). Characterizing the overall benthic community condition with this CBP B-IBI is straight-forward, since there are four previously established categories ranging from good to severely degraded (Please refer to **Table 1** below). It is the preferred and most appropriate index for use within the Chesapeake Bay drainage. An alternate B-IBI, developed for estuaries of the Middle Atlantic Region (Llansó et al., 2002a, 2002b), is used for assessment in the Atlantic coastal estuarine waters of the Delmarva Peninsula. For this B-IBI, index values < 3.0 are considered indicative of stressed benthic assemblages and degraded conditions (SQT Matrix Score = 3 or 2), while scores  $\geq 3.0$  indicate that benthic goals are met (Matrix Score = 0).

**Table 1 - Chesapeake Bay B-IBI Ranges and Benthic Community Condition**

<b>CBP B-IBI Score</b>	<b>Benthic Community Condition</b>	<b>SQT Matrix Score</b>
≥ 3.0	Meets Goal	0
2.7-2.9	Marginal	1
2.1-2.6	Degraded	2
≤ 2.0	Severely Degraded	3

When one or more measurements essential for the calculation of either of these B-IBIs is lacking, or if they are considered geographically inappropriate, a third alternative is available. Paul et al. (2001) developed a benthic-based “Index of Estuarine Condition” (VP-IEC) for the Virginian Biogeographic Province (from Cape Cod to the mouth of Chesapeake Bay), based on the 1990-1993 results of EPA’s Middle Atlantic Integrated Assessment (MAIA) Program. This index is given minimum weight when either of the B-IBIs is available and more appropriate, but is more heavily weighted when neither of the B-IBIs is available. In the original publication of the VP-IEC, calculated as a linear discriminant function, final values greater than zero (> 0.000) were interpreted as an indication of non-degraded conditions and values less than zero (< 0.000) were interpreted as an indication of degraded sites. No indeterminate “gray zone” was specified. For the purpose of weight-of-evidence assessment, discriminant scores of this index between -0.1 and +0.1 are considered “marginal.” Although no systematic salinity-induced bias has been demonstrated for any of these indices, it should be noted that all three are notably less reliable in low-salinity habitats, *i.e.*, oligohaline and tidal fresh waters (salinity < 5.0 ppt and < 0.5 ppt, respectively). Approximately 20% of Virginia’s estuarine probabilistic sites sampled between 2001 and 2018 were within this salinity range.

Back Bay and the North Landing River, in southeastern coastal Virginia, fall within the Carolinian Biogeographic Province. They constitute a unique tidal freshwater/oligohaline region that is so isolated from the Virginia Biogeographical Province, Albemarle Sound and the Atlantic Ocean that none of the previously described benthic indices may be completely appropriate. Most of the benthos in this region is more characteristic of freshwater than of tidal estuarine waters. For the purpose of assessment in this region, all three benthic indices are compared and a tentative characterization is based on the relative degree of concordance among them and other individual metrics of species abundance, taxonomic richness, and diversity. In 2017 EPA’s Atlantic Ecology Division (AED) in Narragansett, RI, announced the further development of a new multivariate benthic index for use in U.S. coastal waters (Pelletier et al., 2018). The multivariate AMBI (M-AMBI) is an extension of the AZTI Marine Biotic Index (AMBI – Borja et al, 2000) that has been used extensively in Europe for a number of years. The multivariate U.S. M-AMBI was added to the benthic evaluations of the WOE assessments, on a trial basis, in 2017 and 2018. If it proves to be a useful tool as an independent line of evidence for benthic characterizations, and resources are available for its continued inclusion, it may be incorporated as a permanent addition.

Since the summer of 2005, the separation, taxonomic identification and enumeration of all benthic samples collected within the Estuarine ProbMon Program has been carried out at the Benthic Ecology Laboratory (BEL) at Old Dominion University (ODU) under the auspices of Dr. Dan Dauer. Dr. Dauer is the principal investigator responsible for Virginia’s portion of the Chesapeake Bay Program’s Probabilistic Benthic Monitoring Program. In addition to providing a complete list of all benthic taxa and their abundances (in terms of numbers of individuals and biomass), Mike Lane (BEL/ODU) uses the BEL database to calculate all of the individual metrics required and the final score for each of the benthic and estuarine indices discussed above. In practice, all three (currently four) benthic indices are calculated and evaluated for all benthic samples. The greatest weight is given to the results of the most appropriate index, but the degree of concordance (or disagreement) among them is also considered for the final characterization.

Tab “(5) Benthic Infauna” of the WOE Assessment Workbook summarizes the integrated scoring and weighting for the three (currently four) benthic and condition indices, along with a number of associated habitat and sediment characteristics that contribute to the final characterization of the benthic community. Several of these complementary characteristics (*e.g.*, bottom DO, bottom temperature, sediment TOC, and habitat type – salinity regime and mud or sand substrate, summarized on page 2 “Summary Sheet” ) are helpful in identifying potential causes of any observed benthic degradation. Comments and the final Matrix Score recorded on Tab (5) are subsequently transferred to the “SQT Evaluation Matrix” Tab (page 3) for integration into the final assessment.

### **Sediment Characterization: Workbook Tabs (1), (2), (3) and (4)**

#### **Sediment Chemistry: Workbook Tab “(1) SedChem Mean ERMq Model”** (Mean ERM Quotient Model – page 4)

At the present time, EPA has not yet established specific criteria for toxic contaminants in sediment, and Virginia has not established sediment quality standards against which to assess sediment contamination. Consequently, a site is rarely assessed as impaired based on sediment chemistry alone. However, numerous empirical studies carried out over the past 20 to 25 years have provided “Sediment Quality Guidelines” (SQGs) or “Screening Values” (SVs) that serve to tentatively identify the range of concentrations of specific contaminants or classes of contaminants that are likely to cause adverse effects in benthic communities. Virginia currently employs two sets of screening values to characterize sediments: consensus-based Probable Effects Concentrations (PECs - MacDonald et al., 2000) for freshwater sediments and Effects Range Median (ER-M - Long et al., 1995) concentrations for estuarine and marine sediments. **APPENDIX F** of this Assessment Guidance Manual lists the “Consensus Based and ERM Sediment Screening Values” currently applied in Virginia. They also appear on Tab “(1) SedChem Mean ERMq Model” of the WOE Assessment Workbook, where they are compared to observed sediment contaminant concentrations. Virginia’s Water Quality Standards (“WQS” - 9 VAC 25-260) provide guidelines for the application of these screening values and indicate that in transitional Class II (oligohaline) waters the “more stringent of either the freshwater or saltwater criteria apply.” Section 9 VAC 25-260-140, Subsection C, of the WQS defines specific, fixed zones of transitional Class II waters for Virginia’s major tidal tributaries (Potomac, Rappahannock, York, and James Rivers) and Back Bay. (Transitional or oligohaline waters vary in salinity from 0.5 ppt. to 5.0 ppt.) Fixed transition zones within the Chesapeake Bay drainage correspond with pre-established Chesapeake Bay Program assessment segments. However, at any specific estuarine site the salinity, the sediment chemistry, and the resultant toxicity of contaminants vary temporally. As a consequence, the bottom salinity observed at the time of sampling is used to define habitat classes for benthic IBI evaluations and for the selection of PEC *vs.* ER-M screening values for WOE assessment. An exceedance of these screening values raises a red flag of warning, but does not in itself result in an “Impaired” assessment (Virginia Assessment Category 5A).

When the appropriate SVs are exceeded for one or more contaminants or when toxicity thresholds are exceeded, and no benthic community data are available to corroborate significant benthic degradation, the site is considered to have insufficient information to assess the aquatic life use (Virginia Assessment Category 3E). In such cases, additional biological monitoring should be scheduled to assess actual aquatic life use support. In practice, for WOE assessment, each SV is evaluated based upon its Sediment Quality Guideline Quotient (SQGQ, sometimes abbreviated as “Q”), which is calculated as the ratio between the observed concentration in the sediment and the screening value:  $SQGQ = \text{observed concentration} / \text{SV}$ . A ratio of 1.0 or greater indicates that the screening value was exceeded. A ratio of 2.0 indicates that the observed concentration was twice the screening value, etc. In the WOE assessment, the magnitude of each exceedance, abbreviated as “Q”, is considered and weighted in scoring the degree of chemical contamination. A summary of the SQT Matrix



Scoring Guidelines for sediment contamination can be found at the top of Tab “(1) - SedChem Mean ERMq Model” of the WOE Assessment Workbook.

The use of such screening values for assessment suffers several limitations. First, although they are available for most of the trace toxic metals, they are only available for a very limited number of organic contaminants. Secondly, each screening value reflects the potential effects of a single contaminant and does not consider possible interactions with other contaminants in the same sediment matrix. The often significant effects of additivity, antagonism, and synergism are not considered. A number of efforts have been made in recent years to integrate SQG quotients across multiple contaminants (see Long et al., 2006 for a critical review). The most successful and commonly applied integrated measure is the mean SQG quotient (mSQGQ). In a study of southeastern estuaries Hyland et al. (1999), applying the methods of Long et al. (1998), demonstrated that sites with mean SQG quotients as low as 0.1 had relatively high probabilities of significant degradation of their benthic communities. Applying the mean ERM quotient (mERM-Q) of eight trace metals (excluding Ni), 13 PAHs (excluding Low Molecular Wt. PAHs, High Molecular Wt. PAHs, and total PAHs), total PCBs, plus total DDT, they found that when the mean ER-M quotient exceeded 0.1, the probability of adverse effects on the benthic community was  $\geq 0.75$ . Similar results were observed when using mean quotients for another set of freshwater sediment quality guidelines, the Probable Effects Level (PEL). Experience with WOE assessments over the past 18 years has shown that mean quotients calculated with Virginia’s freshwater screening values (PECs) and the PELs studied by Hyland et al. (2003) correspond very closely, and that the benthic risk matrix provided for PEL quotients in Hyland’s study (ibid.) is appropriate for use with PECs. Both PEC and PEL screening values are provided on Tab “(1) - SedChem Mean ERMq Model” of the WOE Assessment Workbook, and mean quotients are calculated for both (columns L and S). Hyland’s “benthic risk” matrix for tidal freshwater (ibid.) is provided at the bottom of each column. For the purpose of WOE assessment, therefore, when the mean ERM quotient for the specified saltwater contaminants exceeds 0.022, or the PEL/PEC quotient for the specified freshwater contaminants exceeds 0.040, a positive chemical score is reported, whether an individual screening value is exceeded or not (see Table 2, below).

**Table 2 – Matrix of Benthic Impact Risk for Mean SQG Quotient Ranges in the Virginian Biogeographic Province (adapted from Hyland et al., 2003)**

Risk of benthic impact	Mean ERM-Q	Mean PEC-Q Tidal Freshwater	Site Score Mean SQG Quotient
<b>Low</b>	$\leq 0.022$	$\leq 0.040$	0
<b>Medium</b>	$> 0.022 - 0.098$	$> 0.040 - 0.179$	1
<b>High</b>	$> 0.098 - 0.473$	$> 0.179 - 0.969$	2
<b>Very High</b>	$> 0.473$	$> 0.969$	3

**Sediment Chemistry: Workbook Tab “(2) SedChem Logistic Reg Model”** (Logistic Regression Model – page 5)

EPA recently applied a new method for evaluating sediment chemistry in the National Coastal Condition Assessment 2010 (US EPA, 2015). In that Report (also described as NCCA Report V), EPA introduced the Logistic Regression Model (LRM - Field et al., 1999; Field et al., 2002) and the concept of utilizing the maximum probability ( $P_{max}$ ) of acute toxic effects among the sediment contaminant analytes to characterize the ecological condition of a site. To briefly summarize the procedure, a table of LRM coefficients ( $B_0$  = Intercept,  $B_1$  = Slope), and LRM 25th and 75th percentiles was presented in the 2010 Technical Report (US EPA, 2016) for ten individual metals, 21 individual PAHs plus biphenyl, total PCBs, and four pesticides/pesticide derivatives found in estuarine and marine sediments. The observed sediment concentration of each of these

analytes (x) was used to calculate a logistic regression value (LRM<sub>x</sub>) corresponding to the observed concentration of the analyte. Based on the array of all LRM<sub>x</sub> values calculated for the measured analytes in a sample, the maximum LRM<sub>x</sub> value was identified and the probability was calculated of observing significant toxicity based on the observed concentration of that single analyte. Significant toxicity was defined as control-corrected survivorship < 80% and a statistically significant difference from negative controls. This probability was termed P<sub>max</sub>. EPA subsequently utilized the P<sub>max</sub> value in conjunction with the mERM-Q to characterize individual sites in relation to their ecological condition. In order to be classified as “Good”, a site was required to have a P<sub>max</sub> ≤ 0.5 and a mERM-Q < 0.1 (refer to Table S-6, reproduced below). A classification of “Fair” resulted from a P<sub>max</sub> > 0.5 and < 0.75 or a mERM-Q ≥ 0.1 and ≤ 0.5, and a classification of “Poor” resulted if either P<sub>max</sub> was ≥ 0.75 or the mERM-Q was > 0.5.

**Table S-6. Thresholds for sediment chemistry used in NCCA 2010.\***

Ecological Condition by Site		
Rank	Estuarine	Great Lakes
Good	mERM-Q < 0.1 and LRM P <sub>max</sub> ≤ 0.5	mPEC-Q ≤ 0.1
Fair	mERM-Q ≥ 0.1 - ≤ 0.5 or LRM P <sub>max</sub> > 0.5 - < 0.75	mPEC-Q > 0.1 - ≤ 0.6
Poor	mERM-Q > 0.5 or LRM P <sub>max</sub> ≥ 0.75	mPEC-Q > 0.6

\* Reproduced from the 2015 Technical Report (US EPA, 2016) of the National Coastal Condition Assessment 2010 (US EPA, 2015).

The use of this Logistic Regression Model characterization was added to the Weigh-of-Evidence assessments as an additional line of evidence beginning with the 2016 Integrated Report. Observations and comments from Tab “(2) SedChem Logistic Reg Model” are copied to the SQT Evaluation Matrix where they contribute to the final weighting of the SQT Matrix score given for sediment chemistry from Tab (1).

A secondary output statistic from the Logistic Regression Model has been added to the weight-of-evidence workbooks. The probability of toxic effects for each analyte in the model has been calculated in column “P” of the “(2) SedChem Logistic Reg Model” tab. The mean (arithmetic average) probability (P<sub>avg</sub>) of toxic effects across all analytes was then calculated. This statistic integrates potential effects across all contaminant analytes, as opposed to considering only the potential effect of the single most critical contaminant (P<sub>max</sub>). Evaluation of the P<sub>avg</sub> statistic among 442 probabilistic sites sampled within Virginia’s estuarine water between 2005 and 2014 revealed that P<sub>avg</sub> values as low as 0.1000 indicated an elevated probability of significant benthic effects, as expressed by the EMAP IEC for the Virginia biogeographic Province (Smith, 2016 - unpublished results).

Ratings for the LRM P<sub>avg</sub> statistic used in the WOE Workbooks are based on percentile values of sites characterized as having degraded benthic communities by the EMAP IEC:

- |  |                       |
|--|-----------------------|
| P <sub>avg</sub> < 0.0298 (10th percentile) - “Good”             | Matrix Score “0”,     |
| P <sub>avg</sub> ≥ 0.0298 to < 0.0761 (50th percentile) - “Fair” | Matrix Score “1”,     |
| P <sub>avg</sub> ≥ 0.0760 and ≤ 0.1423 (90th percentile) - Poor” | Matrix Score “2”, and |
| P <sub>avg</sub> > 0.1423 - “Very Poor”                          | Matrix Score “3”.     |

The final Matrix Score for the Logistic Regression Model is calculated as the arithmetic average of the Matrix scores for P<sub>max</sub> and P<sub>avg</sub> (cell L1 of the “(2) SedChem Logistic Reg Model” tab).

**Sediment Chemistry: Workbook Tab “(3) SedChem ESB Model”** (Equilibrium Partitioning Sediment Benchmark for PAH Mixtures – page 6)

The concentration of dissolved contaminants in the interstitial water of sediment may also stress benthic infauna. The interstitial water in sediment is difficult to collect and analyze accurately and this is not commonly carried out during normal monitoring programs. However, the concentrations of dissolved contaminants in interstitial water can be estimated from the concentrations in the sediment itself using their **equilibrium partitioning coefficients** and their integrated effects can be predicted by applying procedures similar to those applied for integrating sediment quality guidelines. EPA has published procedures for the derivation of equilibrium partitioning sediment benchmarks (ESBs) for the protection of benthic organisms from several classes of contaminants (US EPA - 2001, 2003a, 2003b, 2003c, 2005, 2008). The guidance manual “Procedures for the Derivation of Equilibrium Partitioning Sediment Benchmarks (ESBs) for the Protection of Benthic Organisms: PAH Mixtures” (US EPA, 2003a) provides orientation for calculating an integrated ESB for a suite of 34 common PAHs that have been included in various intensive sediment studies, as well as conversion factors to be applied for smaller subsets of 23 and 13 PAHs analyzed in other studies. The suite of PAHs analyzed in DEQ’s Estuarine ProbMon Program includes all 23 PAHs of the 23-analyte subset, for which conversion factors have been provided. Tab “(3) SedChem ESB Model - PAHs” performs the necessary calculations and conversion based on the concentrations of PAHs and total organic carbon (TOC) measured in the sediment. When the converted sum of the 23 individual benchmarks reaches or exceeds 1.0, there is a high probability of adverse chronic effects due to the toxicity of dissolved PAHs. Observations and comments from Tab (3) are copied to the SQT Evaluation Matrix where they contribute to the final weighting of the SQT Matrix score given for sediment chemistry from Tab (1).

Even a moderate amount of total organic carbon (TOC) in the sediment is sufficient to sequester PAHs and inhibit their solubility in interstitial water. Consequently, the results from this line of evidence complement other lines of chemical characterization and are useful for interpreting probable causes of impairment, but they are not adequate in themselves to assess a site as impaired or not.

An additional tool for the identification of potential sources of PAHs is the ratio between the members of each of two pairs of compounds (Neff et al., 2005). Depending upon the value of the ratio (see **Table 3**, below), the source may be identified as probably pyrogenic as opposed to petrogenic in origin. Petrogenic PAHs are found in nature, usually at low concentrations; they may be associated with petroleum spills. Pyrogenic PAHs are combustion byproducts, and usually result from the combustion of petroleum products (*e.g.*, emissions from fuel consumption by outboard motors, etc.).

**Table 3 – The Identification of Pyrogenic vs. Petrogenic Sources of PAH Contaminants based on the ratio of concentrations of Phenanthrene / Anthracene and Fluoranthene / Pyrene (Neff et al., 2005)**

<u>Ratio</u>	<u>Value</u>	<u>Probable Source of PAHs</u>
<b>Phenanthrene / Anthracene Ratio</b>	<b>If &lt;7.0</b>	<b>Probably Pyrogenic</b>
	<b>If &gt;10.0</b>	<b>Probably Petrogenic</b>
<b>Fluoranthene / Pyrene Ratio</b>	<b>If &lt;0.9</b>	<b>Possibly Petrogenic</b>
	<b>If &gt;1.0</b>	<b>Possibly Pyrogenic</b>

The Fluoranthene / Pyrene Ratio is much more variable among pyrogenic and petrogenic sources of PAHs than is the Phenanthrene / Anthracene Ratio. Consequently the resultant classification is considered a possible rather than a highly probable source.

## **Sediment Total Organic Carbon (TOC):** (Summary Sheet – page 2)

The concentration of total organic carbon in the sediment influences the availability and route of uptake of toxic contaminants by benthic organisms. Organic carbon absorbs or sequesters many organic and inorganic contaminants, and many benthic infauna organisms actively feed on the organic detritus where these contaminants concentrate. Elevated amounts of TOC are consequently considered undesirable for benthic infauna that ingest sediment particulates. Sediment quality indices published in a series of National Coastal Condition Reports (US EPA, 2001, 2005, 2008) classify sediments with more than 5% TOC as being of poor quality. More recently, Hyland et al. (2005) reported that sediment TOC concentration as low as 3.5% may induce a high risk to benthic communities. Sediment TOC evaluations are presented under both sets of guidelines included in the Summary Sheet of page 2 - NCCA guidelines in line 18 and Hyland et al. guidelines in line 36.

Conversely, because of the equilibrium partitioning of contaminants between sediment carbon and interstitial water, high TOC concentrations in the sediment tend to lower their ESBs and reduce the risk from dissolved toxics that would diffuse across gills and other semi-permeable membranes. Higher TOC concentrations would be beneficial in reducing toxic effects through this route of uptake. Sediment TOC concentration is consequently maintained as an ancillary line of evidence for the interpretation of sediment contamination and is used in the calculation of ESBs on Tab “(3) SedChem ESB Model”.

The final Matrix Score for sediment chemistry appears in cell C19 of the “SQT Evaluation Matrix” tab. It consists of the maximum score among the three lines of chemical evidence: SedChem Mean ERMq Model – Tab (1), SedChem Logistic Reg Model – Tab (2), and SedChem ESB Model – PAHs – Tab (3).

## **Sediment Toxicity: Workbook Tab “(4) Sediment Toxicity” (Page 7)**

The magnitude of effects observed during sediment toxicity tests can be applied for weighting this line of evidence. The survival of test organisms, expressed by the percent control-corrected survival or control-corrected mortality endpoint, is generally associated with the acute effects of higher levels of toxicants (although chemical additivity, antagonism and synergism among contaminants can also play a role). Sub-lethal test endpoints that provide a measure of chronic exposure effects at an increased level of sensitivity, with lower toxicant concentrations, include organism growth rate (expressed as weight), reburial rate (amphipods), reproductive rate, etc. In relative terms, the ecological significance of these endpoints is not likely to be as critical as the measure of survival. Therefore, less weight is applied in cases where only these endpoints show effects. In situations where the survival endpoint yields statistically significant effects by one or more species, greater weight would be applied accordingly. It is important to factor the number of test species, their taxonomic identifications and the associated sensitivities of each test species into the weighting. During the tests, attention must also be applied to artificial toxicity such as in the case of naturally occurring sediment ammonia. Another complication is salinity adjustment, which can chemically alter the sediment, thus leading to changes in chemical bioavailability and ultimately affecting sediment toxicity (Roberts et al., 2002). In the field, indigenous predators can also significantly alter the test outcome. Most of these factors are considered and controlled during the performance of toxicity tests in the laboratory.

Toxicity tests performed in compliance with the QAPP of the National Coastal Assessment Program (US EPA, 2009b) and continued within the DEQ Estuarine Probabilistic Monitoring Program have generally been limited to a single test format with a single test species: ten-day static acute toxicity tests with an estuarine amphipod, conducted in accordance with standard ASTM guidelines and EPA methods. (Prior to 2010, *Ampelisca abdita* was the amphipod of choice. Beginning in 2010, the NCCA Program, as well as DEQ’s Estuarine Probabilistic Monitoring Program, switched from *Ampelisca abdita* to the amphipod *Leptocheirus plumulosus* as a test

organism.) Beginning in 2012 DEQ started to observe exceptionally high mortalities in toxicity tests of sediment from a number of tidal freshwater sites. Apparently, adjusting the salinities from freshwater (<0.5‰) to a standardized estuarine salinity of 20‰ following estuarine toxicity test protocols, changed sediment chemistry sufficiently to stimulate the proliferation of iron-fixing bacteria, the subsequent precipitation of a reddish-brown flocculant, a precipitous drop in pH and increased ammonia concentrations, resulting in catastrophic amphipod mortality. Consequently, beginning in 2013 we included a parallel freshwater toxicity test using *Hyalella azteca* as a test organism. Subsequently, we reported the results of both tests for sediment from tidal freshwater sites. Often, control-corrected survivorship was comparable and high in both tests, but when poor results from the estuarine test were accompanied by the manifestations described above, sediment toxicity was characterized primarily by the results of the freshwater test.

In either case, the specified end-point was amphipod survival. The results of these tests are provided on the “SedTox Data” Tab (page 11) of the WOE Assessment Workbook and the final evaluation and scoring are carried out on Tab “(4) Sediment Toxicity.” Both statistical significance and ecological significance of the results are considered. The statistical significance of test results is tested at a significance level of  $\alpha = 0.05$  ( $\geq 95\%$  confidence that differences from control are real); ecological significance is assumed only if control-corrected survivorship is  $< 80\%$ . On rare occasion, results may be statistically significant but not ecologically significant, or *vice versa*. In such cases a score of 1 (marginal toxicity = “Fair”) is awarded for evaluation in the SQT matrix and the Summary Sheet.

### **Ancillary Parameter Values and the Identification of Additional Potential Stressors**

**Near Bottom Salinity (‰) and Percent Sand (%):** The near-bottom salinity class observed at the time of sample collection, and the percent sand in the sediment sample are used in several ancillary evaluations. On the “Summary Sheet” Tab of the WOE Assessment Workbook these two measurements are automatically integrated into a “Habitat Type” characterization for application of the CBP B-IBI and the Middle Atlantic B-IBI on Tab “(5) Benthic Fauna”. This integration is also performed by the BEL benthic database at Old Dominion University during the calculation of the CBP and MAIA Benthic IBIs.

In addition, the relative proportions of sand vs. fines (silt/clay) in the sediment (“sand” vs. “mud” substrate) can be used to differentiate high energy from low energy benthic environments. Percent sand  $\geq 98.00\%$  and TOC concentrations  $\leq 0.5\%$  are generally indicative of current-scoured substrate or dynamic habitat (due to wave action). Chemical contamination is always very low and benthic communities are almost always (naturally) degraded at such sites. Both are areas where the substrate is at least periodically in movement and fine particulates tend to be washed away. Filter feeders may predominate in such areas, while deposit feeders may predominate in low energy areas where fine particulates accumulate. Substrate type may also serve as an indication of the relative risk of chemical contamination. Contaminants are more readily absorbed, transported and deposited by fine particles (silt/clay) and associated organic detritus than by sand. (Incidentally, sediment with extremely high sand content often stimulates amphipod mortality that is not associated with chemical contamination during toxicity tests. Some amphipod species respond negatively to high sand concentrations (U.S EPA, 1996; Emery et al., 1997).

**Near Bottom Dissolved Oxygen (DO - mg/L) and Depth:** Low dissolved oxygen presents a direct stress on benthic fauna. This may result from natural thermal and/or salinity stratification that inhibit mixing in deep channels, from nutrient enrichment and eutrophication, or from a combination of both. In warmer, shallower waters low DO concentrations (mg/L) may result simply from oxygen’s lower solubility at higher temperatures. In either case, the observation of a single low near-bottom DO concentration at a probabilistic site is not sufficient to result in an impaired assessment. It is evaluated only as one among various potential causes for any benthic degradation that is observed.

**Bottom Temperature** (°C): Higher water temperature may itself be a significant stressor, in addition to its indirect effect via DO depression.

**Total Organic Carbon (TOC)**: As indicated above, the concentration of Total Organic Carbon in the sediment can influence the degree of exposure to and the uptake route of chemical contaminants by benthic organisms. In association with other water quality characteristics, it may also provide insight into the degree of eutrophication present in the estuarine system.

While potential scenarios based on these integrated lines of evidence are too numerous to list, for many sites the conclusion should be obvious. For example, chemical contaminants are commonly detected but at concentrations below their respective SQGs. If the biological results from the same samples indicate a lack of effects, as demonstrated by lack of significant sediment toxicity and the presence of a healthy benthic community, the resultant listing would be VA Category 2A (fully supporting designated use). On the opposite end of the spectrum, all three components of the triad may show extreme effects and the site would consequently be listed in the 5A category (impaired for toxics – potentially needing a TMDL).

**Ancillary Indices**: Beginning in 2011, several indices used for site evaluation by the National Coastal Condition Assessment (NCCA) Reports were added to the Summary sheets of WOE Assessment workbooks. A Sediment Quality Index (SQI) integrates the results of sediment toxicity, sediment contamination, and sediment TOC content into a rating of “Good”, “Fair” or “Poor.” A Water Quality Index (WQI) integrates near surface dissolved inorganic nutrients (nitrogen and phosphorus), chlorophyll-a, water clarity, and near-bottom dissolved oxygen into a similar rating scale. In 2013, a national workgroup began reevaluating the thresholds for classification of the individual components of these indices for the fifth National Coastal Condition Report (U.S. EPA, 2015). Once a final consensus was reached for the national report, the thresholds in the WOE workbooks were adjusted accordingly (see Table 2-5, below). A résumé of the three types of benthic evaluations (CBP B-IBI, Mid Atlantic B-IBI, EMAP VP-IEC) applied in the WOE Assessment has also been included on the Summary tab to facilitate site characterizations.

**Table 2-5. NCCA guidelines for evaluating the five component indicators used in the water quality index to assess estuarine coastal condition.\***

Estuarine Water Quality Thresholds				
	Region	Good	Fair	Poor
Surface Concentrations of Dissolved Inorganic Nitrogen (DIN): Estuaries	Northeast Southeast Gulf	< 0.1 mg/L	0.1 – 0.5 mg/L	> 0.5 mg/L
	West	< 0.35 mg/L	0.35 – 0.5 mg/L	> 0.5 mg/L
	Tropical <sup>a</sup>	< 0.05 mg/L	0.05 – 0.1 mg/L	> 0.1 mg/L
Surface Concentrations of Dissolved Inorganic Phosphorus (DIP): Estuaries	Northeast Southeast Gulf	< 0.01 mg/L	0.01 – 0.05 mg/L	> 0.05 mg/L
	West	< 0.07 mg/L	0.07 – 0.1 mg/L	> 0.1 mg/L
	Tropical <sup>a</sup>	< 0.005 mg/L	0.005 – 0.01 mg/L	> 0.01 mg/L
Surface Concentrations of Chlorophyll <i>a</i> : Estuaries	Northeast Southeast Gulf	< 5 µg/L	5 – 20 µg/L	> 20 µg/L
	West			
	Tropical <sup>a</sup>	< 0.5 µg/L	0.5 – 1 µg/L	> 1 µg/L
Water Clarity (percent of incident light remaining after passing through 1 meter of water): Estuaries	Waters with naturally high turbidity	> 10%	5 – 10%	< 5%
	Waters with normal turbidity	> 20%	10 – 20%	< 10%
	Waters that support SAV <sup>b</sup>	> 40%	20 – 40%	< 20%
Bottom Water Concentrations of Dissolved Oxygen: Estuaries	All	> 5 mg/L	2 – 5 mg/L	< 2 mg/L

<sup>a</sup> Tropical refers to NCCA Florida Bay sites. <sup>b</sup> Submerged Aquatic Vegetation.

\* Reproduced from the 2015 Technical Report (US EPA, 2016) of the National Coastal Condition Assessment 2010 (US EPA, 2015).

It is preferable that this WOE approach only be applied when all three sediment data components are available from a particular site. It is still possible, however, to implement this process if only two elements are available, as long as data on the condition of the resident benthic community is included (e.g., sediment chemistry and benthic IBI or sediment toxicity and benthic IBI). If both lines of evidence are in agreement as to the condition of the site (e.g., degraded or severely degraded), a corresponding assessment may be attained (i.e., Category 5A with toxics as a potential cause). If such a conclusion is suggested based solely on sediment chemistry and sediment toxicity data, follow-up monitoring should be scheduled (Virginia Assessment Category 3B), even if both chemical and toxicological results are in agreement on the potential existence of a toxic condition. For those instances where the conclusions are not obvious, it will be necessary to obtain consensual agreement between Central Office and the Regional Office responsible for the assessment of that water body. If agreement cannot be attained, advice should be sought from DEQ’s Academic Advisory Committee.

## References

- Buchanan, M.F. 1999. National Oceanic and Atmospheric Administration Screening Quick Reference Tables, NOAA HAZMAT Report 991 Seattle, WA, Hazardous Materials Response and Assessment Division, 12 pages.
- Borja, A., J. Franco, V. Pérez. 2000. A marine biotic index to establish the ecological quality of soft-bottom benthos within European estuarine and coastal environments.

- Mar. Pollut. Bull. 40, 1100–1114.
- Chapman, P.M. 1986. Sediment quality criteria from the sediment quality triad – an example. *Environ. Toxicol. Chem.* 5: 957-964.
- Chapman, P.M., R.N. Dexter, S.F. Cross and D.G. Mitchell. 1986. A field trial of the Sediment Quality Triad in San Francisco Bay. NOAA Technical Memorandum NOS OMA 25. National Oceanic and Atmospheric Administration, San Francisco, CA. 127 pp.
- Chapman, P.M., R.N. Dexter and E.R. Long. 1987. Synoptic measures of sediment contamination toxicity and infaunal community structure (the Sediment Quality Triad). *Mar Ecol. Prog. Ser.* 37:75-96.
- Chapman, P.M., 1992. Sediment Quality Triad Approach. In: *Sediment Classification Methods Compendium*. EPA 823-R-92-006 Ch. 10 pp. 1-18.
- Chapman, P.M, B. Anderson, S. Carr, V. Engle, R. Green, J. Hameedi, M. Harmon, P. Haverland, J. Hyland, C. Ingersoll, E. Long, J. Rodgers Jr, M. Salazar, P. Sibley, P. Smith, R. Swartz, B. Thompson and H. Windom. 1997. General guidelines for using the sediment quality triad. *Mar Pollut Bull* 34(6):368-372.
- Dauer, D.M., and M.F. Lane. 2005. Side-by-Side Comparison of Young Grab and Composite Petite Ponar Grab Samples for the Calculation of the Benthic Index of Biological Integrity (B-IBI). Report to: Chesapeake Bay Program Office, Virginia Department of Environmental Quality, 629 East Main Street, Richmond, Virginia 23230. June, 2005. 46pp.
- Dauer, D.M., M.F. Lane and R.J. Llansó. 2002. Development of Diagnostic Approaches to Determine Sources of Anthropogenic Stress Affecting Benthic Community Condition in the Chesapeake Bay. Final Report to the U.S. Environmental Protection Agency, Chesapeake Bay Program Office, Annapolis, Maryland, 64 pp.
- Emery, V.L., Jr., D.W. Moore, B.R. Gray, B.M. Duke, A.B. Gibson, R.B. Wright, and J.D. Farrar. 1997. Development of a chronic sublethal sediment bioassay using the estuarine amphipod *Leptocheirus plumulosus* (Shoemaker). *Environ. Toxicol. Chem.* 16(9):1912-1920.
- Hall, L.W. and R.W. Alden III. A review of concurrent ambient water column and sediment toxicity testing in the Chesapeake Bay watershed: 1990-1994. *Environ. Toxicol. Chem.* 16:1607-1617.
- Hall, L.W. Jr., and R.D. Anderson. 1997. A Pilot Study for Ambient Toxicity Testing in Chesapeake Bay Year 4 Report. Chesapeake Bay Program, 410 Severn Ave., Suite 109, Annapolis, MD 21403.
- Hyland, J.L., R.F. Van Dolah, and T.R. Snoots. 1999. Predicting stress in benthic communities of Southeastern U.S. estuaries in relation to chemical contamination of sediment. *Environmental Toxicology and Chemistry*, 18(11): 2557-2564.
- Hyland, J, L. Balthis, I. Karakassis, P. Magni, A. Petrov, J. Shine, O. Vestergaard, R. Warwick. 2005. Organic carbon content of sediments as an indicator of stress in the marine benthos. *Marine Ecology Progress Series*. Vol. 295: 91–103, 2005 Published June 23
- Llansó, R.J., and D.M. Dauer. 2003. Methods for Calculating the Chesapeake Bay Benthic Index of Biotic Integrity. Chesapeake Bay Program. 24 pp. [www.baybenthos.versar.com](http://www.baybenthos.versar.com)
- Llansó, R.J., L.C. Scott, J.L. Hyland, D.M. Dauer, D.E. Russell, and F.W. Kutz. 2002. An Estuarine Benthic Index of Biotic Integrity for the Mid-Atlantic Region of the United States. II. Index Development. *Estuaries* Vol. 25, No. 6A, p. 1231–1242 December 2002.
- Long, E.R., and P.M. Chapman. 1985. A sediment quality triad: Measures of sediment contamination, toxicity and infaunal community composition in Puget Sound. *Mar. Pollu. Bull.* 16(10): 405-415.
- Long, E.R., D.D. MacDonald, S.L. Smith, F.D. Calder. 1995. Incidence of adverse biological effects within ranges of chemical concentrations in marine and estuarine sediments. *Environmental Management* vol 19(1):81-97.
- Long, E.R., D.D. MacDonald, C.G. Severn, and C.B. Hong. 2000. Classifying the probabilities of acute toxicity in marine sediments with empirically-derived sediment quality guidelines. *Environmental Toxicology & Chemistry* 19(10): 2598-2601.
- Long, E.R., C.G. Ingersoll, and D.D. MacDonald. 2006. Calculation and uses of mean sediment quality guideline quotients: A critical review. *Environmental Science and Technology* (March 2006 issue).



- Long, E.R., J. Field, and D. MacDonald. 1998. Predicting toxicity in marine sediments with numerical sediment quality guidelines. *Environmental Toxicology and Chemistry* 17:714-727
- MacDonald, D.D., C.G. Ingersoll, T.A. Berger. 2000. Development and Evaluation of Consensus Based Sediment Quality Guidelines for Freshwater Ecosystems. *Arch. Environ. Contam. Toxicol.* 39:2031.
- McGee, B.L., D.J. Fisher, J. Ashley and D. Velinsky. 2001. Using the Sediment Quality Triad to characterize toxic conditions in the Chesapeake Bay (1999): An assessment of tidal river segments in the Bohemia, Magothy, Patuxent, James, and York Rivers. EPA 903-R-01-008, CBP/TRS 257/01, Environmental Protection Agency, Chesapeake Bay Program Office, Annapolis, MD, 35 pp + Appendices.
- Neff, J.M., S.A. Stout, and D.G. Gunsert. 2005. Ecological Risk Assessment of Polycyclic Aromatic Hydrocarbons in Sediments: Identifying Sources and Ecological Hazard. *Integr Environ Assess Manag* 1(1): 22-33. © 2005 SETAC.
- Paul, J.F., K.J. Scott, D.E. Campbell, J.H. Gentile, C.S. Strobel, R.M. Valente, S.B. Weisberg, A.F. Holland and J.A. Ranasinghe. 2001. Developing and applying a benthic index of estuarine condition for the Virginian Biogeographic Province. *Ecological Indicators* 1: 83-99.
- Pelletier, M.C., D.J. Gillett, A. Hamilton, T. Grayson, V. Hansen, E.W. Leppo, S.B. Weisberg, A. Borja. 2018. Adaptation and application of multivariate AMBI (M-AMBI) in US coastal waters. *Ecological Indicators*, Volume 89, June 2018, Pages 818-827
- Roberts, M.H. Jr, M.A. Vogelbein, M.A. Richards, L. Seivard, and P.F. De Lisle. 2002a. Chemical and Toxicological Characterization of Tidal Freshwater Areas in the James River, Virginia. EPA 903-R-02-006, CBP/TRS 264/02, Environmental Protection Agency, Chesapeake Bay Program Office, Annapolis, MD, 123 pp + Appendices.
- Roberts, M.H. Jr, M.A. Vogelbein, M.A. Richards, L. Seivard, and P.F. De Lisle. 2002b. Chemical and Toxicological Characterization of Tidal Freshwater Areas in the James River, Virginia Between Jordan Point and Richmond. Final report to VA Department of Environmental Quality. 49 pp + Appendices.
- Roberts, M.H. Jr, M.A. Richards, P.F. DeLisle. 2003. Chemical and Toxicological Characterization of the Lower Mobjack Bay, York River, Virginia Segment of the Chesapeake Bay. Final Report to VA Department of Environmental Quality. 46 pp + Appendices.
- Smith, D.H. 2016. Comparative Evaluation of Mean ERM-Q (mERM-Q) and the Logistic Regression Model Pmax vs Pavg Output Statistics for Sediment Contamination Characterizations relative to Benthic Community Integrity. Unpublished report prepared for the National Coastal Condition Assessment Steering Committee. March 2016.
- U.S. EPA. 1996. Ecological Effects Test Guidelines: OPPTS 850:1740: Whole Sediment Acute Toxicity Invertebrates, Marine. Technical Report EPA 712 C-96-355. U.S. Environmental Protection Agency, Washington, DC.
- U.S. EPA. 1999. *Targeting Toxics: A Characterization Report. A Tool for Directing Management and Monitoring Actions in the Chesapeake Bay's Tidal Rivers*. CPB/TRS 222/103, US EPA, Chesapeake Bay Program, Annapolis, MD.
- U.S. EPA. 2001. Environmental Monitoring and Assessment Program (EMAP): National Coastal Assessment Quality Assurance Project Plan 2001-2004. United States Environmental Protection Agency, Office of Research and Development, National Health and Environmental Effects Research Laboratory, Gulf Ecology Division, Gulf Breeze, FL. EPA/620/R-01/002.
- U.S. EPA. 2001a. National Coastal Condition Report. EPA/620-R-01-005. U.S. Environmental Protection Agency, Office of Research and Development and Office of Water, Washington, DC.  
<http://water.epa.gov/type/oceb/assessmonitor/nccr/index.cfm>
- U.S. EPA. 2001b. Supporting documentation used in the derivation of selected freshwater Tier 2 ESBs. Office of Research and Development, Atlantic Ecology Division, Narragansett, RI USA. Contribution AED-08-044 of the Office of Research and Development National Health and Environmental Effects Research Laboratory's Atlantic Ecology Division.  
[http://www.epa.gov/nheerl/publications/files/Appendix\\_A\\_Cupdate7\\_2.pdf](http://www.epa.gov/nheerl/publications/files/Appendix_A_Cupdate7_2.pdf)

- U.S. EPA. 2003a. Procedures for the Derivation of Equilibrium Partitioning Sediment Benchmarks (ESBs) for the protection of Benthic Organisms: **PAH Mixtures**. United States Environmental Protection Agency, Office of Research and Development, Washington, DC. EPA-600-R-02-013. November 2003. <http://www.epa.gov/nheerl/publications/files/PAHESB.pdf>
- U.S. EPA. 2003b. Procedures for the Derivation of Equilibrium Partitioning Sediment Benchmarks (ESBs) for the Protection of Benthic Organisms: **Endrin**. EPA-600-R-02-009. Office of Research and Development. Washington, DC 20460. <http://www.epa.gov/nheerl/publications/files/endrin.pdf>
- U.S. EPA. 2003c. Procedures for the Derivation of Equilibrium Partitioning Sediment Benchmarks (ESBs) for the Protection of Benthic Organisms: **Dieldrin**. EPA-600-R-02-010. Office of Research and Development. Washington, DC 20460. <http://www.epa.gov/nheerl/publications/files/dieldrin.pdf>
- U.S. EPA (Environmental Protection Agency). 2005a. National Coastal Condition Report II. EPA-620/R-03/002. U.S. Environmental Protection Agency, Office of Research and Development and Office of Water, Washington, DC. (December 2004) <http://water.epa.gov/type/oceb/assessmonitor/nccr/index.cfm>
- U.S. EPA. 2005b. Procedures for the Derivation of Equilibrium Partitioning Sediment Benchmarks (ESBs) for the Protection of Benthic Organisms: **Metal Mixtures** (Cadmium, Copper, Lead, Nickel, Silver, and Zinc). United States Environmental Protection Agency, Office of Research and Development, Washington, DC. EPA/600/R-02/011. January 2005. [http://epa.gov/nheerl/publications/files/metalsESB\\_022405.pdf](http://epa.gov/nheerl/publications/files/metalsESB_022405.pdf)
- U.S. EPA. 2007. Sediment Toxicity Identification Evaluation (TIE) Phases I, II, and III Guidance Document EPA/600/R-07/080 Office of Research and Development. Washington, DC <http://www.epa.gov/nheerl/publications/files/Sediment%20TIE%20Guidance%20Document.pdf>
- U.S. EPA. 2008. Procedures for the Derivation of Equilibrium Partitioning Sediment Benchmarks (ESBs) for the Protection of Benthic Organisms: Compendium of Tier 2 Values for Nonionic Organics. EPA-600-R-02-016. Office of Research and Development. Washington, DC 20460 [http://www.epa.gov/nheerl/publications/files/ESB\\_Compndium\\_v14\\_final.pdf](http://www.epa.gov/nheerl/publications/files/ESB_Compndium_v14_final.pdf)
- U.S. EPA (Environmental Protection Agency). 2009a. National Coastal Condition Report III. EPA/842-R-08-002. U.S. Environmental Protection Agency, Office of Research and Development and Office of Water, Washington, DC. (December 2008) <http://water.epa.gov/type/oceb/assessmonitor/nccr/index.cfm>
- U.S. EPA. 2009b. National Coastal Condition Assessment Quality Assurance Project Plan 2008-2012. United States Environmental Protection Agency, Office of Water, Office of Wetlands, Oceans and Watersheds. Washington, D.C. EPA/841-R-09-004
- U.S. EPA. 2009c. National Coastal Condition Assessment: Field Operations Manual. EPA-841-R-09-003. U.S. Environmental Protection Agency, Washington, DC.
- U.S. EPA (United States Environmental Protection Agency). 2012. National Coastal Condition Report IV. EPA-842-R-10-003. Office of Research and Development and Office of Water, Washington, DC. <http://water.epa.gov/type/oceb/assessmonitor/nccr/index.cfm>
- U.S. EPA - US Environmental Protection Agency, US Geological Survey, US Fish and Wildlife Service. 2012. Toxic Contaminants in the Chesapeake Bay and its Watershed: Extent and Severity of Occurrence and Potential Biological Effects, USEPA Chesapeake Bay Program Office, Annapolis, MD, December, 2012, 175 pages.
- U.S. EPA. 2014a. National Coastal Condition Assessment Quality Assurance Project Plan. United States Environmental Protection Agency, Office of Water, Office of Wetlands, Oceans and Watersheds. Washington, D.C. EPA 841-R-14-005. 2014.
- U.S. EPA. 2014b. National Coastal Condition Assessment: Field Operations Manual. EPA-841-R-14-007. U.S. Environmental Protection Agency, Washington, DC.
- U.S. EPA. 2015. National Coastal Condition Assessment 2010. U.S. Environmental Protection Agency. Office of Water and Office of Research and Development. (2015). National Coastal Condition Assessment 2010 (EPA 841-R-15-006). Washington, DC. December 2015.

U.S. EPA. 2016. U.S. Environmental Protection Agency. Office of Water and Office of Research and Development. (2016). NCCA 2010 Technical Report. Washington, DC. January 2016.

(Citation format suggested by Hugh Sullivan, NCCA QA Officer, personal communication 04Feb2016 – will be updated at a later time.)

Weisberg, S.B., J.A., Ranasinghe, DM. Dauer, L.C. Schaffner, R.J. Diaz and J.B. Frithsen. 1997. An estuarine benthic index of biotic integrity (B-IBI) for Chesapeake Bay. *Estuaries* 20: 149-158.

## APPENDIX H

### SIGNIFICANT LAKES/RESERVOIRS BY REGION

#### **Northern Regional Office – 18 Reservoirs/Lakes**

<b>Abel Lake</b>	Stafford Co.	174 (Acres)	PWS
<b>Aquia Reservoir</b> (Smith Lake)	Stafford Co.	131	PWS
<b>Beaverdam Reservoir</b>	Loudoun Co.	301	PWS
<b>Breckenridge Reservoir</b>	Prince William Co.	47	PWS
<b>Burke Lake</b>	Fairfax Co.	208	DWR
<b>Curtis Lake</b>	Stafford Co.	58	
<b>Goose Creek Reservoir</b>	Loudoun Co.	40	PWS
<b>Hunting Run Reservoir</b>	Spotsylvania Co.	440	PWS
<b>Lake Anna</b>	Louisa, Spotsylvania, Orange	9,595	
<b>Lake Manassas</b>	Prince William Co.	675	PWS
<b>Lake Orange</b>	Orange Co.	124	
<b>Lake Pelham</b>	Culpeper Co.	250	PWS
<b>Lunga Reservoir</b>	Prince William Co.	477	PWS
<b>Motts Run Reservoir</b>	Spotsylvania Co.	137	PWS
<b>Mountain Run Lake</b>	Culpeper Co.	73	PWS
<b>Ni Reservoir</b>	Spotsylvania Co.	408	PWS
<b>Northeast Creek Reservoir</b>	Louisa Co.	178	PWS
<b>Ocoquan Reservoir</b>	Fairfax, Prince William Co.	1,333	PWS

#### **Piedmont Regional Office – 22 Reservoirs/Lakes**

<b>Amelia Lake</b>	Amelia Co.	98	DWR
<b>Briery Creek Lake</b>	Prince Edward Co.	825	DWR
<b>Brunswick Lake</b>	Brunswick Co.	138	DWR
<b>Lake Chesdin</b>	Chesterfield Co.	3,164	PWS
<b>Chickahominy Lake</b>	Charles City Co.	1,049	PWS
<b>Diascund Creek Reservoir</b>	New Kent Co.	1,055	PWS
<b>Emporia Lake</b>	Greensville Co.	290	PWS
<b>Falling Creek Reservoir</b>	Chesterfield Co.	88	
<b>Fort Pickett Reservoir</b>	Nottoway Co.	319	
<b>Great Creek Reservoir</b>	Lawrenceville	219	
<b>Harrison Lake</b>	Charles City Co.	60	
<b>Holiday Lake</b>	Appomattox Co.	113	
<b>Lake Nottoway</b>	Nottoway Co.	161	
<b>Lakeview Reservoir</b>	Chesterfield Co.	43	
<b>Little Creek Reservoir</b>	James City Co.	926	PWS
<b>Lunenburg Beach Lake</b>	Town of Victoria	12	PWS
<b>Modest Creek Reservoir</b>	Town of Victoria	20	PWS
<b>Powhatan Lake (U &amp; L)</b>	Powhatan Co.	61	
<b>Sandy River Reservoir</b>	Prince Edward Co.	718	
<b>Swift Creek Lake</b>	Chesterfield Co.	102	
<b>Swift Creek Reservoir</b>	Chesterfield Co.	1,581	PWS
<b>Troublesome Creek Reservoir</b> (SCS Impoundment #2)	Buckingham Co.	53	PWS

### **Southwest Regional Office – 11 Reservoirs**

<b>Bark Camp Lake</b>	Scott Co.	29	USFS
<b>Big Cherry Lake</b>	Wise Co.	103	PWS
<b>Hidden Valley Lake</b>	Russell Co.	58	DWR
<b>Hungry Mother Lake</b>	Smyth Co.	100	DCR
<b>J. W. Flannagan Reservoir</b>	Dickenson Co.	1,177	USACE /PWS
<b>Lake Keokee</b>	Lee Co.	97	DWR
<b>Laurel Bed Lake</b>	Russell Co.	312	DWR
<b>North Fork Pound Reservoir</b>	Wise Co.	116	USACE /PWS
<b>Rural Retreat Lake</b>	Wythe Co.	85	DWR
<b>South Holston Reservoir</b>	Washington Co.	1,699	TVA/PWS
<b>Wise Reservoir</b>	Wise Co.	46	WISE/PWS

### **Tidewater Regional Office – 21 Reservoirs/Lakes**

<b>Airfield Pond</b>	Sussex Co.	120	DWR
<b>Harwood Mills Reservoir</b>	York Co.	258	PWS
<b>Lake Burnt Mills</b>	Isle of Wight Co.	638	PWS
<b>Lake Cohoon</b>	Suffolk City	454	PWS
<b>Lake Drummond</b>	Suffolk City	3,242	
<b>Lake Kilby</b>	Suffolk City	200	PWS
<b>Lake Lawson</b>	Virginia Beach	75	
<b>Lake Meade</b>	Suffolk City	490	PWS
<b>Lake Prince</b>	Suffolk City	709	PWS
<b>Lake Smith</b>	Norfolk City	185	PWS
<b>Lake Whitehurst</b>	Norfolk City	495	PWS
<b>Lake Wright</b>	Norfolk City	12	
<b>Lee Hall Reservoir</b>	Newport News	290	PWS
<b>Little Creek Reservoir</b>	Norfolk City	200	PWS
<b>Lone Star Lake F</b>	Suffolk City	19	PWS
<b>Lone Star Lake G</b>	Suffolk City	90	PWS
<b>Lone Star Lake I</b>	Suffolk City	33	PWS
<b>Speights Run Lake</b>	Suffolk City	118	PWS
<b>Stumpy Lake</b>	Virginia Beach	263	
<b>Waller Mill Reservoir</b>	York Co.	288	PWS
<b>Western Branch Reservoir</b>	Norfolk City	1,205	PWS

### **Valley Regional Office – 21 Reservoirs/Lakes**

<b>Beaver Creek Reservoir</b>	Albemarle Co.	96	PWS
<b>Chris Greene Lake</b>	Albemarle Co.	57	PWS
<b>Coles Run Reservoir</b>	Augusta Co.	11	USFS/PWS
<b>Douthat Lake</b>	Bath Co.	47	DCR
<b>Elkhorn Lake</b>	Augusta Co.	51	USFS/PWS
<b>Fluvanna Ruritan Lake</b>	Fluvanna Co.	51	DWR
<b>Lake Albemarle</b>	Albemarle Co.	37	DWR
<b>Lake Arrowhead</b>	Page Co.	36	DWR
<b>Lake Frederick</b>	Frederick Co.	67	DWR
<b>Lake Nelson</b>	Nelson Co.	41	DWR
<b>Lake Robertson</b>	Rockbridge Co.	24	DWR
<b>Mount Jackson Reservoir</b>	Shenandoah Co.	1	PWS
<b>Ragged Mountain Reservoir</b>	Albemarle Co.	71	PWS
<b>Rivanna Reservoir</b>	Albemarle Co.	399	PWS

Final Guidance for 2024 IR  
Appendix H

Lake Shenandoah	Rockingham Co.	36	DWR
Silver Lake	Rockingham Co.	11	PWS
Staunton Dam Lake	Augusta Co.	21	PWS
Strasburg Reservoir	Shenandoah Co.	5	PWS
Switzer Lake	Rockingham Co.	99	USFS/PWS
Sugar Hollow Reservoir	Albemarle Co.	47	PWS
Totier Creek Reservoir	Albemarle Co.	37	PWS

**Blue Ridge Regional Office - Roanoke – 31 Reservoirs/Lakes**

Beaverdam Creek Reservoir	Bedford Co.	70	PWS
Bedford (Stony Cr.) Reservoir	Bedford Co.	28	PWS
Carvin Cove Reservoir	Botetourt Co.	632	PWS
Cherrystone Reservoir	Pittsylvania Co.	104	PWS
Claytor Lake	Pulaski Co.	4,287	PWS
Clifton Forge (Smith Cr.) Reservoir	Alleghany Co.	10	PWS
Conner Lake	Halifax Co.	98	DWR
Fairystone Lake	Henry Co.	127	
Gatewood Reservoir	Pulaski Co.	176	PWS
Georges Creek Reservoir	Pittsylvania Co.	8	PWS
Graham Creek Reservoir	Amherst Co.	40	PWS
Hogan Lake	Pulaski Co.	36	PWS
Kerr Reservoir (Va.'s portion)	Halifax Co.	33,300	USACE /PWS
Keysville Reservoir	Charlotte Co.	36	PWS
Lake Gordon	Mecklenburg Co.	115	DWR
Lake Gaston (Va.'s portion)	Brunswick Co.	5,614	PWS
Leesville Reservoir	Bedford Co.	2,630	PWS
Little River Reservoir	Montgomery Co.	60	PWS
Martinsville Reservoir	Henry Co.	181	PWS
Mill Creek Reservoir	Amherst Co.	190	
Lake Moomaw	Bath Co.	2,389	USACE
Pedlar Lake	Amherst Co.	118	PWS
Phelps Creek Reservoir	Campbell Co.	19	PWS
Philpott Reservoir	Henry Co.	2,813	USACE
Roaring Fork Reservoir	Pittsylvania Co.	19	PWS
Smith Mountain Lake	Bedford Co.	19,820	PWS
Spring Hollow Reservoir	Roanoke Co.	113	PWS
Stonehouse Creek Reservoir	Amherst Co.	34	
Talbott Reservoir	Patrick Co.	141	
Thrashers Creek Reservoir	Amherst Co.	32	
Townes Reservoir	Patrick Co.	28	

**Total 124 = Significant Reservoirs/Lakes statewide**

PWS = Public Water Supply Water Quality Standards Designation

DWR = Virginia Department of Wildlife Resources Owned or Managed

DCR = Virginia Department of Conservation and Recreation Owned or Managed

USACE = U.S. Army Corps of Engineers Owned or Managed

USFS = U.S. Forest Service Owned or Managed

# **APPENDIX I**

## **Memorandum**

TO: John Kennedy, Director of the Office of Ecology  
Sandra Mueller, Water Quality Monitoring and Assessment Manager  
Andrew Garey, Water Quality Monitoring Team Lead  
Mark Richards, TMDL Team Lead  
Amanda Shaver, Water Quality Assessment Coordinator  
David Whitehurst, Water Quality Standards Analyst

From: Tish Robertson, Water Quality Monitoring and Assessment Scientist

Subject: Need for standardized rounding method for water quality assessment

DATE: October 29, 2019

Assessment staff are instructed to list water bodies as impaired when data indicate a greater than 10.5% exceedance rate for conventional parameter criteria (DO, pH, temperature, and bacteria). This practice is based on guidelines from EPA's *Consolidated Assessment and Listing Methodology: Toward a Compendium of Best Practices*, which recommends conventional pollutant criteria be allowed a 10% exceedance frequency. For recreational bacteria criteria, this "10% rule" is codified in the water quality standards (WQS). Since this "10%" is expressed as a whole number in the WQS rather than as a value with decimal places, calculated exceedance frequencies should be rounded to the nearest whole number before evaluating them.

Under the most common method of rounding—wherein if the digit to be dropped is 5 or greater, the preceding digit is rounded up—an exceedance of 10.5% would be rounded up to 11%. Thus, under this method, a water body exhibiting an exceedance rate of 10.5% for the single sample maximum bacteria criterion can be said to be in non-compliance with the WQS.

However, under the "round to even" method an exceedance rate of 10.5% is rounded to 10%. If the digit to be dropped is 5, the preceding digit is rounded up if it is odd and kept the same if the digit is even. When rounding 10.5% to the nearest whole number, "0" (the preceding digit) is kept since it is even. Thus, under this rounding method a water body with an exceedance rate of 10.5% is in compliance with the WQS.

While it can be inferred from the "greater than 10.5% exceedance rate" rule stipulated in the assessment guidance manual that assessors are supposed to be using the "round to even" method, there is currently no reference to this method anywhere in the guidance manual. It is very possible some staff are unaware of this rounding method and thus may be applying it only for the evaluation of exceedance rates rather than for all assessment decisions. The "round to even" method is just as valid as the more commonly used "half round up" method, but it is hard to defend the combined use of those methods.

There is a strong policy justification for adopting the "round to even" method. VPDES staff are instructed to use the "round to even" method in their implementation guidance for permit limits and DMRs (DEQ Guidance memo no: 06-2016). EPA and other organizations involved in the preparation of promulgated analytical methods under Clean Water Act Section 304(h) are instructed to use the "round to even" method as well<sup>11</sup>. Thus, incorporating a statement into the assessment guidance that directs staff to use the "round to even" method would ensure consistency

---

<sup>11</sup> [EPA. \(1996\) Guidelines and Format for Methods to be proposed at 40 CFR Part 136 or Part 141. Office of Science and Technology. Washington, DC.](#)  
*Final Guidance for 2024 IR*  
*Appendix I*

with existing DEQ and EPA policy. A similar policy could also be adopted by the Watersheds program so that TMDLs and IPs are consistent with the other water quality programs.

If a “round to even” rule was incorporated into the assessment guidance manual, automated tools (Access, Excel, SAS, and R tools) will need to be programmed accordingly, since typically the “half round up” is the default method. Additionally, staff who are unfamiliar with this method will need to be introduced to it.