

Disclosures by Licensed Midwives for High-Risk Pregnancy Conditions

Virginia Board of Medicine

The Code of Virginia (Law) requires that licensed midwives “disclose to their patients, when appropriate, options for consultation and referral to a physician and evidence-based information on health risks associated with birth of a child outside of a hospital or birthing center.” Regulations for Licensed Midwives specify that:

Upon initiation of care, a midwife shall review the client's medical history in order to identify pre-existing conditions or indicators that require disclosure of risk for home birth. The midwife shall offer standard tests and screenings for evaluating risks and shall document client response to such recommendations. The midwife shall also continually assess the pregnant woman and baby in order to recognize conditions that may arise during the course of care that require disclosure of risk for birth outside of a hospital or birthing center.

The risk factors or conditions that require disclosures are listed in regulation. If any of these conditions or factors are presented, the midwife is to:

- 1) Request and review the client's medical history, including records of the current or previous pregnancies;*
- 2) Disclose to the client the risks associated with a birth outside of a hospital or birthing center; and*
- 3) Provide options for consultation and referral.*

Regulations require that if the risk factors or criteria have been identified that may indicate health risks associated with birth of a child outside a hospital or birthing center, the midwife must provide evidence-based information on such risks and must document in the client record the assessment of all health risks that pose a potential for a high risk pregnancy and, if appropriate, the provision of disclosures and evidence-based information. **The disclosure for intrapartum risk factors should be given to a client at the first prenatal visit.**

For each of the risk factors or conditions identified, this guidance document provides evidence-based information and a format to record in a client's record the disclosure of information and options for consultation and referral.

To access the evidence-based information and disclosure for a particular conditions or risk factor, click on the link in the index below. The midwife may then print the form for that condition or risk factor for presentation and discussion with the client and have the form signed for inclusion in the client record.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

Intrapartum Risk Factors

1. Abnormal fetal cardiac rate or rhythm
2. Active cancer
3. Acute or chronic thrombophlebitis
4. Anemia (hematocrit less than 30 or hemoglobin less than 10 at term)
5. Any pregnancy with abnormal fetal surveillance tests
6. Blood coagulation defect
7. Body Mass Index (BMI) equal to or greater than 30
8. Cardiac disease
9. Chronic obstructive pulmonary disease including asthma
10. Ectopic pregnancy
11. Essential chronic hypertension over 140/90
12. Genital herpes or partner with genital herpes
13. History of hemoglobinopathies
14. HIV positive status with AIDS
15. Inappropriate fetal size for gestation – Macrosomia (Large for gestational age)
16. Inappropriate fetal size for gestation – IUGR (Small for gestational age)
17. Incomplete spontaneous abortion
18. Isoimmunization to blood factors
19. Multiple gestation
20. Persistent severe abnormal quantity of amniotic fluid
21. Platelet count less than 120,000
22. Position presentation other than vertex at term or while in labor
23. Pre-eclampsia/eclampsia
24. Pregnancy lasting longer than 42 completed weeks with an abnormal non-stress test
25. VBAC (vaginal birth after cesarian) previous uterine incision or myomectomy
26. Psychiatric disorders (Mental Health Disorders)
27. Rupture of membranes 24 hours before the onset of labor
28. Seizure disorder requiring prescriptive medication

29. Severe liver disease -- active or chronic
30. Severe renal disease - active or chronic
31. Significant 2nd or 3rd trimester bleeding
32. Significant glucose intolerance (Preexisting diabetes, gestational diabetes, PCOS)
33. Uncontrolled hyperthyroidism
34. Uterine ablation (endometrial ablation)
35. Uterine anomaly

Intrapartum Risk Factors

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the North American Registry of Midwives).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Conditions requiring on-going medical supervision or on-going use of medications

Clients with chronic medical conditions, on prescribed medications, or under medical care for a time-limited problem that coincides with pregnancy should be advised to consult with their treating healthcare providers regarding the impact of these conditions and medications on pregnancy, as well as any impact pregnancy may have on their other diagnosed conditions. Women who choose not to disclose information regarding any medical conditions they have or medications that they are taking may increase their risk of complications.

Current substance abuse (including alcohol and tobacco)

Obstetrical complications of cigarette smoking include:

- Growth restriction (IUGR)
- Spontaneous abortion (miscarriage)
- Sudden infant death syndrome (SIDS)

Alcohol abuse leads to:

- Nutritional deficiencies
- Fetal alcohol syndrome

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

In addition to increased risk of preterm labor and baby being small for gestational age, complications resulting from abusing other drugs include:

- Heroin and cocaine consumption result in medical, nutritional and social neglect
- Cocaine and amphetamine cause hypertension, placental abruption
- Intravenous abuse also increases the risk of contracting infectious disease.¹
- Maternal substance use of opioids, benzodiazepines, barbiturates, and alcohol can cause NAS (Neonatal abstinence syndrome).² NAS is a set of drug withdrawal symptoms that affect the central nervous, gastrointestinal, and respiratory systems in the newborn when separated from the placenta at birth.

Documented Intrauterine growth retardation (IUGR)/small for gestational age (SGA) at term

Complications³ for the growth-restricted fetus include:

- Prematurity
- Perinatal morbidity
- Stillbirth

"IUGR is a serious problem, regardless of why the baby is small. About 20% of stillborn babies are IUGR, and perinatal mortality for growth-restricted infants may be 6 to 10 times higher than for those of normal size. Most IUGR stillbirths occur after the 36th week of pregnancy and before labor begins."⁴

Suspected uterine rupture

Consequences of uterine rupture:

- There have been no reported maternal deaths due to uterine rupture
- Overall, 14 percent to 33 percent of women will need a hysterectomy when the uterus ruptures
- Approximately 6 percent of uterine ruptures will result in perinatal death
- This is an overall risk of intrapartum fetal death of 20 per 100,000 women undergoing trial of labor after previous cesarean section
- "For term pregnancies, the reported risk of fetal death with uterine rupture is less than 3 percent. Although the risk is similarly low, there is insufficient evidence to quantify the neonatal morbidity directly related to uterine rupture."⁵

Prolapsed cord or cord presentation

Prolapsed cord is a term describing a cord that is passing through the cervix at the same time or in advance of the fetal presenting part. This occurs in approximately 1.4-6.2 per 1000 of pregnancies. Although uncommon, it is considered a true obstetrical emergency most often necessitating a caesarean delivery. Prolapsed cord is associated with other complications of pregnancy and delivery as well.

¹ Pregnancy and substance abuse, G. Fischer, M. Bitschnau, A. Peterzell, H. Eder, A. Topitz. Archives of Women's Mental Health. August 1999, Volume 2, Issue 2, pp 57-65.

² Casper, Tammy, and Megan W. Arbour. "Identification of the Pregnant Woman Who Is Using Drugs: Implications for Perinatal and Neonatal Care." Journal of Midwifery & Women's Health (2013).

³ Lerner, Jodi P. "Fetal growth and well-being." Obstetrics and gynecology clinics of north America 31.1 (2004): 159-176.

⁴ Frye, Anne, *Holistic Midwifery, Volume I*, Labrys Press, Portland, OR, 2006, p. 990

⁵ Guise, Jeanne-Marie, et al. "Vaginal birth after cesarean: new insights." (2010).

Fetal risks:

- Hypoxia
- Stillbirth/death

Suspected complete or partial placental abruption

Placental abruption results from a cascade of pathophysiologic processes ultimately leading to the separation of the placenta prior to delivery. Pregnancies complicated by abruption result in increased frequency⁶ of:

- Low birth weight
- Preterm delivery
- Stillbirth
- Perinatal death

Suspected placental previa

Pregnancies complicated with placenta previa had significantly higher rates⁷ of

- Second-trimester bleeding
- Pathological presentations
- Placental abruption
- Congenital malformations
- Perinatal mortality
- Cesarean delivery
- Apgar scores at 5 minutes lower than 7
- Placenta accreta
- Postpartum hemorrhage
- Postpartum anemia
- Delayed maternal and infant discharge from the hospital

Suspected chorioamnionitis

Chorioamnionitis is a potentially serious complication:⁸

- Chorioamnionitis is a major risk factor in the event of preterm birth, especially at earlier gestational ages, contributing to prematurity-associated mortality and morbidity
- Increased susceptibility of the lung for postnatal injury, which predisposes for bronchopulmonary dysplasia.
- Chorioamnionitis is associated with cystic periventricular leukomalacia, intraventricular hemorrhage and cerebral palsy in preterm infants
- Prenatal inflammation/infection has been shown a risk factor for neonatal sepsis

⁶ Ananth, Cande V., et al. "Placental abruption and adverse perinatal outcomes." *JAMA: the journal of the American Medical Association* 282.17 (1999): 1646-1651.

⁷ Sheiner, E., et al. "Placenta previa: obstetric risk factors and pregnancy outcome." *Journal of Maternal-Fetal and Neonatal Medicine* 10.6 (2001): 414-419.

⁸ Thomas, Wolfgang, and Christian P. Speer. "Chorioamnionitis: important risk factor or innocent bystander for neonatal outcome?." *Neonatology* 99.3 (2010): 177-187.

Pre-eclampsia/eclampsia

Complications of preeclampsia include:

- Eclampsia
- HELLP (hemolysis, elevated liver enzymes, low platelets) syndrome
- Liver rupture
- Pulmonary edema
- Renal failure
- Disseminated intravascular coagulopathy (DIC)
- Hypertensive emergency
- Hypertensive encephalopathy
- Cortical blindness

Maternal complications occur in up to 70% of women with eclampsia and include:⁹

- DIC
- Acute renal failure
- Hepatocellular injury
- Liver rupture
- Intracerebral hemorrhage
- Cardiopulmonary arrest
- Aspiration pneumonitis
- Acute pulmonary edema
- Postpartum hemorrhage
- Maternal death rates of 0-13.9% have been reported

Fetal complications in preeclampsia are directly related to gestational age and the severity of maternal disease and include increased rates of:¹⁰

- Preterm delivery
- Intrauterine growth restriction
- Placental abruption
- Perinatal death

Thick meconium stained amniotic fluid without reassuring fetal heart tones and birth is not imminent

Meconium staining of the amniotic fluid is a common occurrence during labor. Although a large proportion of these pregnancies will have a normal neonatal outcome, its presence may be an indicator of fetal hypoxia and has been linked to the development of:¹¹

- Cerebral palsy

⁹ Norwitz, Errol R., Chaur-Dong Hsu, and John T. Repke. "Acute complications of preeclampsia." *Clinical obstetrics and gynecology* 45.2 (2002): 308-329.

¹⁰ de Souza Rugolo, Ligia Maria Suppo, Maria Regina Bentlin, and Cleide Enoir Petean Trindade. "Preeclampsia: effect on the fetus and newborn." *Neoreviews* 12.4 (2011): e198-e206.

¹¹ Rahman, Shimma, Jeffrey Unsworth, and Sarah Vause. "Meconium in labour." *Obstetrics, Gynaecology & Reproductive Medicine* 23.8 (2013): 247-252.

- Seizures
- Meconium aspiration syndrome

Abnormal auscultated fetal heart rate pattern unresponsive to treatment or inability to auscultate fetal heart tones

Sustained abnormal fetal heart rate patterns include bradycardia (abnormally low heart rate) and decelerations in the baby's heart rate. Additionally, tachycardia (abnormally high heart rate) is abnormal, and can also be an indication for the need for further evaluation. Historically, a 30-minute rule from decision-to-incision time for emergent cesarean delivery in the setting of abnormal FHR pattern has existed; however, the scientific evidence to support this threshold is lacking.

Excessive vomiting, dehydration, or exhaustion unresponsive to treatment

- Sufficient fluid intake during labor may prevent hemoconcentration, starvation, and activation of the thrombogenic and fibrinolytic system¹²
- With extreme exhaustion, the chances of fetal distress and non-progressive labor are greatly increased
- Bleeding during or after the placental birth, followed by shock, are much more likely to occur when the woman and her uterus are exhausted¹³
- Maternal exhaustion is diagnosed with a combination of ketonuria, elevated temperature, and elevated pulse. This condition is also known as ketoacidosis, in that the mother's blood becomes abnormally acidic and less able to carry oxygen. Unless this condition is reversed, fetal distress will result¹⁴

Blood pressure greater than 140/90 which persists or rises and birth is not imminent

Women with chronic hypertension are at increased risk of:¹⁵

- Superimposed preeclampsia (25% risk)
- Preterm delivery
- Fetal growth restriction or demise
- Placental abruption
- Congestive heart failure
- Acute renal failure
- Seizures
- Stroke
- Death

Maternal fever equal to or greater than 100.4°

Fever can indicate infection. Fever in labor is associated with:¹⁶

- Early neonatal and infant death
- Hypoxia

¹² Watanabe, Takashi, et al. "Effect of labor on maternal dehydration, starvation, coagulation, and fibrinolysis." *Journal of perinatal medicine* 29.6 (2001): 528-534.

¹³ Frye, Anne, *Holistic Midwifery, Volume II*, Labrys Press, Portland, OR, 2004, p. 1055.

¹⁴ Davis, Elizabeth, *Heart and Hands: A Midwife's Guide to Pregnancy and Birth*, Celestial Arts, New York, NY, 2004, p. 141.

¹⁵ Hypertension. 2003; 41: 437-445 Published online before print February 10, 2003, doi: 10.1161/01.HYP.0000054981.03589.E9

¹⁶ PETROVA, Anna, et al. "Association of maternal fever during labor with neonatal and infant morbidity and mortality." *Obstetrics and gynecology* 98.1 (2001): 20-27.

- Infection-related death. These associations were stronger among term than preterm infants
- Meconium aspiration syndrome
- Hyaline membrane disease
- Neonatal seizures
- Assisted ventilation

Labor or premature rupture of membrane (PROM) less than 37 weeks according to due date

Premature rupture of membranes before 37 weeks’ gestation (and where there is at least an hour between membrane rupture and the onset of contractions and labor) can have consequences for both the mother and the baby:

Risks to Baby:

- Neurologic injury
- Infection
- Respiratory Distress
- Death
- Increased need for neonatal intensive care services

Maternal Risks:

- Infection
- Prolonged Labor
- C-Section
- Death

Because the out-of-hospital birth setting does not provide for immediate access to medications, surgery, and consultation with a physician, there may be increased risks to mother and/or baby if any of these conditions present during the birth. In some communities, the lack of availability of a seamless, cooperative hospital transfer process adds additional risk during intrapartum transfer.

I understand that the intrapartum risks may not be apparent until labor, and my opportunity for referral to a physician, should I choose that, would be limited to hospital transfer and transfer of care to the physician on call at that facility.

I have received and read this document, discussed it with my midwife, and my midwife has answered my questions to my satisfaction.

Client _____

Date _____

Midwife _____

Date _____

[HOME](#)

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

1. ABNORMAL FETAL CARDIAC RATE OR RHYTHM

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the **North American Registry of Midwives**).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Abnormal fetal cardiac rate or rhythm

Fetal rhythm abnormalities (fetal heart rates that are irregular, too fast or too slow):

- occur in up to 2% of pregnancies
- usually identified by the obstetrical clinician who detects an abnormal fetal heart rate or rhythm using a Doppler or stethoscope
- majority have isolated premature atrial contractions which may spontaneously resolve
- sustained tachyarrhythmia (rapid) or bradyarrhythmia (slow) may be of clinical significance
 - may indicate severe systemic disease
 - may have the potential to compromise the fetal circulation
 - May require intensive antepartum and/or neonatal care

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

[HOME](#)

Congenital heart disease: Rhythm abnormalities of the fetus. Lisa K Hornberger, David J Sahn. Heart 2007;93:10 1294-1300 doi:10.1136/hrt.2005.069369

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

2. ACTIVE CANCER

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the **North American Registry of Midwives**).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Active Cancer

Maternal risks:

- maternal infection due to immune suppression,
- deep vein thrombosis and pulmonary embolism during pregnancy and especially after delivery
- hemorrhage at delivery.

Fetal risks:

- Intrauterine growth restriction
- Preterm birth
- Fetal health effects from exposure to maternal medications

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

<http://www.nlm.nih.gov/medlineplus/cancerandpregnancy.html> J Obstet Gynaecol Can. 2013 Mar;35(3):263-80.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

3. ACUTE OR CHRONIC THROMBOPHLEBITIS

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Deep vein thrombosis (DVT) and pulmonary embolism (PE) are collectively known as venous thromboembolism (VTE). VTE occurs more frequently in pregnant women, with an incidence of 0.5 to 2.0 per 1000 pregnancies, four to five times higher than in the non-pregnant population. The risk for VTE is further elevated in the postpartum period.

The risk for VTE in pregnancy is increased in women with:

- **Prior history of VTE**
- **Advanced maternal age**
- **Collagen-vascular disease, especially antiphospholipid antibody syndrome**
- **Obesity (BMI > 30)**
- **Multiparity**
- **Hypercoagulable state**
- **Nephrotic syndrome**
- **Operative delivery**
- **Prolonged bed rest**
- **Hematologic disorders (hemoglobin SS and SC disease, polycythemia, thrombotic thrombocytopenic purpura, paroxysmal nocturnal hemoglobinuria, and some dysfibrinogenemias).**
- **Maternal medical conditions (diabetes, heart disease, inflammatory bowel disease)**
- **Smoking**
- **Preeclampsia**

Maternal complications:

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- **hypoxemia**
- **post-phlebotic syndrome**
- **pulmonary infarction**
- **death**

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

Chisholm CA, James AH, Ferguson JE. Thromboembolic disorders. In: Evans AE, Manual of Obstetrics, 8th edition. 2014, Wolters Kluwers Health.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

4. ANEMIA (HEMATOCRIT LESS THAN 30 OR HEMOGLOBIN LESS THAN 10 AT TERM)

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

"If a midwife supports a client's choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process." – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Anemia (hematocrit less than 30 or hemoglobin less than 10 at term)

The World Health Organization (WHO) estimates that worldwide, 42% of pregnant women are anemic.¹⁷

Current knowledge indicates that iron deficiency anemia in pregnancy is a risk factor for preterm delivery and subsequent low birth weight, and possibly for inferior neonatal health. Data are inadequate to determine the extent to which maternal anemia might contribute to maternal mortality.¹⁸

...a woman who is already anemic is unable to tolerate blood loss that a healthy woman can.¹⁹

Maternal Risks related to severe or untreated anemia:

- need for blood transfusion(s), resulting from a hemorrhage (significant blood loss) during delivery
- postpartum depression

Fetal/Neonatal Risks related to maternal severe or untreated anemia:

- prematurity
- low-birth-weight
- anemia
- developmental delays

¹⁷ Benoist B, McLean E, Egli I, et al. *Worldwide Prevalence of Anaemia 1993-2005*. Geneva, Switzerland: World Health Organization; 2008.

¹⁸ Allen, Lindsay H. "Anemia and iron deficiency: effects on pregnancy outcome." *The American journal of clinical nutrition* 71.5 (2000): 1280s-1284s.

¹⁹ McCormick, M. L., et al. "Preventing postpartum hemorrhage in low-resource settings." *International journal of gynecology & obstetrics* 77.3 (2002): 267-275.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

5. ANY PREGNANCY WITH ABNORMAL FETAL SURVEILLANCE TESTS

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the North American Registry of Midwives).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Pregnancy with abnormal Fetal Surveillance Tests

There is no benefit in continuing a pregnancy at or post term after fetal surveillance is found to be non-reassuring. The recommendation is delivery (Price, 2014).” Abnormal stress tests at any point in pregnancy are associated with an increased risk of poor outcomes in pregnancy and during labor and delivery. Babies with diagnosed or undiagnosed anomalies are more likely to have abnormal test results requiring specialized care before or after delivery. Antepartum testing results, with regard to the overall clinical picture, should be taken seriously.

Risks to fetus:

- Stillbirth
- Asphyxia
- Fetal Acidosis
- Low Apgar scores
- Respiratory distress
- Surgical delivery
- Meconium Aspiration
- Death

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

O'Neill, E. T. (2012). Antepartum evaluation of the fetus and fetal well-being. *Clinical Obstetrics and Gynecology* , 55 (3), 722.
Preboth, M. (2000). Practice Guidelines ACOG Guidelines on Antepartum Fetal Surveillance . *Am Fam Physician* .
Price, A. (2014, January). MSN CNM. Assistant Clinical Professor VCUMC. (B. Sheets, Interviewer)
Singh, T. (2008). The prediction of intra-partum fetal compromise in prolonged pregnancy. *Journal of Obstetrics and Gynecology* , 28 (8), 779-782.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

6. BLOOD COAGULATION DEFECT

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Blood coagulation defect

Hereditary thrombophilia, or predisposition to thrombosis, ranges from the common (Factor V Leiden heterozygosity, present in 1-15% of pregnant women) to the rare (antithrombin deficiency occurring in 0.02%). The risk of deep vein thrombosis or pulmonary embolism (collectively known as venous thromboembolism or VTE) ranges from 0.1-7% of pregnancies. The maternal medical history determines the management during pregnancy, which can include anticoagulation with injections of heparin throughout the pregnancy and post-partum period.

The presence of one of these disorders may contribute to the risk of obstetric complications as well, including:

- IUGR
- preeclampsia
- stillbirth
- Frequent fetal surveillance is recommended in most cases, as well as timed delivery in the last week before the estimated date of delivery.

Alternatively, disorders of maternal hemostasis (such as von Willebrand disease) increase the risk of blood loss at delivery, and as hereditary disorders also increase the risk for abnormal bleeding in the newborn.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

Inherited Thrombophilia in Pregnancy. Practice Bulletin 138, November 2013. American College of Obstetricians and Gynecologists.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

7. BODY MASS INDEX (BMI) EQUAL TO OR GREATER THAN 30

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the North American Registry of Midwives).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Body Mass Index (BMI) equal to or greater than 30

Obesity is defined as having a BMI of 30 or higher. The number of obese women in the United States has increased greatly during the past 25 years. Obesity has also become a major health concern for pregnant women. More than one half of pregnant women are overweight or obese.

Risks of Obesity Include:

- Birth defects – Babies born to obese mothers have an increased risk of having birth defects, such as heart defects and neural tube defects.
- Macrosomia – In this condition, the baby is larger than normal. This can increase the risk of the baby being injured during birth. For example, the baby’s shoulder can become entrapped after the head is delivered. Macrosomia also increases the risk of cesarean birth.
- Preterm Birth – Problems associated with a mother’s obesity may mean that the baby will need to be delivered early. Preterm infants have an increased risk of health problems, including breathing problems, eating problems, and developmental and learning difficulties later in life.
- Stillbirth – The risk of stillbirth increases the higher the mother’s BMI.
- High Blood Pressure
- Preeclampsia – Preeclampsia is a serious illness for both the woman and her baby. Although gestational hypertension is the most common sign of preeclampsia, this condition affects all organs of the body. The kidneys and liver may fail. In rare cases, stroke can occur. The fetus is at risk of growth problems and problems with the placenta. It may require early delivery, even if the baby is not fully grown. In severe cases, the woman, baby, or both may die.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- Gestational Diabetes – High blood glucose (sugar) levels during pregnancy increase the risk of having a very large baby and a cesarean delivery. Women who have had gestational diabetes have a higher risk of having ~~babies~~ diabetes in the future, as do their children.
- Challenges in Prenatal Care – Obesity can make it more difficult for the midwife to assess fetal position and fetal growth.
- Challenges in Labor Management – Obesity can create challenges in moving the woman quickly in the event of an emergency during the birth, and can make auscultation of fetal heart tones more difficult.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

8. CARDIAC DISEASE

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the **North American Registry of Midwives**).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Cardiac Disease

Most women tolerate the cardiovascular changes of pregnancy without difficulty. Pregnancy in a patient with significant cardiac disease is associated with significant risk. Despite occurring in only 0.2-4% of pregnancies, cardiac disease is associated with up to 30% of maternal deaths. A pregnant patient with cardiac disease will benefit from the coordinated care of a multidisciplinary team including perinatologists, cardiologists and anesthesiologists. In particular, adults with repaired congenital heart disease may pose complex management scenarios. They may require specialized cardiac monitoring during labor and birth, and some cardiac conditions are associated with a high enough risk of labor complications that cesarean is recommended.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

Nanda S, Nelson-Piercy C, Mackillop L. Cardiac disease in pregnancy. Clin Med 2012;12:553-560.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

9. CHRONIC OBSTRUCTIVE PULMONARY DISEASE INCLUDING ASTHMA (1)

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Asthma affects approximately 4% to 6% of adults of all ages and is one of the most common medical conditions complicating pregnancy.

RISKS

- Preterm birth
- Decreased birth weight
- Increased neonatal and maternal death

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

(1) http://www.glowm.com/section_view/heading/Pulmonary%20Disease%20in%20Pregnancy/item/170#1199

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

10. ECTOPIC PREGNANCY (1)

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Today, about 1 in 50 pregnancies is ectopic. An ectopic pregnancy occurs when a fertilized egg grows outside of the uterus most commonly in the tube. As the pregnancy grows, it can rupture (burst). If this occurs, it can cause major internal bleeding. This can be life threatening and needs to be treated with surgery.

RISKS

- Fallopian tube damaged, leading to an increased likelihood of having another ectopic pregnancy in the future.
- Ruptured ectopic pregnancy (when the fallopian tube splits) and severe internal bleeding, which can lead to shock.
- Death

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

(1) <http://www.webmd.boots.com/pregnancy/tc/ectopic-pregnancy-complications-of-ectopic-pregnancy>

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

11. ESSENTIAL CHRONIC HYPERTENSION (1)

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Elevated blood pressure, systolic >140 or diastolic >90 or both, that predates conception or is diagnosed before 20 weeks of gestation.

MATERNAL RISKS

- Preterm delivery
- Placental abruption
- Preeclampsia
- Eclampsia
- Seizures
- Maternal congestive heart failure
- Acute renal failure
- Death

FETAL RISKS

- Fetal growth restriction
- Fetal death

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

(1) http://www.nhlbi.nih.gov/health/public/heart/hbp/hbp_preg.htm

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

12. GENITAL HERPES OR PARTNER WITH GENITAL HERPES

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of Risks Related to: Genital Herpes

Because of its serious and potentially lethal risks to the fetus and neonate, pregnant women and their partners should be tested for **HSV - Herpes Simplex Virus** (HSV1 & HSV2).

In women with a previous diagnosis of genital herpes, cesarean delivery to prevent neonatal HSV infection is not indicated if there are NO genital lesions at the time of labor. In an effort to reduce cesarean deliveries performed for the indication of genital herpes, the use of oral acyclovir or valacyclovir near the end of pregnancy to suppress genital HSV recurrences has become increasingly common in obstetric practice. Several studies with small sample sizes suggest that suppressive acyclovir therapy during the last weeks of pregnancy decreases the occurrence of clinically apparent genital HSV disease at the time of delivery, with an associated decrease in cesarean delivery rates for the indication of genital HSV. **However, because viral shedding still occurs (albeit with reduced frequency), the potential for neonatal infection is not avoided completely, and cases of neonatal HSV disease in newborn infants of women who were receiving antiviral suppression recently have been reported.**²⁰

Genital HSV, especially in primary infections, may be dangerous to the neonate if infected during delivery, as it can cause a severe neonatal disease.²¹

Risks of HSV infection to the fetus include:

²⁰ Kimberlin, David W., et al. "Guidance on management of asymptomatic neonates born to women with active genital herpes lesions." *Pediatrics* 131.2 (2013): e635-e646.

²¹ Meytal Avgil, Asher Ornoy, Herpes simplex virus and Epstein-Barr virus infections in pregnancy: consequences of neonatal or intrauterine infection, *Reproductive Toxicology*, Volume 21, Issue 4, May 2006, Pages 436-445, ISSN 0890-6238, <http://dx.doi.org/10.1016/j.reprotox.2004.11.014>.

- intrauterine fetal demise (the death of the fetus while in the uterus)
- skin scars (cutaneous manifestations),
- ophthalmologic findings (chorioretinitis, microphthalmia),
- neurological involvement (causing brain damage)

The clinical presentation of infants with neonatal HSV infection, that is almost invariably symptomatic and frequently lethal, is a direct reflection of the site and extent of viral replication.²²

Risks of HSV infection to the neonate (newborn) include:

- death
- neurologic (brain) damage (intracranial calcifications, microcephaly, seizures, encephalomacia),
- growth restriction,
- psychomotor development impairment
- skin vesicles or scarring,
- eye lesions resulting in vision loss and/or blindness (chorioretinitis, microphthalmia, cataracts),
- hearing loss and/or deafness

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

²² Anzivino, Elena, et al. "Herpes simplex virus infection in pregnancy and in neonate: status of art of epidemiology, diagnosis, therapy and prevention." *Virology* 6.1 (2009): 1-11.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

13. HISTORY OF HEMOGLOBINOPATHIES

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the **North American Registry of Midwives**).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: History of hemoglobinopathies

Hemoglobinopathies include sickle cell disease and its variants as well as alpha and beta thalassemia. The involvement of a multidisciplinary team including perinatologists, hematologists and anesthesiologists can allow for development of a plan to screen for and manage complications.

Maternal risks include:

- cerebral vein or deep vein thrombosis
- anemia and vaso-occlusive crisis
- pneumonia
- pyelonephritis
- transfusion
- pregnancy induced hypertension
- postpartum infection, sepsis, and systemic inflammatory response syndrome
- cesarean delivery

Fetal risks include:

- preterm birth and its consequences including low birth weight
- intrauterine growth restriction
- abruption placentae
- stillbirth

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- genetic risk assessment is also recommended for individuals identified as carriers for hemoglobinopathy, as they may be at risk to have affected offspring.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

Villers, Margaret S., et al. "Morbidity associated with sickle cell disease in pregnancy." American journal of obstetrics and gynecology 199.2 (2008): 125-e1.
 Naik, Rakhi P., and Sophie Lanzkron. "Baby on board: what you need to know about pregnancy in the hemoglobinopathies." ASH Education Program Book 2012.1 (2012): 208-214.
 John C. Morrison and Marc R. Parrish. "Sickle Cell Disease and Other Hemoglobinopathies" Protocols for High-Risk Pregnancies (2010): 158-159.
 American College of Obstetricians and Gynecologists, Practice Bulletin 78, "Hemoglobinopathy in Pregnancy," January 2007

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

14. HIV POSITIVE STATUS WITH AIDS

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence- based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: HIV positive status with AIDS

HIV transmission from mother to child during pregnancy, labor and delivery, or breastfeeding is known as perinatal transmission and is the most common route of HIV infection in children. When HIV is diagnosed before or during pregnancy, perinatal transmission can be reduced to less than 1% if appropriate medical treatment is given, the virus becomes undetectable, and breastfeeding is avoided.²³

Recommended medical treatment includes antiretroviral medication taken throughout pregnancy and during labor, regular monitoring of the maternal viral load, cesarean delivery for viral load > 1000 copies/mL, and initiation of antiretroviral medication for the newborn shortly after birth.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

15. INAPPROPRIATE FETAL SIZE FOR GESTATION – MACROSOMIA (LARGE FOR GESTATIONAL AGE)

²³ <http://www.cdc.gov/hiv/risk/gender/pregnantwomen/index.html>

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of Risks Related to: Inappropriate Fetal Size for Gestation – Macrosomia (Large for Gestational Age)

Macrosomia (meaning **big body**), is arbitrarily defined as a birth weight of more than 4,000 g (8 lb, 13 oz). Also known as **large for gestational age**, fetal macrosomia complicates more than 10 percent of all pregnancies in the United States.²⁴

Risks to the mother related to macrosomia include:

- increased risk of uterine rupture after previous cesarean section or other uterine surgery;
- increased likelihood of induction at or before 40 weeks;
- increased likelihood of an operative delivery: forceps, vacuum, or cesarean section;
- trauma to vagina and/or perineum; including perineal and/or vulvar lacerations, 3rd or 4th degree episiotomy, short or long-term urinary or fecal incontinence;
- increased blood loss and/or postpartum hemorrhage,
- damage to the coccyx (tailbone)

Risks to the baby related to macrosomia at the time of birth include:

- shoulder dystocia (the baby gets stuck at the shoulders after the delivery of the head), which may result in trauma to the baby including:
 - broken clavicle (collar) bone(s);
 - brachial plexus injury, temporary or permanent nerve damage (sensory and motor) to either one or both shoulders, arms, and hands;
 - cerebral palsy;
 - hypoxia, resulting in permanent brain damage;
 - death.
- injuries related to operative delivery (forceps, vacuum, or cesarean section) including:
 - bruising and/or injury to the scalp, head and/or face;

²⁴ MARK A. ZAMORSKI, M.D., M.H.S.A., and WENDY S. BIGGS, M.D., University of Michigan Medical School, Ann Arbor, Michigan. Am Fam Physician. 2001 Jan 15;63(2):302-307.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- temporary weakness in the facial muscles (facial palsy);
- external eye and/or ear trauma;
- broken clavicle (collar) bone(s);
- brachial plexus injury (see description above);
- cerebral palsy;
- skull fracture;
- bleeding within the skull;
- seizures;
- lacerations (during cesarean section) to the baby’s presenting part
- immature lungs and breathing problems, if the due date has been miscalculated and the infant is delivered before 39 weeks of gestation;
- need for special care in the neonatal intensive care unit (NICU);

Risks to the newborn related to macrosomia and later childhood risks:

- higher than normal blood sugar level (impaired glucose tolerance);
- childhood obesity (research suggests that the risk of childhood obesity increases as birth weight increases);
- metabolic syndrome (a group of conditions: increased blood pressure, a high blood sugar level, excess body fat, abnormal cholesterol levels; that occur together, increasing the risk of heart disease, stroke and diabetes later in life.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

16. INAPPROPRIATE FETAL SIZE FOR GESTATION – IUGR (SMALL FOR GESTATIONAL AGE)

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of Risks Related to: Inappropriate Fetal Size for Gestation – IUGR (Small for Gestational Age)

IUGR (Intrauterine Growth Restriction) is a serious problem, regardless of why the baby is small. About 20% of stillborn babies are IUGR, and perinatal mortality for growth-restricted infants may be 6 to 10 times higher than for those of normal size. Most IUGR stillbirths occur after the 36th week of pregnancy and before labor begins.²⁵

Risks to the baby related to IUGR, known as Small for Gestation Age:

- low birth weight (LBW);
- difficulty handling the stresses of vaginal delivery;
- decreased oxygen levels (hypoxia);
- hypoglycemia (low blood sugar);
- low resistance to infection;
- low APGAR scores (a test given immediately after birth to evaluate the newborn's physical condition and determine need for special medical care);
- meconium aspiration (inhalation of stools passed while in the uterus), which can lead to breathing problems, lung surfactant dysfunction, chemical pneumonitis, and persistent pulmonary hypertension;
- trouble maintaining body temperature (hypothermia);
- abnormally high red blood cell count;
- admission to NICU;
- long-term growth problems;
- intrauterine fetal demise (fetal death prior to labor);

²⁵ Frye, Anne, *Holistic Midwifery, Volume I*, Labrys Press, Portland, OR, 2006, p. 990

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- stillbirth (fetal death during labor or birth).

Risks to the mother related to IUGR:

- increased stress related to fetal monitoring and surveillance (serial ultrasounds and non-stress testing);
- premature labor;
- premature birth (delivery of the fetus before 37 weeks gestation);
- induction and early delivery, before 40 weeks;
- cesarean section.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

<p>A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.</p>
--

17. INCOMPLETE SPONTANEOUS ABORTION OR INCOMPLETE MISCARRIAGE (10)

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the North American Registry of Midwives).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Spontaneous abortion also known as early pregnancy loss refers to a miscarriage that happens before 20 weeks of gestation and is seen in 13% to 20% of all diagnosed pregnancies. Incomplete spontaneous abortion occurs when some tissue is retained in the uterus. Medication or a procedure may be needed to remove the tissue.

STILLBIRTH OR INTRAUTERINE FETAL DEMISE (IUFD)

Fetal death that happens after 20 weeks of gestational age is called stillbirth and has a rate of 3.2 per 1000 births. Medical intervention is needed for delivery.

MATERNAL FETAL RISKS OF EARLY OR LATE FETAL LOSS

- Infection
- Hemorrhage
- Maternal coagulopathy
- Gestational trophoblastic disease
- Rh isoimmunization

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

http://www.acog.org/Resources_And_Publications/Practice_Bulletins/Committee_on_Practice_Bulletins_--_Obstetrics/Management_of_Stillbirth

http://www.acog.org/Resources_And_Publications/Patient_Education_Pamphlets/Files/Early_Pregnancy_Loss

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

18. ISOIMMUNIZATION TO BLOOD FACTORS

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Isoimmunization to blood factors

Pregnant women with a negative Rh blood type (O-, A-, B-, AB-) or with other atypical antibodies have significant fetal and neonatal risk factors. Clinical manifestations of RhD haemolytic disease (HDN) range from asymptomatic mild anemia to hydrops fetalis or stillbirth associated with severe anemia and jaundice.²⁶

Risks to the baby related to maternal isoimmunization include:

- destruction of fetal red blood cells (hemolysis);
 - mild to moderate hemolysis manifests as increased indirect bilirubin (red cell pigment).
 - severe hemolysis leads to red blood cell production by the spleen and liver.
- severe anemia;
- hepatic circulatory obstruction (portal hypertension);
- placental edema, interfering with placental perfusion;
- ascites (accumulation of fluid in the abdominal cavity);
- hepatomegaly (swelling of the liver);
- increased placental thickness;
- polyhydramnios (increased amniotic fluid);
- hydrops (fetal heart failure);
- anasarca (extreme generalized edema);
- effusions (abnormal accumulation of fluid);
- intrauterine fetal demise (fetal death);
- stillbirth.

²⁶ Urbaniak, S. J., and M. A. Greiss. "RhD haemolytic disease of the fetus and the newborn." *Blood reviews* 14.1 (2000): 44-61.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

19. MULTIPLE GESTATION

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Multiple gestation

Maternal risks:

- Anemia
- Hemorrhage
- Preeclampsia
- Gestational diabetes
- Cesarean delivery

Fetal risks:

- Twin-to-twin transfusion syndrome (TTTS) in monochorionic twins
- Vanishing twin/death of one fetus
- Congenital anomalies
- Hydramnios
- Preterm birth
- Malpresentation
- Small for gestational age
- Umbilical cord prolapse
- Neonatal intensive care unit admission

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

Rao, Anita, Shanthi Sairam, and Hassan Shehata. "Obstetric complications of twin pregnancies." *Best Practice & Research Clinical Obstetrics & Gynaecology* 18.4 (2004): 557-576.
Spellacy, W. N. "Antepartum complications in twin pregnancies." *Clinics in perinatology* 15.1 (1988): 79-86.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

20. PERSISTENT SEVERE ABNORMAL QUANTITY OF AMNIOTIC FLUID (OLIGOHYDRAMNIOS AND POLYHYDRAMNIOS)

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Persistent severe abnormal quantity of amniotic fluid

Oligohydramnios (decreased amniotic fluid) may be caused by fetal anomalies (bladder outlet obstruction, renal agenesis), premature rupture of the membranes, or placental insufficiency occurring de novo or as a consequence of maternal conditions such as hypertension.

Maternal risks:

- antepartum hospitalization
- induction of labor
- cesarean delivery

Fetal risks:

- pulmonary hypoplasia (underdevelopment of the lungs)
- limb contractures
- abnormal fetal heart rate patterns
- acidosis
- neonatal intensive care unit admission
- need for surgical intervention if anomalies present
- stillbirth or neonatal death

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

Polyhydramnios (increased amniotic fluid) is most commonly idiopathic (no identifiable cause) but may be seen in maternal diabetes (especially uncontrolled or with large for gestational age fetus) and with fetal anomalies (diaphragmatic hernia, intestinal obstruction).

Maternal risks:

- cesarean delivery
- post-partum hemorrhage

Fetal risks:

- malpresentation
- neonatal intensive care unit admission
- need for surgical intervention if anomalies present
- neonatal hypoglycemia
- stillbirth and neonatal death

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

Shanks, Anthony, et al. "Assessing the optimal definition of oligohydramnios associated with adverse neonatal outcomes." *Journal of Ultrasound in Medicine* 30.3 (2011): 303-307.

Magann EF, Sandlin AT, Ounpraseuth ST. Amniotic fluid and the clinical relevance of the sonographically estimated amniotic fluid volume: oligohydramnios. *J Ultrasound Med* 2011;30:1573-85.

Moore, Thomas R. "Abnormal Amniotic Fluid Volume." *Protocols for High-Risk Pregnancies* (2010): 399.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

21. PLATELET COUNT LESS THAN 120,000

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Platelet count less than 120,000

Platelet disorders in pregnancy include those that are time-limited to pregnancy (gestational thrombocytopenia, HELLP syndrome) and those that may pre-date or be newly diagnosed during the pregnancy (idiopathic thrombocytopenic purpura (ITP), thrombotic thrombocytopenic purpura (TTP)). With the exception of gestational thrombocytopenia, all of these platelet disorders place the mother at increased risk for blood loss and need for transfusion.

Gestational thrombocytopenia: occurs in 7-8% of pregnancies and accounts for 70-80% of cases of thrombocytopenia in pregnancy, typically diagnosed in the third trimester, rarely associated with platelet counts below 70,000, not associated with increased risks of bleeding in the mother or fetus, platelet counts return to normal after delivery.

It is important to differentiate gestational thrombocytopenia from more serious platelet disorders:

- ITP: chronic disorder associated with:
 - fluctuating platelet counts that may be lower than 50,000
 - need for steroid or immune globulin treatment and platelet transfusion to avoid excess blood loss at delivery, particularly surgical delivery.
- TTP: acute or chronic disorder generally associated with:
 - severe thrombocytopenia of 20,000 or less
 - hepatic impairment
 - renal impairment
 - CNS impairment
 - increased risk of death for both mother and fetus.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- HELLP syndrome: an acute condition occurring in up to 2% of pregnancies, usually seen in the setting of preeclampsia, and characterized by:
 - thrombocytopenia
 - elevated liver enzymes
 - hemolytic anemia
 - potential for severe maternal illness including:
 - liver failure
 - hepatic subcapsular hematoma
 - excess maternal blood loss
 - seizure
 - maternal death
 - preterm birth
 - intrauterine growth restriction
 - fetal death

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

Gernsheimer T, James AH, Stasi R. How I treat thrombocytopenia in pregnancy. Blood 2013;121:38-47.
Thrombocytopenia during pregnancy. Importance, diagnosis and management. Boehlen F. Hamostaseologie. 2006 Jan;26(1):72-4

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

22. POSITION PRESENTATION OTHER THAN VERTEX AT TERM OR WHILE IN LABOR

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Presentation Risks

Non-vertex presentations occur in less than 4% of all pregnancies. This would include breech, brow, face, transverse lie, and compound presentations. Non-vertex presentations are associated with congenital abnormalities of the baby, multiple pregnancies, placenta previa, and uterine abnormalities. These associations would increase risk to the mother/baby in addition to the actual risks associated with non-vertex delivery.

C-section has become the standard mode of delivery for babies in non-vertex positions. Physicians and midwives may not have adequate training in the vaginal delivery of non-vertex presentations further increasing the risk of injury or death to both mother and baby. A transverse presentation is considered incompatible with vaginal delivery. Posterior, Brow, and Face presentations are associated with complicated delivery and increased maternal and/or fetal complications and may require C-section if the fetal position cannot be rotated.

Disclosure of risks related to: Position presentation other than vertex at term or while in labor:

Risks to Babies:

- Low APGAR scores
- Ruptured organs (kidney, liver)
- Neck Trauma
- Genital edema
- Prematurity
- Cord Prolapse
- Respiratory distress
- Stillbirth

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- Head entrapment
- Edema to face and skull
- Tracheal damage
- Increased NICU admission rates
- Shoulder/arm trauma
- Hip and leg trauma
- Intracranial hemorrhage
- Death

Maternal Risks:

- C-section
- Prolonged/Dysfunctional labor
- Placenta abruption
- Increased risk of deep lacerations

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

de Leeuw, J. (2002). Mortality and early morbidity for abdominal and vaginal deliveries in breech presentation. *Journal of Obstetrics and Gynaecology* , 22 (2), 127-139.

Tidy, C. R. (2010). *patient.co.uk/doctor/malpresentations*. Retrieved from patient.co.uk.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

23. PRE-ECLAMPSIA/ECLAMPSIA

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to Pre-eclampsia:

Pre-eclampsia is a leading cause of death in pregnant women and occurs in 5% of all pregnancies. The management of pre-eclampsia may require medication and monitoring unavailable in an out of hospital setting.

Maternal Risks:

- Hypertension leading to brain injury
- Liver Failure
- Kidney Failure
- HELLP syndrome
- Clotting problems (DIC)
- Pulmonary edema
- Seizure (Eclampsia)
- Stroke
- Placental Abruption
- C-section
- Death

Fetal Risks:

- Small for gestational age (IUGR)
- Premature Birth
- Stillbirth

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

American College of Obstetricians and Gynecologists. (2011). *Frequently Asked Questions: Pregnancy: High Blood Pressure During Pregnancy*. ACOG.
Cunningham, C. L. (2010). *Williams Obstetrics* (23rd Edition ed.). New York, NY: McGraw-Hill.
Frye, A. (1998). *Holistic Midwifery* (Vol. 1). Portland, OR: Labry's Press.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

24. PREGNANCY LASTING LONGER THAN 42 COMPLETED WEEKS WITH AN ABNORMAL STRESS TEST

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the North American Registry of Midwives).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Pregnancy is considered to be postdates at 42 weeks of gestation. There is limited research available to outline the risks of a pregnancy continuing beyond 42 weeks **with** an abnormal stress test. Current medical standard of practice is that beginning at 41 weeks, a non-stress test (NST) be combined with other indicators of fetal well-being, i.e., amniotic fluid index (AFI) or biophysical profile (BPP). There is no benefit in continuing a pregnancy at or post term after fetal surveillance is found to be non-reassuring. The recommendation is delivery. (Price, 2014)

Maternal Risks:

- Oligohydramnios
- Medical induction
- C-section
- Prolonged labor
- Complicated delivery such as: Shoulder dystocia

Fetal Risk

- Large size leading to risks associated with macrosomia
- uteroplacental insufficiency
- Asphyxia
- Infection
- Neonatal acidemia

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- Low Apgar
- Birth Injury
- Stillbirth
- Postmaturity/Dysmaturity syndrome
- Fetal distress
- Meconium Aspiration
- Death

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

Hilder, C. T. (1998). Prolonged Pregnancy: evaluating gestation-specific risks of fetal and infant mortality. *BJOG* .
 O'Neill, E. T. (2012). Antepartum evaluation of the fetus and fetal well-being. *Clinical Obstetrics and Gynecology* , 55 (3), 722.
 Preboth, M. (2000). Practice Guidelines ACOG Guidelines on Antepartum Fetal Surveillance . *Am Fam Physician* .
 Price, A. (2014, January). MSN CNM. Assistant Clinical Professor VCUMC. (B. Sheets, Interviewer)
 Singh, T. (2008). The prediction of intra-partum fetal compromise in prolonged pregnancy. *Journal of Obstetrics and Gynecology* , 28 (8), 779-782.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

25. VBAC (VAGINAL BIRTH AFTER CESARIAN) PREVIOUS UTERINE INCISION OR MYOMECTOMY (8)

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Because the uterine scar for most caesarian sections is low on the uterus, women who undergo TOLAC (trial of labor after cesarean), are able to give birth vaginally 60–80% of the time. But if problems arise during TOLAC, the baby may need to be born by emergency cesarean delivery. Because uterine rupture can be sudden and unexpected labor outside of a hospital can delay delivery and increase the risk of injury and death for both mother and baby in an emergency. Some surgery for fibroids can result in a similar risk for uterine rupture. An unknown type of prior uterine scar is a contraindication for TOLAC so review of prior surgical records is essential part of the evaluation.

RISKS

Maternal risks

- Maternal hemorrhage
- Infection
- Thromboembolism
- Placenta accreta
- Death
- Emergency hysterectomy

Fetal risks

- Hypoxic Ischemic Encephalopathy
- Stillbirth

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- Perinatal death
- Neonatal death
- Respiratory morbidity
- Transient tachypnea
- Hyperbillirubinemia

The probability that a woman attempting TOLAC will achieve VBAC depends on her individual combination of factors.

Selected Clinical Factors Associated with Trial of Labor after Previous Cesarean Delivery Success

Increased Probability of Success

- Prior vaginal birth
- Spontaneous labor

Decreased Probability of Success

- Recurrent indication for initial cesarean delivery (labor dystocia)
- Increased maternal age
- Non-white ethnicity
- Gestational age greater than 40 weeks
- Maternal obesity
- Preeclampsia
- Short interpregnancy interval
- Increased neonatal birth weight

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

(8) <http://www.webmd.com/baby/tc/vaginal-birth-after-cesarean-vbac-risks-of-vbac-and-cesarean-deliveries>

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

26. PSYCHIATRIC DISORDERS (MENTAL HEALTH DISORDERS)

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Clients with mental health issues such as:

- Depression
- Panic/anxiety
- Obsessive-compulsive traits
- Schizophrenia

should be counseled about the stresses of pregnancy and the postpartum period. Clients who are taking psychiatric medication should be made aware that some potential for birth defects may exist and are advised to discuss the risks and benefits of continuing their drugs during pregnancy with their provider.

Risks associated with pregnancy and psychiatric disorders include:

- Poor maternal health
- Poor outcomes for babies including poor fetal growth and development
- Maternal psychiatric medication side effects
- Increased potential for some birth defects

Clients who are taking psychiatric medication are advised to discuss the risks and benefits of continuing their drugs during pregnancy with their provider.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

Works Cited

Sinclair, C. (2004) A Midwife’s Handbook St. Louis, MO: Saunders

Vesga-Lopez O, B.C. (2008) *Psychiatric Disorders in Pregnant and Postpartum Women in the United States*, Archives of General Psychiatry, 65(7) 805-815

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

27. RUPTURE OF MEMBRANES 24 HOURS BEFORE THE ONSET OF LABOR (7)

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

The risk of prolonged rupture of membranes is chorioamnionitis. The risk increases with the delay between rupture of membranes and delivery.

MATERNAL COMPLICATIONS

- cesarean delivery
- endomyometritis
- wound infection
- pelvic abscess
- bacteremia
- postpartum hemorrhage
- postpartum hemorrhage
- bacteremia most commonly involving GBS

Rarely

- septic shock
- disseminated intravascular coagulation
- adult respiratory distress syndrome

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- maternal death

FETAL COMPLICATIONS

- fetal death
- neonatal sepsis

NEONATAL COMPLICATIONS

- perinatal death
- asphyxia
- early onset neonatal sepsis
- septic shock
- pneumonia
- intraventricular hemorrhage
- cerebral palsy

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

(7) <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3008318/>

(7) <http://www.nejm.org/doi/full/10.1056/NEJM199611143352013>

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

28. SEIZURE DISORDER REQUIRING PRESCRIPTIVE MEDICATION

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Seizure disorder requiring prescriptive medication

Most pregnancies are uneventful in women with epilepsy, and most babies are delivered healthy with no increased risk of obstetric complications in women. When controlled, there does not appear to be an increased risk for intrauterine growth restriction, preeclampsia, preterm birth or stillbirth compared to women without seizure disorder.

Fetal risks:

- With uncontrolled seizures:
 - Intrauterine growth restriction
 - Preterm birth
 - Stillbirth
- Some medications are associated with an increased risk of birth defects

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

Best practice guidelines for the management of women with epilepsy. Crawford, P., *Epilepsia*. 2005;46 Suppl 9:117-24.

McPherson JA, harper LM, Odibo AO, et al. Maternal seizure disorder and risk of adverse pregnancy outcomes. *Am J Obstet Gynecol* 2013;208:378.e1-5.

Management of epilepsy during pregnancy. Battino D., Tomson T. *Drugs*, 2007;67(18):2727-46.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

29. SEVERE LIVER DISEASE -- ACTIVE OR CHRONIC

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the **North American Registry of Midwives**).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Severe liver disease -- active or chronic

Liver disease occurs in approximately 3% of pregnancies. It may be chronic or occurring coincident with pregnancy, such as viral hepatitis or drug-induced hepatotoxicity, or pregnancy specific such as HELLP syndrome, intrahepatic cholestasis of pregnancy or acute fatty liver of pregnancy.

Severe liver disease:

- is usually acute in onset
- can be life-threatening to the mother
- associated with a high risk of stillbirth
- If hypertension has preceded the onset of HELLP syndrome, fetal growth restriction may also be present.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

Liver Disease in Pregnancy, Cleveland Clinic Disease Management Project, Jamilé Wakim-Fleming, August 10, 2010.
Joshi D, James A, Quaglia A et al. Liver Disease in Pregnancy. Lancet 2010;375:594-605.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

30. SEVERE RENAL DISEASE -- ACTIVE OR CHRONIC

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Severe Renal Disease — Active or Chronic

Renal disease is associated with increased risks of both maternal and fetal adverse outcomes. These risks, which rise with the severity of preexisting renal disease, include:

Maternal:

- Hypertension
- abruptio placentae
- deterioration of renal function including permanent end-stage renal failure;

Fetal:

- Intrauterine growth restriction
- abruptio placentae
- stillbirth

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

Williams DJ, Davison JM. Renal Disorders. In: Creasy & Resnick's Maternal-Fetal Medicine, Principles and Practice. 6th edition, 2009: Saunders Elsevier.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

31. SIGNIFICANT 2ND OR 3RD TRIMESTER BLEEDING

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the **North American Registry of Midwives**).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Significant 2nd or 3rd trimester bleeding is often associated with potentially serious conditions, including placenta previa, placenta abruption, and vasa previa.

Medical management and ultrasound is indicated to rule out and/or monitor potentially serious conditions associated with significant bleeding.

Maternal Risk Factors:

- C-section
- Hemorrhage
- Anemia
- Hypovolemic Shock
- Death
- Coagulation Defects (DIC)
- Damage to Kidneys and Brain

Fetal Risk Factors:

- Poor fetal growth (IUGR)
- Birth Defects

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- Premature Birth
- Anemia
- Hypovolemic Shock
- Stillbirth

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

American College of Obstetricians and Gynecologists. (2011). *Frequently Asked Questions in Pregnancy: Bleeding During Pregnancy*. ACOG.
 Karim, S. e. (1998). Effects of first and second trimester vaginal bleeding on pregnancy outcome.". *JPMA* .
 Nielson, E. M. (1991). The Outcome of Prengancies complicated by bleeding during the second trimester. *Surgery, Gynecology, & Obstetrics* .
 Oylese, Y. (2010). Third Trimester Bleeding. *Protocols for High Risk Pregnancies* .

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

32. SIGNIFICANT GLUCOSE INTOLERANCE (PREEXISTING DIABETES, GESTATIONAL DIABETES, PCOS)

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Significant glucose intolerance

Pre-gestational diabetes mellitus (Type 1 or Type 2) affects approximately 1% of pregnancies, with an incidence rising with the incidence of type 2 diabetes in younger adults. Gestational diabetes is diagnosed in 5-7% of pregnancies.

Risk factors for GDM: occurs more commonly in women with a family history of diabetes, prior personal history of glucose intolerance including prior gestational diabetes, obesity, and maternal age over 25.

Maternal risks:

- Hypertension
- Antepartum hospitalization
- Induction of labor
- Cesarean delivery
- Uncontrolled diabetes may result in:
 - kidney damage
 - retinopathy resulting in vision loss
 - peripheral nerve damage.

Fetal risks:

- Even when controlled, pre-gestational diabetes is associated with an increased risk of miscarriage and major congenital anomalies. This risk rises with poorer control around the time of conception.
- Throughout pregnancy, diabetes is associated with increased risks of:

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

- hypertensive disorders
- large for gestational age babies
- stillbirth
- abnormal progression of labor
- cesarean delivery
- shoulder dystocia with resultant brachial plexus injury
- Due to these risks, more frequent ultrasound examinations and antepartum testing of fetal well-being ~~are prescribed~~ may be indicated.
- In the newborn period
 - hypoglycemia
 - hyperbilirubinemia
 - polycythemia

Timing of delivery:

- Pre-gestational diabetes, and uncontrolled gestational diabetes: between 37 and 39 weeks, individualized
- Controlled gestational diabetes: between 39 and 41 weeks, individualized

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

Pre-gestational Diabetes Mellitus. American College of Obstetricians and Gynecologists, Practice Bulletin 60, March 2005.
 Gestational Diabetes Mellitus. American College of Obstetricians and Gynecologists, Practice Bulletin 137, August 2013.
 Landon MB, Gabbe SG. Gestational Diabetes Mellitus. Obstet Gynecol 2011;118:1379-93.

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

33. UNCONTROLLED HYPERTHYROIDISM

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Hyperthyroidism occurs in 0.2% of pregnancies; Graves' disease accounts for 95% of these cases.

The signs and symptoms of hyperthyroidism include nervousness, tremors, tachycardia, frequent stools, excessive sweating, heat intolerance, weight loss, goiter, insomnia, palpitations, and hypertension.

RISKS

- Premature delivery
- Severe preeclampsia
- Heart failure
- Maternal death
- Low birth weight
- Fetal death
- Abnormal thyroid function in the newborn

Thyroid storm is a medical emergency and occurs in 1% of pregnant patients with hyperthyroidism and can be triggered by infection, labor or delivery.

RISKS

- Shock
- Stupor
- Coma

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

(1)[http://www.acog.org/Resources And Publications/Practice Bulletins/Committee on Practice Bulletins --
_Obstetrics/Thyroid Disease in Pregnancy](http://www.acog.org/Resources_And_Publications/Practice_Bulletins/Committee_on_Practice_Bulletins_-_Obstetrics/Thyroid_Disease_in_Pregnancy)

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

34. UTERINE ABLATION (ENDOMETRIAL ABLATION)

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. **The risks listed below apply to birth in any setting, and are not all-inclusive.** The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to Uterine Ablation (Endometrial Ablation):

Endometrial Ablation is a procedure accompanied by sterilization or the strong recommendation for continuous contraception. Pregnancy after ablation is rare and therefore there is little research and the maternal and fetal complications are poorly defined.

Maternal Risks:

- Miscarriage
- Placenta accreta
- Manual/Surgical removal of placenta
- Hemorrhage
- Uterine rupture
- C-Section
- Hysterectomy
- Death

Fetal Risks:

- Prematurity
- Death
- Possible increase in anomalies

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

American College of Obstetricians and Gynecologists. (2013). *Frequently Asked Questions: Special Procedures: Endometrial Ablation*. ACOG.
Jenny, S. L. (2006). Pregnancy after endometrial ablation: English literature review and case report . *The Journal of Minimally Invasive Gynecology* , 13 (2), 88-91.
Laberge P. (2008, Oct). Serious and deadly complications from pregnancy after endometrial ablation reports and review of the literature. *J Gynecology Obstetrics Biological Reproduction (Paris)* .

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.

35. UTERINE ANOMALY

Preamble:

The Midwives Model of Care® recognizes the client/patient as the primary decision maker in all aspects of her care and respects her autonomy. This is supported within a model of well-informed, shared decision-making in order to achieve optimal clinical outcomes. Disclosure of risks is an integral part of the informed consent process, as outlined by NARM (the *North American Registry of Midwives*).

“If a midwife supports a client’s choices that are outside of her Plan of Care, she must be prepared to give evidence of informed consent. The midwife must also be able to document the process that led to the decision and show that the client was fully informed of the potential risks and benefits of proceeding with the new care plan. It is the responsibility of the midwife to provide evidence-based information, clinical expertise, and when appropriate, consultation or referral to other providers to aid the client in the decision making process.” – NARM

Licensed midwives are trained experts in the management of low-risk pregnancy and birth outside of the hospital. Certain conditions may present increased risk to mother and/or baby. The risks listed below apply to birth in any setting, and are not all-inclusive. The condition/risk factor listed may require medications and treatments outside of the scope of practice of Virginia Licensed Midwives and, thus may necessitate consultation with a physician, additional testing, and careful consideration for the appropriateness of birth in an out-of-hospital setting. Some conditions in pregnancy should be optimally managed and supported by a multidisciplinary team that may include midwives, obstetricians, perinatologists, family physicians, psychologists, social workers, and spiritual advisors.

Disclosure of risks related to: Uterine anomaly

Women with a uterine anomaly (uterine septum, unicornuate uterus, bicornuate uterus, uterine didelphys) are at risk for

- PTB (preterm birth)
- Fetal presentation other than vertex
- Hemorrhage
- Retained placenta
- Kidney malformation

As required by the regulations for her practice as a Virginia Licensed Midwife, my midwife has discussed this information with me and has provided me with options for consultation and referral to a physician for the risk factors she has indicated apply to me. I have decided to:

- Consult with a physician regarding my risk factors.
- Decline consultation with a physician regarding my risk factors.

Client _____

Date _____

Midwife _____

Date _____

A Work Group comprised of members of the Board of Medicine and the Advisory Board on Midwifery has developed this information to assist licensed midwives in satisfying the requirements of Code Section 54.1-2957.9(iv), which requires midwives to disclose to their patients options for consultation and referral to a physician and evidence-based information on health risks associated with the birth of a child outside of a hospital. This information does not constitute medical advice, diagnosis, opinion or treatment. Individuals should consult a qualified health care provider for advice regarding a medical condition.